

*Proceedings of the Young Researchers
Workshop on*

Positivity in Lie Groups

Edited by

Xenia Flamm

ETH Zürich, Switzerland

Arnaud Maret

Ruprecht-Karls-Universität Heidelberg, Germany

November 2022

Foreword

The workshop on Theta-positivity (hereafter abbreviated Θ -positivity) and Higher Teichmüller Theory took place in January 2022 in Heidelberg. The original initiative to host a workshop on Θ -positivity was put forward by Xenia Flamm and Mareike Pfeil in the fall of 2019. Soon after, the COVID-19 pandemic started and the event had to be postponed a couple of times. By the time it was possible to host the workshop in person, Mareike already defended her PhD thesis and had left Heidelberg. I then offered to help Xenia with the local organization in Heidelberg in replacement of Mareike.

There were twenty-five participants, mostly PhD students, who all took an active part by either giving a talk or preparing an exercise class. The event took place at the Internationales Wissenschaftsforum Heidelberg (IWH) which provided both lecture rooms and bedrooms for the participants. The location is ideal. It is quiet, with bright rooms and a large garden. On the last day of the workshop, we had the pleasure of welcoming Anna Wienhard for a Q&A session. Anna took all our questions for two hours and provided some expertise on many aspects of Θ -positivity.

Xenia and I later had the idea of writing up a set of notes about the workshop. We are extremely pleased that all the speakers accepted to type in the content of their presentations. After some editing work, we were able to compile all the contributions in a single document. We tried our best to homogenize notations. Shall there remain mistakes or typos, we would greatly appreciate these to be communicated to us.

I want to express my gratitude to Xenia and Mareike for the initial work they put in when they first attempted to organise the workshop during the pandemic. The workshop could have never taken place without the financial contributions from STRUCTURES - Cluster of Excellence Young Researcher's Convent to which we are extremely grateful.

November 2022

Arnaud Maret

*

The idea for this workshop was born in the fall 2019, when Mareike Pfeil and I were visiting MSRI for the semester program *Holomorphic Differential Equations in Mathematics and Physics*. Together with Ivo Slegers we started to learn about Lusztig's positivity. Studying together inspired us to organize a workshop to create an environment in which we can share our knowledge to learn something new. Unfortunately, both Mareike and Ivo could not attend the final workshop, but I am confident they would have enjoyed it as much as I did.

At this point I must thank Arnaud Maret who so willingly offered to help organize the workshop when Mareike could not do it anymore. Without his bold decisions the workshop would probably not have taken place at the time it did. Thank you so much, Arnaud!

I follow Arnaud's words when thanking Anna Wienhard, the IWH and its team, as well as the STRUCTURES - Cluster of Excellence Young Researcher's Convent.

Lastly I need to express my deep gratitude to all the participants of the workshop, without who it would not have been possible. Thank you for the time, work and effort you put into your preparations, presentations and now in these final notes. I hope they will provide a great reference for us and others in the future.

November 2022

Xenia Flamm

Contents

Foreword	ii
Introduction	1
I. Higher Teichmüller Theory	2
1. Introduction	3
<i>Sofia Amontova</i>	
1.1. Geometric realization	3
1.2. Algebraic realization	5
1.3. Tool to detect Teichmüller space as a special connected component in the character variety	8
1.4. Generalization	10
2. Maximal Representations	11
<i>Lisa Ricci</i>	
2.1. Hermitian Lie groups: two examples	11
2.2. Toledo number: definition and examples	12
2.3. Maximal representations	13
2.4. Relation to Theta-positivity	15
3. Hitchin Representations	16
<i>Thomas Le Floch</i>	
3.1. Definition	16
3.2. Positivity	19
4. Exercises	21
<i>Marta Magnani, Arnaud Maret, Enrico Trebeschi</i>	
II. Background in Lie Theory	25
5. Lie Groups, Lie Algebras and Their Symmetric Spaces	26
<i>Luca De Rosa, Victor Jaeck</i>	
5.1. Lie groups and Lie algebras	26
5.2. Left invariant vector fields and the Lie algebra of a Lie group	28
5.3. The exponential map and the adjoint representation	29
5.4. Symmetric spaces and the Cartan decomposition	30
6. Restricted Root Systems and Parabolic Subgroups	32
<i>Max Riestenberg</i>	
6.1. Constructing the associated symmetric space	32
6.2. Maximal flats	33
6.3. Restricted root space decomposition	33

6.4. Weyl group and Weyl chambers	35
6.5. The KAK decomposition	35
6.6. The visual boundary	36
6.7. Flag manifolds and parabolic subgroups	37
6.8. Levi subgroups	38
6.9. Parallel sets	38
7. Representations and Dynkin Diagrams	39
<i>Daniel Soskin</i>	
7.1. Representations	39
7.2. Dynkin diagrams and related classifications	41
8. The Indefinite Special Orthogonal Group	43
<i>Jacques Audibert</i>	
8.1. The Lie group and its Lie algebra	43
8.2. The restricted root system	45
9. Exercises - Part 1	47
<i>Alex Moriani</i>	
9.1. Closed linear groups	47
9.2. Exercises	48
10. Exercises - Part 2	51
<i>Colin Davalo</i>	
10.1. Hermitian symmetric spaces, in general	51
10.2. An interesting example	51
III. Known Notions of Positivity	54
11. Totally Positive Matrices	55
<i>Romeo Troubat</i>	
11.1. Positivity for the special linear group	55
11.2. Totally positive matrices	56
12. Lusztig's Total Positivity	59
<i>Xian Dai</i>	
12.1. Some motivation for total positivity	59
12.2. Lusztig's total positivity for matrices	60
12.3. Lusztig's total positivity for split real Lie groups	60
13. Positivity of Lie Groups of Hermitian Type	63
<i>Jingyi Xue</i>	
13.1. Examples of Hermitian type Lie groups	63
13.2. Maslov index and a semigroup	70
14. Positivity of Triples of Flags	76
<i>Raphael Appenzeller, Francesco Fournier-Facio</i>	
14.1. Flags	76
14.2. Positivity of triples of flags via total positivity of groups	79
14.3. Positivity via triple ratios	81
14.4. Equivalence of the two definitions	83
14.5. Generalizations	84

Contents

14.6. The flag variety for the symplectic group	85
15. The Maslov Index	87
<i>Samuel Bronstein, Ilia Smilga</i>	
15.1. Maslov index in the symplectic group	87
15.2. Maslov index in Hermitian Lie groups of tube type	89
IV. Theta-Positivity	92
16. Introduction to Theta-Positivity	93
<i>Clarence Kineider</i>	
17. The Classification Theorem	97
<i>Pierre-Louis Blayac, Balthazar Fléchettes</i>	
17.1. A theorem by Yves Benoist	98
17.2. Proof of the classification theorem	99
18. Analysis of Groups With a Theta-Positive Structure	102
<i>Dani Kaufman</i>	
18.1. Examples of Theta-positive structures	102
18.2. The Theta-Weyl group	102
18.3. A Theta-positive semigroup	103
18.4. Theta-positive triples of flags	104
19. Positivity in Higher Teichmüller Theory	105
<i>Fernando Camacho Cadena</i>	
19.1. Diamonds	106
19.2. Metrics on diamonds	109
19.3. Basics of Anosov representations	111
19.4. Proof sketch of Theorem 19.4	112
20. Exercises	114
<i>Merik Niemeyer</i>	
20.1. The indefinite special orthogonal group	114
20.2. The symplectic group	117
20.3. Partial solutions	118
Bibliography	126

Introduction

These notes transcribe a workshop about the notion of total positivity and Θ -positivity and its relation to Higher Teichmüller Theory. Θ -positivity is a notion of positivity in semisimple Lie groups and was recently introduced by Guichard and Wienhard in [GW18] as a generalization of Lusztig’s total positivity. It is believed to be the cathartic notion to classify higher Teichmüller spaces. Without doubt, substantial progress will be achieved in the near future on the study of Θ -positive structures. These notes provide an account of the state of the art as of 2021. For the latest developments and further references (some of which were not treated in the workshop), we refer to [GW22].

Working in representation theory, one often comes across the concept of “positivity”. It appears in different contexts, for example the positive reals, the order on the circle, total positivity for matrices, positivity of triples in flag varieties and the Maslov index. An especially important role is played by Lusztig’s total positivity in the context of split real Lie groups. The new notion of Θ -positivity generalizes this to other types of semisimple Lie groups and, in particular, includes the notion of positivity for Lie groups of Hermitian type.

The notes are organized into four chapters. The first two chapters are reminders of classical material. Chapter **I** is an introduction to *Higher Teichmüller Theory*: the study of connected components of discrete and faithful representations of surface groups into semisimple Lie groups. It focuses on some classical examples of higher Teichmüller spaces consisting of maximal and Hitchin representations. Chapter **II** recalls some important definitions from Lie theory, including root systems, Dynkin diagrams, and representation theory of Lie algebras. The notions of split real Lie groups and Lie groups of Hermitian type are introduced along with an in-depth study of the particular case of the Lie group $SO(p, q)$.

The last two chapters deal with the various notions of positivity in Lie groups. Chapter **III** introduces the already known concepts of positivity with a special focus on Lusztig’s positivity. This is a preparation for the definition of Θ -positivity for semisimple Lie groups which is treated in Chapter **IV**. The main motivation for studying positivity lies in its close relation to representation theory and the question of existence of higher Teichmüller spaces. The two well-known family of examples of higher Teichmüller spaces - given by maximal representations and by Hitchin representations - can in fact be characterized by their positive structures. The guiding conjecture of Guichard and Wienhard is that so-called Θ -positive representations provide a new class of higher Teichmüller spaces. The conjecture has, by now, been proven for the most part in [GLW21] and [BP21], building up on results of [BCGP⁺21] on magical \mathfrak{sl}_2 -triples.

Part I.
Higher Teichmüller Theory

Sofia Amontova
Université de Genève

Lisa Ricci
ETH Zürich

Thomas Le Fils
Sorbonne Université

Marta Magnani
Ruprecht-Karls-Universität Heidelberg

Arnaud Maret
Ruprecht-Karls-Universität Heidelberg

Enrico Trebeschi
Università di Pavia

1. Introduction

by Sofia Amontova

Higher Teichmüller theory can be roughly speaking understood as the study of representations

$$\boxed{\text{surface groups}} \longrightarrow \boxed{\text{interesting Lie groups}}.$$

The purpose of this introductory section is to present the *classical Teichmüller theory* as a prototype: we first characterize it on the one hand via its standard definition as a geometric object (see Section 1.1) and on the other hand as an algebraic object (see Section 1.2). The latter perspective allows then to detect the classical Teichmüller space as a special object in the representation-theoretical setting (see Section 1.3) which is precisely the viewpoint that motivates to define the more general notion of a higher Teichmüller space (see Section 1.4) and serves as a transition to the next section on maximal representations by Lisa Ricci.

Setting: For the entirety of this section, let Σ be a closed connected oriented surface of genus $g \geq 2$.

1.1. Geometric realization

We first remind of the geometric picture of the Teichmüller space and start with its folklore characterization:

Definition 1.1. The (*classical*) Teichmüller space $\mathcal{T}(\Sigma)$ is given by

$$\mathcal{T}(\Sigma) = \{\text{marked conformal structure } (X, f) \text{ on } \Sigma\} / \sim$$

where

- X is a Riemann surface,
- with the homotopy equivalence $f : \Sigma \rightarrow X$ as its marking,
- $(X, f) \sim (X', f')$ is an equivalence of two marked conformal structures if and only if there exists a biholomorphism $\phi : X \rightarrow X'$ so that

$$\begin{array}{ccc} & & X \\ & \nearrow f & \downarrow \phi \\ \Sigma & & \\ & \searrow f' & \\ & & X' \end{array}$$

homotopy-commutes.

Notably, the diversity of Teichmüller theory is for one due to different perspectives of the Teichmüller space $\mathcal{T}(\Sigma)$ allowing to endow it with various structures, such as complex, hyperbolic, symplectic, algebraic structures, including several Riemannian metrics, e.g.

1. Introduction

- *Weil-Petersson metric* which is a Kähler metric,
- unique Kähler-Einstein metric,
- Bergmann metric

and non-Riemannian metrics, e.g.

- *Teichmüller metric* (which coincides with the Kobayashi metric) that roughly measures the distortion between conformal structures,
- *Thurston metric* that roughly measures the distortion between hyperbolic structures,
- Carathéodory metric.

Moreover, there is a natural discrete action by the mapping class group on the Teichmüller space, interesting geodesic and horocyclic flows on its quotient Riemann moduli space, a quantization theory of its Poisson structure, to mention a few. As such Classical Teichmüller theory is a rich theory where techniques from geometry, analysis and dynamics confluence; it is an active field in both pure mathematics and theoretical physics.

Now, for our purposes, regarding $\mathcal{T}(\Sigma)$ as the parameter space of conformal structures is not the viewpoint we would like to consider but rather the analogue parameter space of hyperbolic structures known as the Fricke space:

Definition 1.2. The *Fricke space* $\mathcal{F}(\Sigma)$ is given by

$$\mathcal{F}(\Sigma) = \{\text{marked hyperbolic structure } (M, h) \text{ on } \Sigma\} / \sim$$

where

- M is a complete hyperbolic surface,
- with the orientation preserving homeomorphism $h: \Sigma \rightarrow M$ as its marking,
- $(M, h) \sim (M', h')$ is an equivalence of two marked conformal structures if and only if there exists a isometry $i: M \rightarrow M'$ so that

$$\begin{array}{ccc} & & M \\ & \nearrow h & \downarrow i \\ \Sigma & & \\ & \searrow h' & \\ & & M' \end{array}$$

isotopy-commutes.

More precisely, we can make the change of viewpoints thanks to the uniformization theorem which implies that $\mathcal{T}(\Sigma)$ can be identified with $\mathcal{F}(\Sigma)$.

Abuse of notation: From now and on the Teichmüller space $\mathcal{T}(\Sigma)$ refers to the Definition 1.2.

We shall take a closer look at complete hyperbolic structures in the next subsection.

Remark 1.3. *To get a better geometric feel for the object $\mathcal{T}(\Sigma)$ we outline the idea of the following fact*

The Teichmüller space $\mathcal{T}(\Sigma)$ is homeomorphic to \mathbb{R}^{6g-6} .

Idea of proof. Using hyperbolic geometry one can see this for instance by parametrising the Teichmüller space by *Fenchel-Nielsen coordinates*: Let (M, h) be a marked hyperbolic structure. The closed hyperbolic surface M has a collection of pairwise disjoint simple closed geodesics that decompose the surface into a union of $3g - 3$ disjoint pairs of pants¹ (see Figure 1.1).




Figure 1.1.: Pants decomposition.

We call the corresponding lengths by l_i *length parameters* and the twists that determine the gluings pairwise τ_i *twist parameters* for $1 \leq i \leq 3g - 3$. In fact these parameters determine the hyperbolic structure of M . In particular, it holds that the following map is a homeomorphism

$$\begin{aligned} \mathcal{T}(\Sigma) &\rightarrow \mathbb{R}_{>0}^{3g-3} \times \mathbb{R}^{3g-3}, \\ [(M, h)] &\mapsto (l_1, \dots, l_{3g-3}, \tau_1, \dots, \tau_{3g-3}), \end{aligned}$$

where the tuple $(l_1, \dots, l_{3g-3}, \tau_1, \dots, \tau_{3g-3})$ is called the *Fenchel-Nielsen coordinates* associated to (M, h) following the procedure above. \square

1.2. Algebraic realization

The goal of this subsection is essentially to explain the following

Idea: Any hyperbolic structure on Σ gives rise to a representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$.

This then permits to view the Teichmüller space $\mathcal{T}(\Sigma)$ as an algebraic object by identifying it with a connected component of the *character variety*

$$\chi(\Sigma, \mathrm{PSL}(2, \mathbb{R})) := \mathrm{Hom}(\pi_1(\Sigma), \mathrm{PSL}(2, \mathbb{R})) / \mathrm{PSL}(2, \mathbb{R}).$$

Ingredients to realise the idea above. We first fix a marked hyperbolic structure (M, h) on Σ ; we refer to it as merely h for short. Further,

- let s be a basepoint of M and
- let $p: \widetilde{M} \rightarrow M$ be a universal cover.

Also recall that the group of orientation-preserving isometries of the hyperbolic plane $\mathrm{Isom}^+(\mathbb{H}^2)$ can be identified with $\mathrm{PSL}(2, \mathbb{R})$.

¹A pair of pants is a compact hyperbolic surface homeomorphic to a sphere with three boundary components.

1. Introduction

Motto: Globalization. Recall that M comes equipped with the coordinate atlas $\{(U_i, \psi_i)\}_{i \in I}$. In particular, for each connected $U_i \cap U_j \neq \emptyset$ there exists a unique transition map $g_{i,j} \in \text{Isom}^+(\mathbb{H}^2)$ such that for the coordinate charts one has that

$$\psi_i|_{U_i \cap U_j} = g_{i,j} \circ \psi_j|_{U_i \cap U_j}. \quad (1.1)$$

The strategy now is to globalize the coordinate charts in terms of the universal cover $p: \widetilde{M} \rightarrow M$ in order to produce an orientation preserving isometry

$$f_h: \widetilde{M} \rightarrow \mathbb{H}^2,$$

the so-called *developing map*. This in turn will then induce a representation

$$\rho_h: \pi_1(M) \rightarrow \text{Isom}^+(\mathbb{H}^2).$$

The construction of the pair (f_h, ρ_h) can be outlined in following two steps:

- (1) **Construct f_h as a local isometry.** Recall that \widetilde{M} is the set of homotopy classes of paths starting at the basepoint s in M . Let $[\gamma] \in \widetilde{M}$ and the path $\gamma \in M$ with endpoints s and e be a choice of representative of $[\gamma]$.

We now formalize the idea of globalising the local hyperbolic structure by extending a coordinate chart containing the basepoint s along the path γ with help of transition maps $g_{i,j}$ (see Figure 1.2):

We cover γ with a finite collection of coordinate charts $\{(U_i, \psi_i)\}_{1 \leq i \leq n}$ and suppose w.l.o.g. $s \in U_1$ and $U_i \cap U_j \neq \emptyset$ is connected for any consecutive i and j . Using (1.1) we can extend $\psi_1: U_1 \cap \gamma \rightarrow \mathbb{H}^2$ to an isometry $\psi: U_1 \cup U_2 \cap \gamma \rightarrow \mathbb{H}^2$ such that

$$\psi(x) = \begin{cases} \psi_1(x), & \text{if } x \in U_1 \cap \gamma \\ g_{1,2} \circ \psi_2(x), & \text{if } x \in U_2 \cap \gamma. \end{cases}$$

We continue this procedure inductively for the remaining coordinate charts $\{(U_i, \psi_i)\}_{3 \leq i \leq n}$ until we reach the chart U_n containing the endpoint e and we set

$$\psi(e) = g_{1,2} \circ \cdots \circ g_{n-1,n} \circ \psi_n|_{U_n \cap \gamma}(e) \in \mathbb{H}^2.$$

Finally, we define the map $f_h: \widetilde{M} \rightarrow \mathbb{H}^2$ by setting

$$f_h([\gamma]) := \psi(e).$$




Figure 1.2.: Construction of a developing map f_h as a local isometry.

Fact 1.4. The map f_h depends only on the initial chart (composition by isometry) and the homotopy class of paths (look at succession of small homotopies).

We conclude that $f_h: \tilde{M} \rightarrow M$ is well-defined (in particular independent of chart-refinement) and a local orientation-preserving isometry with respect to the hyperbolic structure on \tilde{M} inherited from M .

- (2) **Obtain holonomy ρ_h via f_h .** Recall that $\pi_1(M)$ can be identified with the group of deck transformations of the universal cover $p: \tilde{M} \rightarrow M$ that acts on \mathbb{H}^2 by isometries (see exercise 1 for more details).

Now for any deck transformation $\gamma \in \pi_1(M)$ there exists a unique $g_\gamma \in \text{Isom}^+(\mathbb{H}^2)$ so that the following diagram commutes

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{f_h} & \mathbb{H}^2 \\ \downarrow \gamma & & \downarrow g_\gamma \\ \tilde{M} & \xrightarrow{f_h} & \mathbb{H}^2 \end{array}$$

That is f_h is a ρ_h -equivariant map, where ρ_h is given by

$$\begin{aligned} \rho_h: \pi_1(M) &\rightarrow \text{Isom}^+(\mathbb{H}^2), \\ \gamma &\mapsto g_\gamma. \end{aligned}$$

In fact ρ_h is a homomorphism called the *holonomy representation* of the hyperbolic structure (M, h) .

Remark 1.5.

1. Introduction

a) The pair (f_h, ρ_h) is unique up to a $\mathrm{PSL}(2, \mathbb{R})$ -action defined by

$$(f_h, \rho_h) \xrightarrow{g} (g \circ f_h, g\rho_h g^{-1}).$$

b) In accordance with our motto stated above notice that indeed

- f_h globalizes the coordinate charts,
- ρ_h globalizes the coordinate changes.

Finally, observe that completeness of M implies that $f_h: \widetilde{M} \rightarrow \mathbb{H}^2$ is in fact a *global* orientation preserving isometry.

Idea of the proof. Completeness of M implies completeness of \widetilde{M} so that $f_h: \widetilde{M} \rightarrow \mathbb{H}^2$ is a surjective covering via the path-lifting property. Since \widetilde{M} is simply connected and f_h is a homeomorphism and a local isometry, then indeed f_h is a global isometry. \square

Conversely, it is an exercise to prove that $M \cong \mathbb{H}^2/\rho_h(\pi_1(M))$.

Upshot 1.6. We have constructed the injective map

$$\begin{aligned} \mathrm{hol}: \mathcal{T}(\Sigma) &\rightarrow \chi(\Sigma, \mathrm{PSL}(2, \mathbb{R})) \\ [(M, h)] &\mapsto [\rho := \rho_h \circ h_*: \pi_1(\Sigma) \cong \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{R})]. \end{aligned}$$

The following important fact is to be checked

Fact 1.7. $\mathcal{T}(\Sigma)$ can be identified with a connected component consisting entirely of

$$\boxed{\text{faithful and discrete}} \tag{1.2}$$

representations in the representation variety $\chi(\Sigma, \mathrm{PSL}(2, \mathbb{R}))$, i.e. every holonomy is a discrete embedding into $\mathrm{PSL}(2, \mathbb{R})$.

Remark 1.8. $\mathcal{T}(\Sigma)$ and $\mathcal{T}(\overline{\Sigma})$ (Teichmüller space associated to a closed surface with reversed orientation $\overline{\Sigma}$) are the only two connected components with the special property (1.2).

1.3. Tool to detect Teichmüller space as a special connected component in the character variety

We shall introduce a tool that will later generalize to a broader setting to single out special analogues of $\mathcal{T}(\Sigma)$ in the sense of (1.2) when passing from $\mathrm{PSL}(2, \mathbb{R})$ to some more general Lie group G .

Let $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ be a representation and $p: \widetilde{\Sigma} \rightarrow \Sigma$ a universal cover.

Fact 1.9. There exists a smooth ρ -equivariant map $f: \widetilde{\Sigma} \rightarrow \mathbb{H}^2$.

Idea of the proof. We construct the associated flat $(\mathrm{PSL}(2, \mathbb{R}), \mathbb{H}^2)$ -bundle

$$E_\rho := \widetilde{\Sigma} \times \mathbb{H}^2 / \pi_1(\Sigma) \rightarrow \Sigma.$$

Since the fibres \mathbb{H}^2 are contractible, there exists a smooth section $\Sigma \rightarrow E_\rho$ which lifts to a smooth ρ -equivariant map $f: \widetilde{\Sigma} \rightarrow \widetilde{\Sigma} \times \mathbb{H}^2 \rightarrow \mathbb{H}^2$. \square

1.3. Tool to detect Teichmüller space as a special connected component in the character variety

Since f is ρ -equivariant, the volume form $\omega_{\mathbb{H}^2} \in \Omega^2(\mathbb{H}^2, \mathbb{R})^{\mathrm{PSL}(2, \mathbb{R})}$ pullbacks to $f^*\omega_{\mathbb{H}^2} \in \Omega^2(\tilde{\Sigma}, \mathbb{R})^{\pi_1(\Sigma)}$. We then may take the pushforward $\overline{f^*\omega_{\mathbb{H}^2}} := p_*f^*\omega_{\mathbb{H}^2}$ summarizing in the following diagram:

$$\begin{array}{ccc} (\tilde{\Sigma}, f^*\omega_{\mathbb{H}^2}) & \xrightarrow{f} & (\mathbb{H}^2, \omega_{\mathbb{H}^2}) \\ \downarrow p & & \\ (\Sigma, \overline{f^*\omega_{\mathbb{H}^2}}) & & \end{array}$$

We now may define our detection tool:

Definition 1.10. The *Toledo number* associated to the representation ρ is given by

$$\tau(\rho) = \frac{1}{2\pi} \int_{\Sigma} \overline{f^*\omega_{\mathbb{H}^2}} \in \mathbb{R}.$$

Remark 1.11. Notice that $\tau(\rho)$ is well-defined as any two ρ -equivariant maps are homotopic.

Now in case $\rho = \rho_h$ is a *holonomy* representation associated to a hyperbolic structure (M, h) and we rerun the procedure above, then one can check that in the Fact 1.9 we obtain a ρ_h -equivariant map f_h as the unique *developing* map (and thus an isometry) resulting from a lift of a *developing* section $\Sigma \rightarrow E_{\rho_h}$.

Remark 1.12. As a continuation of the philosophy in Remark 1.5, notice that the developing section can be obtained as a graph by globalizing the coordinate atlas.

In particular, we can take f_M as the lift of the orientation-preserving homeomorphism (marking) $h: \Sigma \rightarrow M$ and the diagram above extends to the following:

$$\begin{array}{ccc} (\tilde{\Sigma}, f^*\omega_{\mathbb{H}^2}) & \xrightarrow{f_M} & (\mathbb{H}^2, \omega_{\mathbb{H}^2}) \\ \downarrow p & & \downarrow \\ (\Sigma, \overline{f^*\omega_{\mathbb{H}^2}}) & \xrightarrow{h} & (M, \omega_{\mathbb{H}^2}), \end{array}$$

which together with Gauß-Bonnet gives that

$$\tau(\rho_M) = \frac{1}{2\pi} \int_{f(\Sigma)} \omega_{\mathbb{H}^2} = \frac{1}{2\pi} \int_M \omega_{\mathbb{H}^2} = -\frac{2\pi}{2\pi} |\chi(\Sigma)| = 2g - 2.$$

This is in fact the maximal absolute number the Toledo number can take:

Theorem 1.13 (Milnor-Wood inequality). *For any representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ we have that*

$$|\tau(\rho)| \leq 2g - 2.$$

Crucially, the following theorem by Goldman characterizes this numerical invariant as a detection tool for the special property (1.2):

Theorem 1.14 ([Gol88]). • $\tau(\rho)$ distinguishes connected components in $\chi(\Sigma, \mathrm{PSL}(2, \mathbb{R}))$ and has values in $\mathbb{Z} \cap [\chi(\Sigma), -\chi(\Sigma)]$. That is, there are $4g - 3$ components.

1. Introduction

- ρ is a holonomy representation of a hyperbolic structure if and only if $\tau(\rho) = 2g - 2$.

Upshot 1.15. $\mathcal{T}(\Sigma)$ is the connected component singled out by the maximal Toledo number (in analogy $\mathcal{T}(\bar{\Sigma})$ by the minimal Toledo number).

Remark 1.16. *The theorems above were actually proven for the so-called Euler number $e(\rho)$ of a representation, but in fact $\tau(\rho) = e(\rho)$.*

The maximality property of the Toledo invariant holds in a more general setting leading to the study of maximal representations (see next Chapter 2).

1.4. Generalization

Let G be a Lie group. This section culminates in generalizing the special property (1.2) in this more general setting:

Definition 1.17. A *higher Teichmüller space* is a subset of $\chi(\Sigma, G) := \text{Hom}(\pi_1(\Sigma), G)/G$ which is a union of connected components that consists entirely of discrete and faithful representations.

Remark 1.18. *The existence of higher Teichmüller spaces is a priori not clear. In fact unless G is locally isometric to $\text{PSL}(2, \mathbb{R})$, the set of discrete and faithful representations is only a closed set in $\text{Hom}(\pi_1(\Sigma), G)/G$. A non-example is when G is a simply-connected complex Lie group.*

2. Maximal Representations

by Lisa Ricci

2.1. Hermitian Lie groups: two examples

In the following G is a Lie group of Hermitian type. In particular, it has an associated symmetric space $X = G/K$, where K is a maximal compact subgroup, such that

- (1) X admits a G -invariant metric $\langle \cdot, \cdot \rangle$,
- (2) X has a complex manifold structure with an almost complex structure J such that
 - the metric is Hermitian: $\langle v, w \rangle = \langle J_x v, J_x w \rangle$ for all $v, w \in T_x X$,
 - J is G -invariant: $d_x L_g \circ J_x = J_{gx} \circ d_x L_g$.

Recall that an almost complex structure J assigns to each $x \in X$ an endomorphism $J_x \in \text{End}(T_x)$ such that $J_x^2 = -\text{id}$.

It follows that if one sets $\omega_X(\cdot, \cdot) := \langle J \cdot, \cdot \rangle$ one obtains the so called *Kähler form* $\omega_X \in \Omega^2(X)$, which is

- G -invariant: $L_g^* \omega = \omega$ for all $g \in G$,
- non-degenerate,
- closed.

Example 2.1. (1) $G = \text{SL}(2, \mathbb{R})$, $K = \text{O}(2)$, $X = G/K \simeq \mathbb{H}^2$. The Riemannian metric is given by $ds^2 = \frac{dx^2 + dy^2}{y^2}$ and identifying $T_x \mathbb{H}^2 \simeq \mathbb{R}^2$ the almost complex structure is

$$J_x \begin{pmatrix} v \\ w \end{pmatrix} = \begin{pmatrix} -w \\ v \end{pmatrix}. \text{ Therefore in } i \in \mathbb{H}^2$$

$$\begin{aligned} (\omega_{\mathbb{H}^2})_i \left(\begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \right) &= \left\langle J_i \begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \right\rangle_i = \begin{pmatrix} -w_1 \\ v_1 \end{pmatrix} \cdot \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \\ &= v_1 w_2 - w_1 v_2 = (dx \wedge dy) \left(\begin{pmatrix} v \\ w \end{pmatrix}, \begin{pmatrix} v \\ w \end{pmatrix} \right) \end{aligned}$$

- (2) $G = \text{Sp}(2n, \mathbb{R})$, $K = \text{Sp}(2n, \mathbb{R}) \cap \text{O}(2n, \mathbb{R}) \cong \text{U}(n)$ and the associated symmetric space is the *Siegel upper half-space*

$$\mathcal{X}_n = \{A + iB : A, B \in \text{Sym}(n, \mathbb{R}), B \gg 0\}.$$

The G -action on \mathcal{X}_n is by Möbius transformations: given $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{Sp}(2n, \mathbb{R})$ and $Z \in \mathcal{X}_n$:

$$g \cdot Z = (AZ + B)(CZ + D)^{-1}.$$

2. Maximal Representations

Under the identification $T_{A+iB}\mathcal{X}_n \simeq \{V+iW : V, W \in \text{Sym}(n, \mathbb{R})\}$ the Riemannian metric is given by

$$\langle U_1, U_2 \rangle_{iI_n} = \frac{1}{2} \text{Tr}(U_1 \overline{U_2} + \overline{U_2} U_1).$$

The almost complex structure is multiplication by i . Then the Kähler form is

$$(\omega_X)_{iI_n}(V_1 + iW_1, V_2 + iW_2) = \text{Tr}(V_1 W_2 - W_1 V_2).$$

2.2. Toledo number: definition and examples

Let Σ be a closed¹ surface of genus $g \geq 2$, G be a Lie group of Hermitian type with associated symmetric space X and let $\rho: \pi_1(\Sigma) \rightarrow G$ be a homomorphism. Let $p: \tilde{\Sigma} \rightarrow \Sigma$ be the universal cover of Σ .

Fact 2.2. There exists a smooth ρ -equivariant map $f_\rho: \tilde{\Sigma} \rightarrow X$.

The pullback $f_\rho^* \omega_X \in \Omega^2(\tilde{\Sigma})$ of the Kähler form is $\pi_1(\Sigma)$ -invariant and therefore defines a 2-form $\overline{f_\rho^* \omega_X} \in \Omega^2(\Sigma)$, which satisfies

$$p^* \overline{f_\rho^* \omega_X} = f_\rho^* \omega_X.$$

Definition 2.3. Retain the above notation. The *Toledo number* of ρ is

$$\tau(\rho) := \frac{1}{2\pi} \int_{\Sigma} \overline{f_\rho^* \omega_X}.$$

Fact 2.4. (1) The existence of a ρ -equivariant map is equivalent to the existence of a section of the bundle

$$\pi_1(\Sigma) \backslash (\tilde{\Sigma} \times X) \rightarrow \pi_1(\Sigma) \backslash \tilde{\Sigma} \cong \Sigma$$

with contractible fiber X .

(2) $\tau(\rho)$ is well-defined since any two ρ -equivariant maps $\tilde{\Sigma} \rightarrow X$ are homotopic via a ρ -equivariant homotopy (this follows from the fact that X is contractible).

(3) For all $g \in G$ it holds $\tau(g\rho g^{-1}) = \tau(\rho)$.

Example 2.5 (The holonomy representation $\pi_1(\Sigma) \rightarrow \text{PSL}(2, \mathbb{R})$). Let Σ be a closed surface of genus $g \geq 2$, $p: \tilde{\Sigma} \rightarrow \Sigma$ the universal cover, h an hyperbolic structure on Σ with associated developing map $f_h: \tilde{\Sigma} \rightarrow \mathbb{H}^2$. This is an orientation-preserving isometry, in particular $f_h \in \text{PSL}(2, \mathbb{R})$ and the holonomy representation is

$$\begin{aligned} \rho_h: \pi_1(\Sigma) &\rightarrow \text{PSL}(2, \mathbb{R}) \\ \gamma &\mapsto f_h \circ \gamma \circ f_h^{-1}, \end{aligned}$$

where $\gamma \in \text{PSL}(2, \mathbb{R})$ denotes the action of $\pi_1(\Sigma)$ on $\tilde{\Sigma}$ by deck-transformations.

We see immediately that f_h is ρ_h -equivariant: for all $\gamma \in \pi_1(\Sigma)$, $x \in \tilde{\Sigma}$ it holds $f_h(\gamma \cdot x) = \rho_h(\gamma) \cdot f_h(x)$.

¹That is compact and without boundary.

Let $\omega_\Sigma, \omega_{\tilde{\Sigma}}$ and $\omega_{\mathbb{H}^2}$ be the volume forms on $\Sigma, \tilde{\Sigma}$ and \mathbb{H}^2 , respectively. Then since local isometries preserve the volume form it holds $p^*\omega_\Sigma = \omega_{\tilde{\Sigma}}$ and $f_h^*\omega_{\mathbb{H}^2} = \omega_{\tilde{\Sigma}}$. Therefore

$$p^*\omega_\Sigma = \omega_{\tilde{\Sigma}} = f_h^*\omega_{\mathbb{H}^2}.$$

Therefore

$$\tau(\rho_h) = \frac{1}{2\pi} \int_\Sigma \overline{f_h^*\omega_{\mathbb{H}^2}} = \frac{1}{2\pi} \int_\Sigma \omega_\Sigma = \frac{2\pi}{2\pi} |\chi(\Sigma)| = 2g - 2.$$

In the second-to-last equality we used the Gauß-Bonnet theorem.

Example 2.6 (The diagonal embedding). Recall that

$$\mathrm{Sp}(2n, \mathbb{R}) = \{x \in \mathrm{SL}(2n, \mathbb{R}) : {}^t x J_{n,n} x = J_{n,n}\},$$

where $J_{n,n} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$. Consider the diagonal embedding

$$\begin{aligned} d : \mathrm{SL}(2, \mathbb{R}) &\rightarrow \mathrm{Sp}(2n, \mathbb{R}) \\ A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} &\mapsto \begin{pmatrix} aI_n & bI_n \\ cI_n & dI_n \end{pmatrix}. \end{aligned}$$

Let X be the Siegel upper half space and define $\varphi : \mathbb{H}^2 \rightarrow X, z \mapsto zI_n$. Then φ is d -equivariant:

$$d(A)\varphi(z) = (aI_n\varphi(z) + bI_n)(cI_n\varphi(z) + dI_n)^{-1} = (az + b)I_n((cz + d)I_n)^{-1} = \frac{az + b}{cz + d}I_n = \varphi(Az).$$

Let $\rho : \pi_1(\Sigma) \rightarrow \mathrm{SL}(2, \mathbb{R})$ be any holonomy with developing map $f : \tilde{\Sigma} \rightarrow \mathbb{H}^2$. We compute the Toledo number $T(d \circ \rho)$. It is easy to check that $\varphi \circ f : \tilde{\Sigma} \rightarrow X$ is $(d \circ \rho)$ -equivariant. Moreover, $\varphi^*\omega_X = n\omega_{\mathbb{H}^2}$. Indeed,

$$\begin{aligned} (\varphi^*\omega_X)_i \left(\begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \right) &= (\omega_X)_{iI_n} \left(d_i\varphi \begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, d_i\varphi \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \right) \\ &= (\omega_X)_{iI_n} \left((v_1 + iw_1)I_n, (v_2 + iw_2)I_n \right) \\ &= \mathrm{Tr}(v_1w_2I_n - w_1v_2I_n) = n(v_1w_2 - w_1v_2) \\ &= n(\omega_{\mathbb{H}^2})_i \left(\begin{pmatrix} v_1 \\ w_1 \end{pmatrix}, \begin{pmatrix} v_2 \\ w_2 \end{pmatrix} \right). \end{aligned}$$

Thus $(\varphi \circ f)^*\omega_X = f^*\varphi^*\omega_X = nf^*\omega_{\mathbb{H}^2}$ and

$$\tau(d \circ \rho) = \frac{1}{2\pi} \int_\Sigma \overline{(\varphi \circ f)^*\omega_X} = \frac{n}{2\pi} \int_\Sigma \overline{f^*\omega_{\mathbb{H}^2}} = n\tau(\rho) = n|\chi(\Sigma)|.$$

2.3. Maximal representations

The Toledo number has the following properties.

Proposition 2.7 ([BIW14, Corollary 5.7, Corollary 5.9]). *Let G be a Lie group of Hermitian type and Σ be a closed surface of genus $g \geq 2$.*

(1) *The map*

$$\begin{aligned} \tau : \mathrm{Hom}(\pi_1(\Sigma), G) &\rightarrow \mathbb{R} \\ \rho &\mapsto \tau(\rho) \end{aligned}$$

is continuous.

2. Maximal Representations

- (2) τ takes discrete values. More precisely, it takes values in $\frac{1}{n_X}\mathbb{Z}$, where $n_X \in \mathbb{N}$ depends only on G .
- (3) $|\tau(\rho)| \leq \text{rank}(X)|\chi(\Sigma)|$.

Definition 2.8. A representation $\rho: \pi_1(\Sigma) \rightarrow G$ is *maximal* if $\tau(\rho) = \text{rank}(X)|\chi(\Sigma)|$.

Since τ is continuous and takes discrete values, it is constant on connected components of $\text{Hom}(\pi_1(\Sigma), G)$ and the set of maximal representations is a union of connected components.

By Fact 2.4 (3) the Toledo number gives a well-defined continuous map on $\chi(\Sigma, G)$ and the set of maximal representations in $\chi(\Sigma, G)$ is also a union of connected components.

Definition 2.9. A *higher Teichmüller space* is a union of connected components of $\chi(\Sigma, G)$ which consists entirely of discrete and faithful representations.

For $G \neq \text{PSL}(2, \mathbb{R})$ the set of discrete and faithful representations is only a closed subset of $\chi(\Sigma, G)$, so it is not even clear that higher Teichmüller spaces exist. However, we have the following.

Proposition 2.10 ([BIW10, §4]). *Maximal representations are injective and have discrete image.*

The proof relies on the following theorem.

Theorem 2.11 ([BIW10, Theorem 8]). *Let G be a Lie group of Hermitian type with associated symmetric space X and let $\rho: \pi_1(\Sigma) \rightarrow G$ be a representation. Then ρ is maximal if and only if there exists a continuous ρ -equivariant map $\varphi: \partial\mathbb{H}^2 \rightarrow \check{\Sigma}$ which sends positively oriented triples in $\partial\mathbb{H}^2 \cong S^1$ to maximal triples in the Shilov boundary $\check{\Sigma}$ of X .*

Remark 2.12. (1) Here the action of $\pi_1(\Sigma)$ on $\partial\mathbb{H}^2$ is given by choosing an hyperbolization of $\pi_1(\Sigma)$ as a lattice in $\text{PSL}(2, \mathbb{R})$.

- (2) In the setting of G being an Hermitian Lie group, a triple of elements $(L_1, L_2, L_3) \in \check{\Sigma}^3$ is maximal if its Maslov index $M_{\check{\Sigma}}(L_1, L_2, L_3)$ takes the maximal value possible, which is $\text{rk}(X)$.

For example, when $G = \text{Sp}(2n, \mathbb{R})$ the Shilov boundary $\check{\Sigma}$ consists of the Lagrangian subspaces of \mathbb{R}^{2n} and a triple of Lagrangian subspaces (L_1, L_2, L_3) is maximal if and only if its Maslov index is equal to n .

We now sketch the proof of Proposition 2.10.

Proposition 2.13 ([BIW10, §4.4]). *Let G be a Lie group of Hermitian type and Σ a closed surface of genus $g \geq 2$. Then maximal representations $\rho: \pi_1(\Sigma) \rightarrow G$ are injective.*

Proof. Suppose by contradiction that $\ker(\rho) \neq \{1\}$. Then there exists $\gamma \in \ker(\rho)$ of infinite order and we can find an open interval $I \subset \partial\mathbb{H}^2 \cong S^1$ such that $I, \gamma I, \gamma^2 I$ are pairwise disjoint and positively oriented. Choose $I_1, I_2, I_3 \subset I$ pairwise disjoint and positively oriented.

Let $\varphi: \partial\mathbb{H}^2 \rightarrow \check{\Sigma}$ be the continuous, ρ -equivariant map which preserves positivity given by Theorem 2.11.

Let $(x, y, z) \in I_3 \times \gamma I_2 \times \gamma^2 I_1$. Since this triple is positively oriented in $\partial\mathbb{H}^2$, we have

$$M_{\check{\Sigma}}(\varphi(x), \varphi(y), \varphi(z)) = \text{rk}(X).$$

On the other hand, the triple $(x, \gamma^{-1}y, \gamma^{-2}z)$ is negatively oriented and therefore

$$M_{\check{\Sigma}}(\varphi(x), \varphi(\gamma^{-1}y), \varphi(\gamma^{-2}z)) = -\text{rk}(X).$$

By ρ -equivariance and since $\rho(\gamma) = e$:

$$\varphi(\gamma^{-1}y) = \rho(\gamma)^{-1}\varphi(y) = \varphi(y),$$

and

$$\varphi(\gamma^{-2}z) = \rho(\gamma^{-2})\varphi(z) = \varphi(z).$$

But this implies $\text{rk}(X) = -\text{rk}(X)$, which is the desired contradiction. \square

The proof of the fact that maximal representations have discrete image can be found in [BIW10], Section 4.3.

2.4. Relation to Theta-positivity

Lie groups of Hermitian type are an example of a broader class of Lie groups, namely those which admit a Θ -positive structure. If G is such a Lie group, there is a notion of Θ -positive representations $\pi_1(\Sigma) \rightarrow G$ and Theorem 2.11 tells us that maximal representations into Hermitian Lie groups are Θ -positive.

It turns out that for any simple Lie group G admitting a Θ -positive structure there exist higher Teichmüller spaces in $\chi(\Sigma, G)$.

Theorem 2.14 ([GLW21, Theorem A]). *Let G be a simple Lie group admitting a Θ -positive structure. Then there exists a higher Teichmüller space in $\chi(\Sigma, G)$.*

The theorem is a corollary of the following two facts.

Theorem 2.15 ([GLW21, Theorem B and Theorem E]). *Let G be a simple Lie group admitting a Θ -positive structure.*

(1) *There is the following inclusion of sets of representations*

$$\{\Theta\text{-positive}\} \subseteq \{\Theta\text{-Anosov}\} \subseteq \{\text{discrete and faithful representations}\}.$$

(2) *There exists a union of connected components of $\chi(\Sigma, G)$ consisting entirely of Θ -positive representations.*

It is still open whether or not the set of Θ -positive representations forms higher Teichmüller spaces.

Conjecture 2.16 ([Wie18]). *The set of Θ -positive representations is open and closed in $\chi(\Sigma, G)$.*

3. Hitchin Representations

by *Thomas Le Fils*

3.1. Definition

Let Σ be a connected closed oriented surface of genus $g \geq 2$. Recall that the character variety $\chi(\Sigma, \mathrm{PSL}(2, \mathbb{R}))$ has two connected components $\mathcal{T}(\Sigma)$ and $\mathcal{T}(\overline{\Sigma})$ consisting of discrete and faithful representations.

For every $n \geq 2$, there exists a unique irreducible representation

$$\iota_n: \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(n, \mathbb{R})$$

up to conjugacy. Namely we can take ι_n as the map that sends $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ to the matrix of the linear map f_A in the basis \mathcal{B} where f_A is the following:

$$\begin{aligned} f_A: \mathbb{R}_{n-1}[X, Y] &\longrightarrow \mathbb{R}_{n-1}[X, Y] \\ P &\longmapsto P(aX + cY, bX + dY) \end{aligned}$$

and $\mathcal{B} = (X^{n-1}, X^{n-2}Y, \dots, Y^{n-1})$.

Remark 3.1. The matrix $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$ is sent to the matrix $\begin{pmatrix} \lambda^{n-1} & 0 & 0 & 0 \\ 0 & \lambda^{n-3} & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \lambda^{1-n} \end{pmatrix}$. Therefore hyperbolic matrices are sent to diagonalizable ones with pairwise distinct eigenvalues.

Therefore the map ι_n induces $\iota_n: \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PSL}(n, \mathbb{R})$.

Example 3.2. If $n = 2$, then $\mathcal{B} = (X, Y)$. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. The map f_A sends X to $aX + cY$ and Y to $bX + dY$ hence ι_2 is the identity.

Remark 3.3. The Veronese embedding

$$\begin{aligned} V: \mathbb{RP}^1 &\longrightarrow \mathbb{RP}^{n-1} \\ [x : y] &\longmapsto [x^{n-1} : x^{n-2}y : \dots : y^{n-1}] \end{aligned}$$

satisfies for all $A \in \mathrm{SL}(n, \mathbb{R})$, for all $p \in \mathbb{RP}^1$

$$V(A \cdot p) = \iota_n(A) \cdot V(p).$$

Let us now define Fuchsian representations.

Definition 3.4. A *Fuchsian representation* is a representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ of the form $\iota_n \circ j$ where the conjugacy class of j belongs to $\mathcal{T}(\Sigma)$.

We can now define the Hitchin component.

Definition 3.5. A *Hitchin representation* is a representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ that can be deformed into a Fuchsian representation. In other words, its conjugacy class lies in the same connected component of $\chi(\Sigma, \mathrm{PSL}(n, \mathbb{R}))$ as that of a Fuchsian representation.

Let us call this component the *Hitchin component* and denote it by Hit_n . Hitchin classified the connected components of $\chi(\Sigma, \mathrm{PSL}(n, \mathbb{R}))$ in [Hit92].

Theorem 3.6 (Hitchin). *Let $n \geq 3$. The number of connected components of $\chi(\Sigma, \mathrm{PSL}(n, \mathbb{R}))$ is 3 if n is odd, 6 if n is even.*

In the same article, Hitchin showed that this component has trivial topology.

Theorem 3.7. *The Hitchin component Hit_n is homeomorphic to $\mathbb{R}^{(2g-2)(n^2-1)}$.*

Remark 3.8. *If n is even, there exists another copy $\overline{\mathrm{Hit}}_n$ of the Hitchin component. The sets Hit_n and $\overline{\mathrm{Hit}}_n$ are obtained from each other by conjugation by an element of $\mathrm{GL}(n, \mathbb{R})$ with negative determinant.*

Exercise 3.9. *Let $j: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ be such that $[j] \in \mathcal{T}(\overline{\Sigma})$. The conjugacy class of the representation $\iota_n \circ j$ is in $\overline{\mathrm{Hit}}_n$ if $n \equiv 2 \pmod{4}$ and in Hit_n otherwise.*

Hitchin components have simple geometric interpretations for small n . For example let us remark that for $n = 2$ it coincides with Teichmüller space.

Remark 3.10. *If $n = 2$ then $\iota_2 = I_2$ thus $\mathrm{Hit}_2 = \mathcal{T}(\Sigma)$.*

For $n = 3$, a theorem of Choi-Goldman [CG93, CG97] gives a geometric interpretation of the Hitchin component.

Theorem 3.11 (Choi-Goldman). *The space Hit_3 parametrizes the marked convex real projective structures on Σ . Such a structure is an identification*

$$\Sigma \cong \mathcal{O} / \rho(\pi_1(\Sigma))$$

where $\mathcal{O} \subset \mathbb{RP}^2$ is a properly convex open set and $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(3, \mathbb{R})$ is a Hitchin representation.

3. Hitchin Representations

3.1.1. Properties

Let us denote by $\text{Flag}(\mathbb{R}^n)$ the variety of full flags of \mathbb{R}^n :

$$\text{Flag}(\mathbb{R}^n) = \{(F_0, F_1, \dots, F_n) : F_i \subset F_{i+1}, \dim(F_i) = i\}.$$

We say that two flags E and F in $\text{Flag}(\mathbb{R}^n)$ are transverse if we have $E_i \oplus F_{n-i} = \mathbb{R}^n$ for all $0 \leq i \leq n$. The group $\text{GL}(n, \mathbb{R})$ acts naturally on flags and this action is transitive on pairs of transverse flags. Therefore for any pair of transverse flags (E, F) , there exists a matrix $A \in \text{GL}(n, \mathbb{R})$ such that $(A \cdot E, A \cdot F) = (E^0, F^0)$ where E_i^0 is the vector space spanned by the i -th first vectors of the canonical basis of \mathbb{R}^n and F_i^0 is the one spanned by the i -th last ones, for all $0 \leq i \leq n$.

Let us recall the following consequence of the Švarc-Milnor Lemma. Let $j: \pi_1(\Sigma) \rightarrow \text{PSL}(2, \mathbb{R})$ be such that $[j] \in \mathcal{T}(\Sigma)$. The map $\pi_1(\Sigma) \rightarrow \mathbb{H}^2$ defined by $\gamma \mapsto j(\gamma) \cdot x_0$ is a quasi-isometry and allows us to identify $\partial\pi_1(\Sigma)$ with $\partial\mathbb{H}^2 = \mathbb{RP}^1$. The action of $\gamma \in \pi_1(\Sigma)$ on $\partial\pi_1(\Sigma)$ in this identification translates into the action of $j(\gamma)$ on \mathbb{RP}^1 . Therefore the action of a non trivial $\gamma \in \pi_1(\Sigma)$ on the boundary $\partial\pi_1(\Sigma)$ has two fixed points that we will denote by γ^+ and γ^- .

Labourie showed in his seminal work [Lab06] that Hitchin components form higher Teichmüller spaces, i.e. they contain only discrete and faithful representations. Indeed Labourie showed for each Hitchin representation the existence of special equivariant limit map, a *Frenet* map.

Theorem 3.12 (Labourie). *If ρ is a Hitchin representation, then there exists a continuous map $\xi: \partial\pi_1(\Sigma) \rightarrow \text{Flag}(\mathbb{R}^d)$ that is ρ -equivariant, $\xi = (\xi^1, \dots, \xi^n)$ and, such that for all integers $n_1, \dots, n_k \geq 1$ that add up to $\sum_i n_i = m \leq d$,*

$$\bigoplus_{i=1}^k \xi^{n_i}(x_i) \xrightarrow[x_i \neq x_j]{x_i \rightarrow x} \xi^m(x)$$

The map $\xi^1: \partial\pi_1(\Sigma) \rightarrow \mathbb{P}(\mathbb{R}^d)$ is called a *hyperconvex curve*: it satisfies for any $x_1, \dots, x_d \in \partial\pi_1(\Sigma)$ that are pairwise distinct, the following sum is direct:

$$\bigoplus_{i=1}^d \xi^1(x_i) = \mathbb{R}^d.$$

Observe that the map ξ completely determines ξ^1 by the continuity property. Guichard [Gui08] showed that actually the existence of such a continuous curve that is ρ -equivariant implies that ρ is in the Hitchin component.

Theorem 3.13 (Guichard). *There exists a ρ -equivariant continuous hyperconvex map $\partial\pi_1(\Sigma) \rightarrow \mathbb{P}(\mathbb{R}^d)$ if and only if ρ is Hitchin.*

Let us now show that the existence of a ρ -equivariant Frenet map as in Labourie's theorem implies that ρ is discrete and faithful. Let us observe before that if $x \neq y$ are in $\partial\pi_1(\Sigma)$, then $\xi(x)$ is transverse to $\xi(y)$ and in particular ξ is injective.

- Let us show that ρ is faithful. Let $\gamma \in \pi_1(\Sigma) \setminus \{1\}$. Pick any point $x \in \partial\pi_1(\Sigma) \setminus \{\gamma^+, \gamma^-\}$. We have $\gamma \cdot x \neq x$ thus $\xi(\gamma \cdot x) \neq \xi(x)$. But $\xi(\gamma \cdot x) = \rho(\gamma) \cdot \xi(x)$ hence $\rho(\gamma)$ is not trivial.

- Let us show that $\rho(\gamma)$ is diagonalizable for any $\gamma \in \pi_1(\Sigma)$. Let V_i be the vector space $\xi^i(\gamma^+) \cap \xi^{n-i+1}(\gamma^-)$ for all $1 \leq i \leq n$. Observe that for all i , $\dim(V_i) = 1$ and that $\mathbb{R}^n = \bigoplus_i V_i$. Indeed since $\xi(\gamma^+)$ is transverse to $\xi(\gamma^-)$, there exists a basis f_1, \dots, f_d of \mathbb{R}^n such that $\xi^i(\gamma^+) = \langle f_1, \dots, f_i \rangle$ and $\xi^{n-i+1}(\gamma^-) = \langle f_n, \dots, f_i \rangle$ thus $V_i = \langle f_i \rangle$. Moreover $\rho(\gamma)$ preserves V_i . Indeed $\rho(\gamma) \cdot \xi^i(\gamma^+) = \xi_i(\gamma \cdot \gamma^+) = \xi_i(\gamma^+)$ and $\rho(\gamma) \cdot \xi^{n-i+1}(\gamma \cdot \gamma^-) = \xi^{n-i+1}(\gamma \cdot \gamma^-) = \xi^{n-i+1}(\gamma^-)$.
- Let us show that ρ is discrete. Let $(\gamma_k)_{k \in \mathbb{N}}$ be such that $\rho(\gamma_k) \rightarrow \pm I_n$. For all $x \in \partial\pi_1(\Sigma)$ we have $\gamma_k \cdot x \rightarrow x$. Indeed suppose $\gamma_{k_\ell} \rightarrow y$. We have $\xi(\gamma_{k_\ell} \cdot x) \rightarrow \xi(y)$ and $\xi(\gamma_{k_\ell} \cdot x) = \rho(\gamma_{k_\ell}) \cdot \xi(x) \rightarrow \xi(x)$. Therefore $\xi(x) = \xi(y)$ and $x = y$. The sequence $\gamma_k \cdot x$ has only x as an accumulation point thus it converges to x . Let us now pick three distinct points $x_1, x_2, x_3 \in \partial\pi_1(\Sigma)$. We have $\gamma_k \cdot x_i \rightarrow x_i$ for all $1 \leq i \leq 3$. The action of $\mathrm{PSL}(2, \mathbb{R})$ on \mathbb{RP}^1 is fully determined by its action on three points hence $\gamma_k \rightarrow 1$. By discreteness we have $\gamma_k = 1$ for k large enough.

Corollary 3.14. *The Hitchin component Hit_n is a higher Teichmüller space.*

We can moreover show that the eigenvalues of an element in the image of a Hitchin representation are all distinct.

Remark 3.15. *Each Hitchin representation ρ is purely hyperbolic. Namely for every $\gamma \in \pi_1(\Sigma) \setminus \{1\}$, the eigenvalues λ_i associated with V_i satisfy $|\lambda_i| > |\lambda_{i+1}|$.*

We can define concretely ξ as follows. For $\gamma \in \pi_1(\Sigma) \setminus \{1\}$ then we can define $\xi(\gamma^+)$ to be the flag associated with the ordered eigenvalues of $\rho(\gamma)$. This defines ξ on a dense set of $\partial\pi_1(\Sigma)$ and we can then extend ξ uniquely by continuity.

Let us now give examples of maps ξ^1 in particular settings.

Example 3.16. Suppose that $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ is in the Fuchsian locus: $\rho = \iota_n \circ j$, with $j: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ such that $[j] \in \mathcal{T}(\Sigma)$. The map j allows us to identify $\partial\pi_1(\Sigma)$ with $\mathbb{P}(\mathbb{R}^2) = \mathbb{RP}^1$. Then the map $\xi^1(p) = V(p)$ is a hyperconvex curve and is ρ -equivariant:

$$\xi^1(\gamma \cdot p) = V(j(\gamma) \cdot p) = \iota_n(j(\gamma)) \cdot V(p) = \rho(\gamma) \cdot \xi^1(p).$$

Example 3.17. In the case $n = 3$, the theorem of Choi-Goldman allows us to identify $\partial\pi_1(\Sigma)$ with $\partial\mathcal{O}$ and thus gives a map $\xi^1: \partial\pi_1(\Sigma) \rightarrow \mathbb{P}(\mathbb{R}^3)$ that is a ρ -equivariant hyperconvex curve.

3.2. Positivity

3.2.1. Positivity of flags

Let us define positivity for triples of flag.

Definition 3.18. A triple of flags of the form (E^0, T, F^0) with T transverse to E^0 is said to

be *positive* if $T = u_T \cdot E^0$, with $u_T = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ * & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ * & \cdots & * & 1 \end{pmatrix}$ that has all its minors positive, unless

they are zero because of its shape.

3. Hitchin Representations

A triple (E, T, F) is positive if there exists $A \in \mathrm{SL}(n, \mathbb{R})$ such that $A \cdot (E, T, F) = (E^0, A \cdot T, F^0)$ is positive.

Hitchin representations are characterized by the existence of an equivariant limit map that sends positive triples to positive triples.

Theorem 3.19 (Labourie, Guichard, Fock-Goncharov). *A representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(n, \mathbb{R})$ is Hitchin if and only if there exists a ρ -equivariant limit map $\xi: \partial\pi_1(\Sigma) \rightarrow \mathrm{Flag}(\mathbb{R}^n)$ that sends positive triples to positive triples.*

3.2.2. More general setting

One can define Hitchin representation in a more general context. Namely when G is an adjoint real split semi-simple Lie group.

Example 3.20. These conditions are met for $G = \mathrm{SL}(n, \mathbb{R}), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{SO}(n, n+1)$.

For this class of groups, there exists an embedding $\iota: \mathrm{SL}(2, \mathbb{R}) \rightarrow G$ unique up to conjugation.

Example 3.21. For the groups $G = \mathrm{SL}(n, \mathbb{R}), \mathrm{Sp}(2n, \mathbb{R}), \mathrm{SO}(n, n+1)$, the map ι is just the embedding ι_n .

We can define use this map to define Fuchsian representations as representation of the form $\iota \circ j$ where $j: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ has its conjugacy class in $\mathcal{T}(\Sigma)$. Then we can also define Hitchin representations as the deformation of those. The theorems we saw still hold.

Theorem 3.22. *The Hitchin components are homeomorphic to $\mathbb{R}^{(2g-2)\dim(G)}$. Hitchin representations are discrete and faithful.*

We can also generalize the notion of positivity to G/B where B is a Borel subgroup of G .

Theorem 3.23 (Labourie, Guichard, Fock-Goncharov). *A representation $\rho: \pi_1(\Sigma) \rightarrow G$ is Hitchin if and only if there exists a ρ -equivariant $\xi: \partial\pi_1(\Sigma) \rightarrow G/B$ that sends positive triples to positive triples.*

For more information on these generalizations we refer to [Wie18] and references therein.

4. Exercises

by Marta Magnani, Arnaud Maret, Enrico Trebeschi

We want to explore the following facts throughout this list of exercises.

- (1) The holonomies of hyperbolic structures are discrete and faithful. Any discrete and faithful representation can be realized as a hyperbolic structure.
- (2) The boundary of $\pi_1(\Sigma)$ is homeomorphic to the boundary of the hyperbolic plane.
- (3) The limit map for Teichmüller space.
- (4) The subspace of discrete and faithful representations is closed in the space of representations.

Exercise 4.1. Let (M, h) be a marked hyperbolic structure over a closed surface Σ of genus $g \geq 2$. We recall that if $p: (\tilde{M}, \tilde{x}_0) \rightarrow (M, x_0)$ is a universal cover, $\pi_1(M, x_0)$ acts discretely and faithfully on (\tilde{M}, \tilde{x}_0) by deck transformations. There exists an isometry $f_h: \tilde{M} \rightarrow \mathbb{H}^2$, which induces an isomorphism $f_h^*: \text{Isom}(\tilde{M}) \rightarrow \text{Isom}(\mathbb{H}^2)$.

The holonomy of a point $[(M, h)] \in \mathcal{T}(\Sigma)$ is the data of the representation

$$[\rho_h] \in \chi(\Sigma, \text{PSL}(2, \mathbb{R})),$$

where $[\rho_h]$ is the class of conjugacy of the following homomorphism:

$$\pi_1(\Sigma, s_0) \xrightarrow{h_*} \pi_1(M, h(s_0)) \xrightarrow{\text{Deck}} \text{Isom}^+(\tilde{M}, \tilde{x}_0) \xrightarrow{f_h^*} \text{Isom}^+(\mathbb{H}^2) = \text{PSL}(2, \mathbb{R}).$$

To prove that ρ_h is well defined one have to check that the above construction is well defined up to conjugation.

- (a) It is a fact that the isomorphism

$$\text{Deck}: \pi_1(M, x_0) \rightarrow \text{Isom}(\tilde{M}, \tilde{x}_0) < \text{Isom}(\tilde{M})$$

only depends on the basepoint x_0 and that changing basepoint change the conjugacy class of $\text{Isom}(\tilde{M}, \tilde{x}_0)$ in $\text{Isom}(\tilde{M})$. Deduce that the morphism $\pi_1(\Sigma) \rightarrow \text{Isom}(\tilde{M})$ is well defined, up to conjugation.

- (b) Describe explicitly the behaviour of f_h^* to prove that the choice of the isometry $\tilde{M} \rightarrow \mathbb{H}^2$ only changes the conjugacy class of the representation.
- (c) Let $(M, h) \sim (M', h')$ be two hyperbolic structures, i.e. $(h')^{-1}h: M' \rightarrow M$ is homotopic to an isometry. Use the homotopy lifting property and the previous point to deduce that the two deck transformation are conjugate by the lifted isometry.

4. Exercises

Let $\rho: \pi_1(\Sigma) \rightarrow \mathrm{PSL}(2, \mathbb{R})$ be a discrete and faithful representation. We want to show that this is a covering action on \mathbb{H}^2 , so that $\mathbb{H}^2/\rho(\pi_1(\Sigma))$ is a surface with fundamental group $\rho(\pi_1(\Sigma)) \cong \pi_1(\Sigma)$, hence diffeomorphic to Σ because of the classification theorem for closed surfaces, and it inherits a hyperbolic structure by the projection $p: \mathbb{H}^2 \rightarrow \mathbb{H}^2/\rho(\pi_1(\Sigma))$.

It is a fact that an action is a covering one if and only if it is free and properly discontinuous.

- (a) Prove that a discrete subgroup G of $\mathrm{Isom}(X)$ acts properly discontinuously if (X, d) is a complete metric space.

For this point use Arzelà-Ascoli Theorem, that is

Theorem 4.1 (Arzelà-Ascoli). Let K be a compact Hausdorff space and X a metric space. Then $F \subset C(K, X)$ is compact in the compact-open topology if and only if it is equicontinuous, pointwise relatively compact and closed.

We recall that F is pointwise relatively compact if $\forall x \in K$, the set $Fx = \{f(x), f \in F\}$ is relatively compact in X .

F is equicontinuous if $\forall \varepsilon > 0, \exists \delta > 0$ such that

$$d_X(f(x), f(y)) < \varepsilon, \quad \forall f \in F, \forall x, y \in K \text{ such that } d_K(x, y) < \delta.$$

Hint: in a complete metric space compact is equivalent to closed and bounded.

- (b) Deduce from the previous point that a discrete and faithful action is properly discontinuous.
- (c) Describe the stabilizer of a point in $\mathrm{Isom}^+(\mathbb{H}^2) = \mathrm{PSL}(2, \mathbb{R})$. Without loss of generality, consider the stabilizer of i in the half-plane model.
- (d) It is a fact that a group having a one-relator presentation $\langle s_i, i \in I : r \rangle$, with r cyclically reduced is with torsion if and only if the r is a proper power.

The fundamental group of a closed orientable surface of genus g can be presented as

$$\pi_1(\Sigma) = \langle a_1, b_1, \dots, a_g, b_g : [a_1, b_1] \dots [a_g, b_g] \rangle,$$

hence is without torsion. Deduce from the previous point that a discrete and faithful action has to be free.

Exercise 4.2. Step 1: The (Gromov) boundary of \mathbb{H}^2 is

$$\partial\mathbb{H}^2 := \{r: [0, \infty) \rightarrow \mathbb{H}^2 \text{ geodesic ray}\} / \text{finite distance}$$

where

$$r_1 \sim r_2 \iff \sup_t d_{\mathbb{H}^2}(r_1(t), r_2(t)) < \infty$$

Convince yourself that the Gromov boundary of \mathbb{H}^2 is homeomorphic to S^1 . A basis for the topology on $\partial\mathbb{H}^2$ is

$$\mathcal{U}_{[\alpha], c, t} = \{\beta: [0, \infty) \rightarrow \mathbb{H}^2 \text{ geod. rays} : \beta(0) = \alpha(0), B_c(\alpha(t)) \text{ intersects } \beta\}$$

Step 2: It is well known that $\pi_1(\Sigma) = \langle a_1, b_1, \dots, a_g, b_g : \prod_{i=1}^g [a_i, b_i] = 1 \rangle$. We can define the boundary $\partial\pi_1(\Sigma)$ using the same definition of Step 1, where instead of \mathbb{H}^2 we consider the Cayley graph of $\pi_1(\Sigma)$ equipped with the word metric d_S for a finite generating set S .

Step 3: We want to show that $\partial\mathbb{H}^2$ and $\partial\pi_1(\Sigma)$ are homeomorphic. The key ingredient is the following

Lemma 4.2 (Milnor-Švarc). *Let (X, d) be a geodesic space and let Γ be a group acting properly discontinuously, cocompactly and by isometries on X . Then Γ is finitely generated and for every finite generating set S and every point $x \in X$ the map*

$$\begin{aligned} f: (\Gamma, d_S) &\rightarrow (X, d) \\ \gamma &\mapsto \gamma \cdot x \end{aligned}$$

is a quasi-isometry.

In our case $X = \mathbb{H}^2$. How does the quasi-isometry extend to the boundary?

Remark 4.3. *The intuitive thing to do would be looking at the map*

$$\begin{aligned} \hat{f}: \partial\Gamma &\rightarrow \partial\mathbb{H}^2 \\ [\gamma] &\mapsto [f \circ \gamma] \end{aligned}$$

but in general quasi-isometries send geodesics to quasi-geodesics, so \hat{f} is not well defined.

The result will follow from the following

Theorem 4.4 (Stability of quasi-geodesics in hyperbolic spaces). *Let X be a hyperbolic metric space, $\gamma: [0, L] \rightarrow X$ a quasi-geodesic and $\gamma': [0, L'] \rightarrow X$ a geodesic with same starting and ending point as γ . Then $\exists \Delta \geq 0$ such that*

$$\text{Im}(\gamma') \subset B_\Delta(\text{Im} \gamma) \text{ and } \text{Im}(\gamma) \subset B_\Delta(\text{Im} \gamma')$$

(that is, one is contained in the Δ -neighbourhood of the other)

Exercise 4.3. *When $G = \text{PSL}(2, \mathbb{R})$ the Hitchin component and the space of maximal representations both agree with Teichmüller space. The existence of a nice limit map tells us something about the representation. More precisely:*

Let $\rho: \pi_1(\Sigma) \rightarrow \text{PSL}(2, \mathbb{R})$ be a representation and let $\xi: \partial\pi_1(\Sigma) \rightarrow \partial\mathbb{H}^2$ be an injective continuous ρ -equivariant map, i.e. $\xi(\gamma \cdot r) = \rho(\gamma)\xi(r)$. Then ρ is injective and discrete.

The group $\pi_1(\Sigma)$ acts on its Cayley graph by isometries. This action extends to the boundary (definition in Exercise 2): for $[r(t)]$ geodesic ray $\in \partial\pi_1(\Sigma)$ and $\gamma \in \pi_1(\Sigma)$, $\gamma \cdot [r(t)] = [\gamma \cdot r(t)]$.

Injectivity: *Since $\rho \in \text{Hom}(\pi_1(\Sigma), \text{PSL}(2, \mathbb{R}))$ it suffices to prove*

$$\gamma \in \pi_1(\Sigma), \gamma \neq 1 \Rightarrow \rho(\gamma) \neq I_2.$$

Use injectivity and ρ -equivariance of ξ to prove injectivity of ρ .

Discreteness: ρ discrete $\iff \nexists$ sequence $\rho(\alpha_n)$ accumulating on I_2 . *Use properties of ξ to show the non-existence of such a sequence.*

Exercise 4.4. *The goal of this exercise is to understand the following statement.*

The subspace of discrete and faithful representations is closed in the space of representations.

4. Exercises

Formally, let G be a Lie group and Γ be a non-cyclic, torsion-free, hyperbolic group. The fundamental groups of hyperbolic surfaces are examples of such groups Γ .

The key ingredient is the so called Margulis Lemma (also known as Margulis-Zassenhaus Lemma).

Theorem 4.5 (Margulis Lemma). *Let G be a Lie group. There exists a neighbourhood U of the identity such that, given a discrete subgroup $J \subset G$ such that $J \cap U$ generates J , then J is nilpotent.*

Recall that J being nilpotent means that the sequence $J_1 := [J, J]$, $J_{i+1} := [J_i, J]$ is eventually the trivial group.

We will apply Margulis Lemma to prove that if $\rho_n: \Gamma \rightarrow G$ is a sequence of discrete and faithful representations that converge to a representation $\rho: \Gamma \rightarrow G$, then ρ is discrete and faithful. First, assume that ρ is not faithful, i.e. there is $g \in \Gamma \setminus \{1\}$ such that $\rho(g) = e$.

- (a) Let $g^+ := \lim_{n \rightarrow +\infty} g^n$ in the boundary of Γ . What kind of subgroup is $\text{Stab}(g^+) < \Gamma$? Conclude that there exists $h \in \Gamma \setminus \text{Stab}(g^+)$.
- (b) What is $\text{Stab}(g^+) \cap \text{Stab}(h^+)$?

We consider the subgroup of Γ defined by $J := \langle g, hgh^{-1} \rangle$.

- (c) Prove that $\rho_n(J)$ is nilpotent for some n large enough and deduce that J is nilpotent.

It is a fact that a nilpotent subgroup of a torsion-free hyperbolic group is either trivial or isomorphic to \mathbb{Z} . So, we conclude that $J \cong \mathbb{Z}$.

- (d) (harder) Prove that $\langle g, h^2 \rangle \cong \mathbb{Z}^2$.

This is a contradiction because $\langle g, h^2 \rangle$ is a nilpotent subgroup of Γ that is not trivial or a copy of \mathbb{Z} . Hence, ρ is faithful.

Assume now that ρ is not discrete. So, there exists a sequence $g_n \in \Gamma \setminus \{1\}$ such that $\rho(g_n)$ converges to the identity.

- (e) Convince yourself that there exists h not contained in $\text{Stab}(g_n^+)$ for all n (but maybe finitely many).
- (f) (somewhat tricky) Deduce that $J_n := \langle g_n, h g_n h^{-1} \rangle$ is nilpotent for some n . Conclude.

Part II.
Background in Lie Theory

Luca De Rosa
ETH Zürich

Victor Jaeck
ETH Zürich

Max Riestenberg
Ruprecht-Karls-Universität Heidelberg

Daniel Soskin
The University of Notre Dame

Jacques Audibert
Sorbonne Université

Alex Moriani
Université Côte d'Azur

Colin Davalo
Ruprecht-Karls-Universität Heidelberg

5. Lie Groups, Lie Algebras and Their Symmetric Spaces

by Luca De Rosa, Victor Jaeck

5.1. Lie groups and Lie algebras

Definition 5.1. A *Lie group* G is a group endowed with the structure of smooth manifold, such that the operations of multiplication $G \times G \rightarrow G$ and inverse $G \rightarrow G$ are smooth.

Remark 5.2. In particular, a Lie group is a locally compact Hausdorff second countable topological group.

In the remaining of the workshop we will probably be mostly interested in *linear groups*, i.e. Lie groups arising as subgroups of $\mathrm{GL}(n, \mathbb{R})$, for some n .

Exercise 5.3. Let G be a connected Lie group and $U \subseteq G$ a neighborhood of the identity $e \in G$. Show that U generates G .

Examples 5.4. (1) $(\mathbb{R}, +)$ and $(\mathbb{R} \setminus \{0\}, \cdot)$.

(2) $G = \mathrm{GL}(n, \mathbb{R})$. The linear group G is an open subset of \mathbb{R}^{n^2} , and hence it inherits a smooth structure from it. The usual matrix multiplication makes it a group. Notice moreover that matrix multiplication is a polynomial function of the entries, and the inverse of a matrix is a rational function. Hence both are smooth. It follows that G is a Lie group, called the *linear group*.

(3) Non-example: Consider the space of homeomorphisms $\mathrm{Homeo}(X)$ with X , for instance, the regular tree of degree $d \geq 3$, endowed with the compact-open topology. One can show that $\mathrm{Homeo}(X)$ is not even a locally compact topological group. In fact, a neighbourhood of the identity contains maps that fix balls of X of larger and larger radius.

(4) For more interesting examples (e.g. $\mathrm{SL}(n, \mathbb{R})$) we will use the inverse function theorem, as we will see in the first exercise session.

Definition 5.5. A *Lie algebra* \mathfrak{g} is a vector space together with a skew-symmetric bilinear map

$$[\cdot, \cdot]: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$$

satisfying the Jacobi identity: for all $X, Y, Z \in \mathfrak{g}$

$$[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] = 0 \tag{5.1}$$

Definition 5.6. Let $\mathfrak{g}, \mathfrak{h}$ be Lie algebras. A *Lie algebra homomorphism* is a linear map $\varphi: \mathfrak{g} \rightarrow \mathfrak{h}$ respecting the bracket operation, i.e.

$$\varphi([X, Y]) = [\varphi(X), \varphi(Y)]$$

for all $X, Y \in \mathfrak{g}$.

A *Lie subalgebra* $\mathfrak{m} \subseteq \mathfrak{g}$ is a subspace of \mathfrak{g} such that for all $X, Y \in \mathfrak{m}$, $[X, Y] \in \mathfrak{m}$.

Definition 5.7. Let A be a commutative algebra over a field k . A *derivation* on A is an endomorphism $\delta: A \rightarrow A$ of the underlying k -vector space A which satisfies the *Leibniz rule* $\delta(ab) = \delta(a)b + a\delta(b)$ for all $a, b \in A$. We write $\text{Der}(A)$ for the set of derivations on A .

Note that smooth vector fields $\text{Vect}^\infty(M)$ on a manifold M can be identified with derivations on $\mathcal{C}^\infty(M)$. Precisely, given a point $p \in M$ and a tangent vector $X_p \in T_pM$, we can associate

$$\delta_{X_p}: \mathcal{C}^\infty(p) \rightarrow \mathbb{R}, \quad [f] \mapsto (d_p f)(X_p),$$

where $\mathcal{C}^\infty(p)$ denotes the space of germs of functions around p . It follows that vector fields correspond to endomorphisms of $\mathcal{C}^\infty(M)$.

Proposition 5.8. *Let M be a smooth manifold. The map*

$$\alpha: \text{Vect}^\infty(M) \rightarrow \text{End}(\mathcal{C}^\infty(M)), \quad X \mapsto (f \mapsto (Xf: p \mapsto (d_p f)(X)))$$

is an isomorphism onto its image which consists of derivations $\text{Der}(\mathcal{C}^\infty(M))$ of the \mathbb{R} -algebra $\mathcal{C}^\infty(M)$

Remark 5.9. *Notice that $\text{Der}(\mathcal{C}^\infty(M))$ is not a subalgebra with respect to the natural operation of composition. Consider for instance $M = \mathbb{R}$ and a derivation X such that $Xf = f'$. Then $X \circ X(f) = Xf' = f''$. Hence the element X^2 sends f to its second derivative f'' . This endomorphism of $\mathcal{C}^\infty(M)$ clearly does not satisfy the Leibniz rule $\delta(ab) = \delta(a)b + a\delta(b)$. Nevertheless there is another operation on $\text{Der}(\mathcal{C}^\infty(M))$ that, differently from the composition, gives back a derivation.*

Definition 5.10. For $X, Y \in \text{Vect}^\infty(M)$, define

$$[X, Y] := [\delta_X, \delta_Y] := \delta_X \circ \delta_Y - \delta_Y \circ \delta_X$$

where $\delta_X: \mathcal{C}^\infty(M) \rightarrow \mathcal{C}^\infty(M)$ is the derivation corresponding to X .

Exercise 5.11. *Let δ_1 and δ_2 be derivations. Show that $[\delta_1, \delta_2]$ is a derivation as well, i.e. it satisfies the Leibniz rule.*

We have shown that $\text{Vect}^\infty(M)$ endowed with the brackets operation above is a Lie algebra. It is in fact a very important and somehow prototypical example of a Lie algebra, as we will see in the following subsection.

Our next goal is to define a Lie algebra \mathfrak{g} canonically attached to a given Lie group G .

5.2. Left invariant vector fields and the Lie algebra of a Lie group

Let G be a Lie group, acting smoothly on a smooth manifold M . Given $g \in G$ we denote with L_g the *left action map*

$$L_g: M \rightarrow M, \quad m \mapsto g \cdot m.$$

The induced map at the level of vector fields is given by

$$(L_g)_*: \text{Vect}^\infty(M) \rightarrow \text{Vect}^\infty(M), \quad X \mapsto (m \mapsto (d_m L_g)(X_m)).$$

Definition 5.12. A vector field $X \in \text{Vect}^\infty(M)$ is said to be *G-invariant*, or more precisely *G-left-invariant* if for any $g \in G$, we have $(L_g)_*(X) = X$. More explicitly, $X \in \text{Vect}^\infty(M)$ is *G-invariant* if for any $g \in G$ and $m \in M$

$$(d_m L_g)(X_m) = X_{L_g(m)} = X_{gm}$$

From now on we will denote with $\text{Vect}^\infty(M)^G \subseteq \text{Vect}^\infty(M)$ the Lie subalgebra of *G-invariant* vector fields on M .

Recall that a Lie group G is also a smooth manifold, and its group structure provides a natural action of G on itself. Precisely, we now put $G = M$, and take as smooth action $G \times M \rightarrow M$ the multiplication action

$$G \times G \rightarrow G, \quad (g, h) \mapsto gh$$

The following lemma is key to the definition of a Lie algebra of a Lie group.

Lemma 5.13. *Let G be a Lie group. The map*

$$\text{Vect}^\infty(G)^G \rightarrow T_e G, \quad X \mapsto X_e$$

is an isomorphism of vector spaces.

Proof. Injectivity: Let $X_e = 0$. Then for all $g \in G$ we have $(d_e L_g)(X_e) = X_g = 0$ and hence the vector field X is identically zero.

Surjectivity: Given $v \in T_e G$, define the vector field X as follows: for any $g \in G$

$$X_g := (d_e L_g)(v)$$

Then X is left invariant by construction, and $X_e = v$. □

From now on, given $v \in T_e G$, let v^L be the corresponding left-invariant vector field constructed above.

Definition 5.14. Let G be a Lie group. The *Lie algebra* \mathfrak{g} of G is the vector space $T_e G \cong \text{Vect}^\infty(M)^G$ consisting of *left-invariant* vector fields, endowed with the brackets operation

$$[v, w] := [v^L, w^L]_e$$

for $v, w \in T_e G$.

Lie algebras are probably the central object in Lie theory. The Lie algebra \mathfrak{g} encodes in its *G*-invariance most of the structure of the Lie group G . Moreover, Lie algebras are linear objects, hence much easier to study.

5.3. The exponential map and the adjoint representation

Exercise 5.15. The Lie algebra of $G = \mathrm{GL}(n, \mathbb{R})$ is given by $\mathrm{Mat}(n, \mathbb{R})$ with bracket operation given by the commutator

$$[A, B] = AB - BA.$$

Definition 5.16. A *simple* Lie group is a connected non-abelian Lie group G which does not have nontrivial closed connected normal subgroups.

Definition 5.17. A Lie algebra \mathfrak{g} is said *semisimple* if it is direct sum of simple Lie algebras.

5.3. The exponential map and the adjoint representation

Definition 5.18. Let M be a manifold and X a vector field on M . An *integral curve* of X through $p \in M$ is a smooth curve $\gamma: (-\delta, \delta) \rightarrow M$ such that $\gamma(0) = p$ and $\gamma'(t) = X_{\gamma(t)}$.

A vector field X on M is *complete* if for every $q \in M$ the integral curve of X through q is defined on all \mathbb{R} .

The following result will motivate the definition of the exponential map.

Theorem 5.19. Let G be a Lie group. The following hold

- (1) Left invariant vector fields on G are complete;
- (2) For every $v \in T_e G$, let $\phi_v: \mathbb{R} \rightarrow G$ be the integral curve of v^L through $e \in G$. Then ϕ_v is a smooth homomorphism $\mathbb{R} \rightarrow G$. In particular for all $t_1, t_2 \in \mathbb{R}$ we have

$$\phi_v(t_1 + t_2) = \phi_v(t_1)\phi_v(t_2).$$

- (3) The flow $\Phi: \mathbb{R} \times G \rightarrow G$ of v^L is given by

$$\Phi(t, g) = g\phi_v(t).$$

We are now ready to define the exponential map.

Definition 5.20. The G be a Lie group with Lie algebra \mathfrak{g} . The *exponential map* is defined by

$$\exp_G: \mathfrak{g} \rightarrow G, \quad v \mapsto \phi_v(1).$$

Examples 5.21. If $G = \mathrm{GL}(n, \mathbb{R})$, then $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{R}) = \mathrm{Mat}(n, \mathbb{R})$ and the Lie exponential map turns out to be

$$\exp_{\mathrm{GL}(n, \mathbb{R})}: \mathrm{Mat}(n, \mathbb{R}) \rightarrow \mathrm{GL}(n, \mathbb{R}), \quad A \mapsto \sum_{n=0}^{\infty} \frac{A^n}{n!}$$

For the remaining of this subsection, let G be a Lie group with Lie algebra \mathfrak{g} . The map $\mathrm{int}(g): G \rightarrow G$ that sends x to gxg^{-1} is a smooth automorphism of G and the associated map $G \rightarrow \mathrm{Aut}(G)$, $g \mapsto \mathrm{int}(g)$ is a homomorphism.

Definition 5.22. The *adjoint representation* of G is $\mathrm{Ad} := d_e \mathrm{int}$. For every $g \in G$, $\mathrm{Ad}(g) := d_e \mathrm{int}(g)$ is an element of $\mathrm{GL}(\mathfrak{g})$ and the map $\mathrm{Ad}: G \rightarrow \mathrm{GL}(\mathfrak{g})$ is a homomorphism.

5. Lie Groups, Lie Algebras and Their Symmetric Spaces

Exercise 5.23. For every $t \in \mathbb{R}$ and $X \in \mathfrak{g}$ it holds

$$g \exp_G(tX)g^{-1} = \exp_G(t\text{Ad}(g)X).$$

Moreover, if $G = \text{GL}(n, \mathbb{R})$ then $\text{Ad}(g)X = gXg^{-1}$.

Definition 5.24. Let \mathfrak{g} be a Lie algebra. A *representation* of \mathfrak{g} into a finite-dimensional vector space V is a Lie algebra homomorphism from \mathfrak{g} to $\text{End}(V)$.

Definition 5.25. We also call *adjoint representation* the map $\text{ad}: \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$ that sends $X \mapsto [X, -]$.

Exercise 5.26. The adjoint representation ad is a representation of the Lie algebra \mathfrak{g} . Moreover, the derivative of Ad at $e \in G$ is the Lie algebra representation ad .

Definition 5.27. The *Killing form* of a Lie algebra \mathfrak{g} is the symmetric bilinear form

$$B_{\mathfrak{g}}(x, y) := \text{Tr}(\text{ad}(x)\text{ad}(y))$$

The following criterion motivates alone the relevance of the Killing form:

Remark 5.28 (Cartan's Criterion). A real Lie algebra is semisimple if and only if its Killing form is non-degenerate.

If \mathfrak{g} is a semisimple Lie algebra, then any non-degenerate symmetric invariant bilinear form on \mathfrak{g} is a scalar multiple of the Killing form.

5.4. Symmetric spaces and the Cartan decomposition

Definition 5.29 (Riemannian viewpoint). A Riemannian manifold M is *locally symmetric* if for every $p \in M$, there is a normal neighborhood $p \in U$ and an isometry $S_p: U \rightarrow U$ such that $S_p^2 = e$ and p is the only fixed point of S_p in U . Moreover, M is *globally symmetric* if each S_p can be extended to an isometry of M .

We consider now a *Lie group viewpoint* of the notion of symmetric space. We are mostly interested in the case when G is a semisimple Lie group with finite center. The following theorem states that one can construct symmetric spaces from a Lie group and its maximal compact subgroups.

Theorem 5.30. Let G be a connected semisimple Lie group with finite center, and let K be a maximal compact subgroup. Then $X = G/K$ admits an essentially unique G -invariant metric (e.g. the one induced by the Killing form). The space X endowed with this metric is a non-positively curved symmetric space, hence complete and contractible (by Cartan-Hadamard).

We can finally introduce the Cartan decomposition of a Lie algebra.

Definition 5.31. An *involution* on \mathfrak{g} is a non-trivial Lie algebra automorphism θ of \mathfrak{g} whose square is equal to the identity. Such an involution is called a *Cartan involution* on \mathfrak{g} if $-B_{\mathfrak{g}}(X, \theta(Y))$ is a positive definite bilinear form.

5.4. Symmetric spaces and the Cartan decomposition

Definition 5.32. Let G and K be as in theorem 5.30. Since $\theta^2 = \text{id}$, the Cartan involution θ is diagonalizable and has eigenvalues $+1$ and -1 . We denote with \mathfrak{k} and \mathfrak{p} the corresponding eigenspace, and hence obtain the decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$$

named the *Cartan decomposition* with respect to θ .

Theorem 5.33. *Every real semisimple Lie algebra has a Cartan involution that is unique up to inner automorphism.*

Exercise 5.34. *Show that, with the above notations, $[\mathfrak{k}, \mathfrak{k}] \subseteq \mathfrak{k}$, $[\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{k}$ and $[\mathfrak{k}, \mathfrak{p}] \subseteq \mathfrak{p}$.*

6. Restricted Root Systems and Parabolic Subgroups

by Max Riestenberg

Assumptions: We assume G is a connected, semisimple Lie group with finite center. K denotes a maximal compact subgroup. The associated symmetric space $X = G/K$ is of noncompact type.

Review: I am assuming we know about Lie groups G , Lie algebras \mathfrak{g} , symmetric spaces $X = G/K$, and the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$.

Goal: We need to cover the restricted root space decomposition, flag manifolds, and parabolic subgroups. In particular we need to give the description of parabolic subgroups in terms of restricted roots.

6.1. Constructing the associated symmetric space

I added this subsection after my talk, since there was some discussion about how to associate a symmetric space to a semisimple Lie group. I will only summarize the construction.

Suppose G is a real semisimple connected Lie group with finite center. Let K be a maximal compact subgroup. Form the quotient $X = G/K$. The orbit map $G \rightarrow X$ given by $g \mapsto gK$ has differential at the identity $\mathfrak{g} \rightarrow T_{[K]}G/K$. The kernel of this linear map is \mathfrak{k} . Define \mathfrak{p} to be the Killing form perpendicular of \mathfrak{k} . Then the differential of the orbit map restricts to an isomorphism $\mathfrak{p} \rightarrow T_{[K]}G/K$. The Killing form restricts to a positive definite inner product on \mathfrak{p} , and K acts on \mathfrak{p} via the adjoint action by isometries for this inner product. The inner product may be extended to all of $X = G/K$ by left translation by the action of G .

The resulting Riemannian manifold turns out to be a symmetric space of non-compact type. We define the Cartan involution θ on $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ by setting \mathfrak{k} to be the $+1$ -eigenspace and \mathfrak{p} to be the -1 -eigenspace. This involution integrates to a unique involution $\sigma: G \rightarrow G$. The geodesic symmetry at $[K]$ is then given by $gK \mapsto \sigma(g)K$.

Let me emphasize that a few points in my summary are not obvious (to me). In particular, it is not obvious that the restriction of B to \mathfrak{k} is negative definite and that the definition of θ yields a Cartan involution. (Recall that a *Cartan involution* is a Lie algebra involution θ such that the modified Killing form $-B(X, \theta Y)$ is an inner product.) Let me not attempt to prove this here.

Example 6.1. Our running example will be $\mathrm{SL}(n, \mathbb{R})$. Its associated symmetric space X_n can be modelled as the space of $n \times n$ real symmetric positive definite matrices of determinant 1. The action $G \times X \rightarrow X$ may be given by $g \cdot x = gx^t g$. Then X has a basepoint $p = I_n$, the identity matrix, with stabilizer $K = \mathrm{SO}(n)$, with involution $\sigma(g) = {}^t g^{-1}$ and Cartan involution $\theta(X) = -{}^t X$. The Cartan decomposition is given by $\mathfrak{k} = \mathfrak{so}(n)$ and \mathfrak{p} is the traceless symmetric matrices.

6.2. Maximal flats

A *flat* in X is a totally geodesic submanifold isometric to Euclidean space. A flat is *maximal* if it is maximal with respect to inclusion.

Example 6.2. The set of diagonal matrices with positive entries on the diagonal, multiplying to 1 is a maximal flat in X_n .

Recall that a point p in X determines the Cartan decomposition $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{k}$ and a canonical identification $\mathfrak{p} = T_p X$. Via this identification, the Riemannian exponential map corresponds to the Lie theoretic exponential map:

$$\exp_p(X) = e^X p.$$

Moreover, for a nonpositively curved symmetric space, the Riemannian exponential map $\exp_p: \mathfrak{p} \rightarrow X$ is a diffeomorphism.

Proposition 6.3.

$$\{\text{maximal flats through } p\} \leftrightarrow \{\text{maximal abelian subspaces of } \mathfrak{p}\}$$

Example 6.4. The set of traceless diagonal matrices form a maximal abelian subspace of \mathfrak{p} in $\mathfrak{sl}(n, \mathbb{R})$.

Proof. Let \mathfrak{a} be an abelian subspace of \mathfrak{p} . Via the identification $\mathfrak{p} = T_p X$, the Riemann curvature tensor at p is given by $R: \mathfrak{p} \times \mathfrak{p} \times \mathfrak{p} \rightarrow \mathfrak{p}$,

$$R(X, Y)Z = -[[X, Y]Z],$$

which clearly restricts to zero on \mathfrak{a} . Hence $\exp_p(\mathfrak{a})$ is a flat.

On the other hand, the sectional curvature of the plane spanned by $X, Y \in \mathfrak{p}$ is given by

$$\kappa(X, Y) = \frac{\langle R(X, Y)Y, X \rangle}{\langle X, X \rangle \langle Y, Y \rangle - \langle X, Y \rangle^2}$$

which reduces to $B([X, Y], [X, Y])$ if X, Y are orthonormal. If X, Y correspond to vectors tangent to a flat, this must be zero. On the other hand, the restriction of B to \mathfrak{k} is negative definite, so $[X, Y] = 0$. It follows that the preimage of a flat under \exp_p is an abelian subspace of \mathfrak{p} . \square

6.3. Restricted root space decomposition

Let $B: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ be the Killing form of \mathfrak{g} . Recall that it is a symmetric bilinear form, nondegenerate if and only if \mathfrak{g} is semisimple (We assume that G is semisimple in this section). The point $p \in X$ defines the Cartan involution $\theta: \mathfrak{g} \rightarrow \mathfrak{g}$ so that \mathfrak{k} is the $(+1)$ -eigenspace of θ and \mathfrak{p} is the (-1) -eigenspace of θ . We may define $B_p: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ by $B_p(X, Y) = -B(X, \theta(Y))$.

Lemma 6.5. B_p is an inner product. For $X \in \mathfrak{k}$, $\text{ad } X$ is skew-symmetric with respect to B_p . For $X \in \mathfrak{p}$, $\text{ad}(X)$ is symmetric with respect to B_p .

6. Restricted Root Systems and Parabolic Subgroups

Example 6.6. For $\mathrm{SL}(n, \mathbb{R})$ and $p = I_n$, $B_p(X, Y) = 2n \operatorname{tr}(X {}^t Y)$. Note that $\operatorname{tr}(X {}^t Y)$ is the entrywise dot product for matrices, also known as the Frobenius inner product.

By the spectral theorem of linear algebra, each $\operatorname{ad}(X): \mathfrak{g} \rightarrow \mathfrak{g}$ is real-diagonalizable over \mathbb{R} .

*

Here's a brief aside to prepare us for root theory. The takeaway is that roots are natural generalizations of eigenvalues and root spaces are natural generalizations of eigenvectors to the setting of a commuting family of linear transformations.

Exercise 6.7. Let f_1, \dots, f_n be a finite set of commuting, diagonalizable linear transformations of a vector space V . Show that they admit a common diagonalization, i.e. there exists a basis $\{e_i\}$ of V such that each f_i is diagonal in the basis $\{e_i\}$.

Recall that an eigenvalue of a linear transformation is a number λ so that $f v = \lambda v$. When you play with the previous exercise, you will discover that the eigenvalues of linear combinations of the f_i is the linear combination of the eigenvalues of the f_i . In other words, rather than think of an eigenvalue as a number, it is better to think of it as a gadget that eats linear combinations of f_i and spits out a number. And this operation should be linear, i.e. it should commute with taking linear combinations. In other words, an eigenvalue of a commuting family of linear transformations f_i is a linear functional on the span of f_i .

*

We apply these observations to a maximal abelian subspace \mathfrak{a} of \mathfrak{p} . \mathfrak{a} acts on \mathfrak{g} via the adjoint action, and this is a commuting family of diagonalizable linear transformations. Therefore we may consider its simultaneous eigenspace decomposition, which in this case is called the *restricted root space decomposition*:

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}_\alpha.$$

where $\mathfrak{g}_\alpha = \{X \in \mathfrak{g} : \forall A \in \mathfrak{a}, [A, X] = \alpha(A)X\}$ and $\Sigma = \Sigma(\mathfrak{g}, \mathfrak{a}) = \{\alpha \in \mathfrak{a}^* \setminus \{0\} : \mathfrak{g}_\alpha \neq 0\}$ is the set of *restricted roots*.

Example 6.8. For \mathfrak{a} as above in $\mathfrak{sl}(n, \mathbb{R})$, $\mathfrak{g}_0 = \mathfrak{a}$ and each root $\alpha: \mathfrak{a} \rightarrow \mathbb{R}$ is given by the difference of two diagonal entries. Each root space is one-dimensional, spanned by an off-diagonal elementary matrix.

Exercise 6.9. Give an example where $\mathfrak{a} \neq \mathfrak{g}_0$.

6.4. Weyl group and Weyl chambers

The restricted root space decomposition depends only on \mathfrak{p} and \mathfrak{a} . Therefore

$$N_K(\mathfrak{a}) = \{k \in K : \text{Ad}(k)(\mathfrak{a}) = \mathfrak{a}\}$$

acts on \mathfrak{a} and also the set of roots (recall that K denotes the stabilizer of \mathfrak{p} in G). The kernel of the action on \mathfrak{a} is

$$Z_K(\mathfrak{a}) = \{k \in K : \forall A \in \mathfrak{a}, \text{Ad}(k)(A) = A\}$$

and the quotient $W = N_K(\mathfrak{a})/Z_K(\mathfrak{a})$ is called the *(restricted) Weyl group*.

Each root $\alpha \in \Sigma$ has a *wall*

$$w_\alpha = \ker \alpha = \{A \in \mathfrak{a} : \alpha(A) = 0\}.$$

The components of $\mathfrak{a} \setminus \cup_{\alpha \in \Sigma} w_\alpha$ are called *(open) Euclidean Weyl chambers*. Elements of \mathfrak{a} in open Euclidean Weyl chambers are called *regular*. We often choose an open Euclidean Weyl chamber and denote it \mathfrak{a}^+ (sometimes called the *positive Weyl chamber*).

Example 6.10. For $\text{SL}(n, \mathbb{R})$, the restricted Weyl group W is isomorphic to the permutation group S_d on d symbols. It acts on \mathfrak{a} via simultaneously permuting the rows and columns.

Via the exponential map, we can talk about walls and Euclidean Weyl chambers in a maximal flat as well. It turns out that each wall of a maximal flat is an intersection of maximal flats.

Choose a regular element $X \in \mathfrak{a}^+$. There is an associated set of *positive roots* $\Sigma^+ = \{\alpha \in \Sigma : \alpha(X) > 0\}$ and a similarly defined set of negative roots. Every root is positive or negative. There is also a set of *simple roots* Δ , uniquely defined by the property that each positive root can be written as a linear combination of simple roots by nonnegative integers. Each wall bounding \mathfrak{a}^+ is the kernel of a unique simple root in Δ . In particular, $|\Delta| = \dim \mathfrak{a} = \text{rank } X$.

6.5. The KAK decomposition

Theorem 6.11. Let $\overline{\mathfrak{a}^+}$ be a closed Euclidean Weyl chamber in a maximal abelian subspace of \mathfrak{p} . Then

$$\mathfrak{p} = \bigcup_{k \in K} \text{Ad}(k)(\overline{\mathfrak{a}^+}).$$

Moreover, if $\text{Ad}(k)(\overline{\mathfrak{a}^+}) = \overline{\mathfrak{a}^+}$ then $\text{Ad}(k)$ fixes $\overline{\mathfrak{a}^+}$ pointwise.

Example 6.12. For $\text{SL}(n, \mathbb{R})$, the stabilizer of the point-chamber pair $(p, \overline{\mathfrak{a}^+})$, i.e.

$$M = \{g \in G : gp = p, \text{Ad}(g)(\overline{\mathfrak{a}^+}) = \overline{\mathfrak{a}^+}\}$$

is given by diagonal matrices with entries ± 1 , with an even number of -1 's.

Theorem 6.11 implies that G acts transitively on point-chamber pairs, but in general this action is not simply transitive.

Exercise 6.13. Give an example of a symmetric space G/K where the stabilizer of a point-chamber pair in G has positive dimension.

6. Restricted Root Systems and Parabolic Subgroups

Unfortunately, the following decomposition is also sometimes referred to as the ‘‘Cartan decomposition.’’ Fortunately, it can be unambiguously referred to as the ‘‘ KAK decomposition.’’

Corollary 6.14. *If G is a connected semisimple Lie group with finite center, K is a maximal compact and $\bar{\mathfrak{a}}^+$ is a closed Euclidean Weyl chamber, then $G = K \exp(\bar{\mathfrak{a}}^+)K$. Moreover, for any two factorizations $g = kak'$, $g = lbl'$ we have $a = b$.*

Example 6.15. Applied to $SL(n, \mathbb{R})$, this is just the *singular value decomposition*.

Proof. One first shows that any two points of X are related by a transvection. Moreover, each transvection lies in a 1-parameter subgroup corresponding to an element of \mathfrak{p} . It follows that $G = K \exp(\mathfrak{p})$. Now apply Theorem 6.11. \square

The previous corollary implies that the *Cartan projection* $\mu: G \rightarrow \bar{\mathfrak{a}}^+$ is well-defined. (It depends on the basepoint p .)

6.6. The visual boundary

In this section we introduce the visual boundary $\partial_{\text{vis}}X$ of X . There is a natural action of G on $\partial_{\text{vis}}X$. The orbits of this action are called *flag manifolds* and the point stabilizers are proper parabolic subgroups. We describe this picture in more detail, then give a description of parabolic subgroups in terms of the restricted root space decomposition.

We say that two unit-speed geodesic rays $c, c': [0, \infty) \rightarrow X$ are *asymptotic* if $\sup_t d(c(t), c'(t))$ is finite. The set of equivalence classes of asymptotic rays is called the *visual boundary* of X , denoted $\partial_{\text{vis}}X$. There is a natural topology on $X \sqcup \partial_{\text{vis}}X$ homeomorphic to a closed ball called the *visual compactification* of X . G acts on $\partial_{\text{vis}}X$ via $g[c] = [g \circ c]$.

Exercise 6.16. *Prove this is indeed an equivalence relation and that this action is well-defined.*

Since the action of G on X is transitive, we can move the basepoint of any geodesic ray to a fixed point p , and by Theorem 6.11 in the last section, we may use K to move the geodesic ray into (the exponential of) a preferred Weyl chamber $\bar{\mathfrak{a}}^+$. Moreover, there is an essentially unique way of doing this.

Definition 6.17. Let $\bar{\mathfrak{a}}^+$ be a closed Euclidean Weyl chamber. Its ideal boundary σ is called an *ideal/spherical Weyl chamber*.

Theorem 6.18. *Let σ be an ideal Weyl chamber, let K be a maximal compact, and let ξ be an ideal point. Then $G \cdot \xi = K \cdot \xi$ and*

$$|(G \cdot \xi) \cap \sigma| = |(K \cdot \xi) \cap \sigma| = 1.$$

The previous Theorem is a consequence of Theorem 6.11. An ideal Weyl chamber is a spherical simplex, and its faces are also simplices. If $g \in G$ takes a simplex τ to itself, it must fix τ pointwise. (Warning: this uses our assumption that G is connected!)

6.7. Flag manifolds and parabolic subgroups

Definition 6.19. The stabilizer $P = G_\xi = \{g \in G : g\xi = \xi\}$ of an ideal point $\xi \in \partial_{\text{vis}}X$ is called a *parabolic subgroup* of G . The homogeneous space $G/P \cong G \cdot \xi$ is called a *flag manifold*.

Example 6.20. We describe the visual boundary, flag manifolds, and parabolic subgroups for $\text{SL}(n, \mathbb{R})$. In this case, each ideal Weyl chamber σ corresponds to a full flag in \mathbb{R}^n , i.e. a chain of subspaces $0 = V_0 \subset V_1 \subset \cdots \subset V_{n-1} \subset V_n = \mathbb{R}^n$ with $\dim V_i = i$. Each simplex τ corresponds to a partial flag. Each parabolic subgroup stabilizes a partial flag. The maximal parabolics correspond to Grassmannians, and the minimal parabolic subgroups (a.k.a. Borel subgroup) stabilize a full flag, hence are conjugate to the subgroup of upper triangular matrices.

Each ideal point $\xi \in \partial_{\text{vis}}X$ lies in the interior of some simplex τ . It corresponds to an *eigenvalue-flag* pair, i.e. a partial flag (V_i) together with a vector $(\lambda_i) \in \mathbb{R}^n$, with the constraints that $\sum \lambda_i = 0$ and $\sum \lambda_i^2 = 1$ and $\lambda_j = \lambda_{j+1}$ when a subspace of dimension j is missing from the partial flag. Given a point $p \in X$, we can construct a geodesic ray from p and an eigenvalue-flag pair by using the inner product on \mathbb{R}^n corresponding to p to construct an orthonormal basis compatible with the partial flag, then using the eigenvalue data to construct a matrix $X \in \mathfrak{p}$ with the appropriate eigenvalues. See Eberlein's book [Ebe96] for more details.

We have seen that a pair of ideal points have the same stabilizers if and only if they span the same simplex of $\partial_{\text{vis}}X$. In other words, parabolic subgroups correspond to simplices of $\partial_{\text{vis}}X$.

By Theorem 6.18, we may assume (up to translating by an element of K) that our ideal point $\xi \in \partial_{\text{vis}}X$ is represented by the geodesic ray

$$c(t) = e^{tX}p$$

for $X \in \overline{\mathfrak{a}^+} \subset \mathfrak{a} \subset \mathfrak{p}$. We can describe G_ξ explicitly in terms of the restricted roots Σ . Let Δ be the simple roots corresponding to $\overline{\mathfrak{a}^+}$ and let $\Theta \subset \Delta$ be the simple roots that are positive on X . Let Σ^+ be the positive roots and Σ^- be the negative roots.

The roots which are zero on X lie in $\text{Span}(\Delta \setminus \Theta)$, so the set of roots positive on X is $\Sigma_\Theta^+ = \Sigma^+ \setminus \text{Span}(\Delta \setminus \Theta)$. We define

$$\mathfrak{u}_\Theta := \sum_{\alpha \in \Sigma_\Theta^+} \mathfrak{g}_\alpha \quad \text{and} \quad \mathfrak{u}_\Theta^{\text{opp}} := \sum_{\alpha \in \Sigma_\Theta^+} \mathfrak{g}_{-\alpha}.$$

Exercise 6.21. \mathfrak{u}_Θ and $\mathfrak{u}_\Theta^{\text{opp}}$ are nilpotent subalgebras of \mathfrak{g} .

Example 6.22. For $\text{SL}(n, \mathbb{R})$, the standard subalgebras \mathfrak{u}_Θ are strictly upper triangular.

Theorem 6.23. Let P_Θ denote the normalizer of \mathfrak{u}_Θ in G . Then $P_\Theta = G_\xi$.

We skip the proof, but mention the main idea: each group is equal to the set of elements $g \in G$ such that the limit $\lim_{t \rightarrow \infty} e^{-tX} g e^{tX}$ exists. See Eberlein's book [Ebe96] for more details.

6. Restricted Root Systems and Parabolic Subgroups

6.8. Levi subgroups

The subgroup P_Θ is called the *standard parabolic* associated to Θ . ('Standard' means standard with respect to the choices of fixed basepoint, flat, and chamber.) Let P_Θ^{opp} denote the normalizer of u_Θ^{opp} in G . The intersection $L_\Theta := P_\Theta \cap P_\Theta^{opp}$ is called the *Levi subgroup*, and its Lie algebra is given by

$$\mathfrak{l}_\Theta = \mathfrak{g}_0 \oplus \sum_{\alpha \in \text{Span}(\Delta \setminus \Theta) \cap \Sigma^+} (\mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}).$$

The Lie algebra \mathfrak{p}_Θ of P_Θ decomposes as

$$\mathfrak{p}_\Theta = \mathfrak{l}_\Theta \oplus \mathfrak{u}_\Theta.$$

Example 6.24. For $\text{SL}(d, \mathbb{R})$, a Levi subgroup is a block diagonal subgroup. It acts on the parallel set of block diagonal symmetric matrices.

Guichard-Wienhard allow the subset Θ to be empty, in which case $P_\emptyset = G$, so G itself is considered a parabolic subgroup. At the other extreme, P_Δ is a minimal parabolic subgroup, a.k.a. a *Borel subgroup*.

Remark 6.25. *Warning! The subspace \mathfrak{p} and subalgebra \mathfrak{p}_Θ have essentially nothing to do with each other. Guichard-Wienhard avoid the notation \mathfrak{p} in the Cartan decomposition by replacing it with \mathfrak{k}^\perp .*

6.9. Parallel sets

Recall that we considered an ideal point ξ in the previous section. Let $c: \mathbb{R} \rightarrow X$ be the unit-speed geodesic with $c(0) = p$ in the asymptote class of ξ . Define Θ and P_Θ as before. The Levi subgroup $L_\Theta = P_\Theta \cap P_\Theta^{opp}$ stabilizes a *parallel set*, which is the union of biinfinite geodesics parallel to c , equivalently, it is the union of maximal flats containing c . If c is regular, it lies in a unique maximal flat, $\Theta = \Delta$, $P_\Theta = P_\Delta$ is a Borel subgroup and the parallel set is a maximal flat. In general, a parallel set is a totally geodesic subspace containing p . It is a nonpositively curved symmetric space, but not necessarily of non-compact type i.e. it has a Euclidean de Rham factor.

The presence of a Euclidean factor means that \mathfrak{l}_Θ is no longer semisimple, but it is still reductive, i.e. the direct sum of its center with a semisimple subalgebra. Observe that L_Θ acts on \mathfrak{u}_Θ via the adjoint representation. This representation is crucial for the definition of Θ -positivity!

◇

Exercise 6.26. *Give an example of a symmetric space with an isometry g which fixes p and preserves $\overline{\mathfrak{a}^+}$ but does not fix $\overline{\mathfrak{a}^+}$ pointwise. (Hint: such an isometry cannot be in the identity component.)*

Exercise 6.27. *The action of L_Θ on G/P_Θ^{opp} fixes the identity coset, so L_Θ also acts on the tangent space $T_{[P_\Theta^{opp}]G/P_\Theta^{opp}}$. Show that this representation is equivalent to the adjoint action of L_Θ on \mathfrak{u}_Θ .*

7. Representations and Dynkin Diagrams

by Daniel Soskin

7.1. Representations

Let \mathfrak{g} be a complex Lie algebra and V a vector space over the field of complex numbers, unless another field is introduced explicitly.

Definition 7.1. A *representation of a Lie algebra* \mathfrak{g} is a Lie algebras homomorphism $\phi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$, where $\mathfrak{gl}(V)$ is the general linear algebra, i.e. the algebra of endomorphisms of a vector space V with product given by $[X, Y] = XY - YX$.

Definition 7.2. A vector space V , endowed with an operation $\mathfrak{g} \times V \rightarrow V$, is called a *\mathfrak{g} -module* if the following conditions are satisfied: for any $X, Y \in \mathfrak{g}$, $v, w \in V$, and $a, b \in F$ we have

- (1) $(aX + bY).v = a(X.v) + b(Y.v)$,
- (2) $X.(av + bw) = a(X.v) + b(X.w)$,
- (3) $[Y, X].v = XY.v - YX.v$.

Given a representation $\phi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$, we can view V as a \mathfrak{g} -module via the action $X.v = \phi(X)(v)$. Conversely, given a \mathfrak{g} -module V , this equation defines a representation $\phi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$.

Definition 7.3. A *homomorphism of \mathfrak{g} -modules* is a linear map $\phi: V \rightarrow W$ such that $\phi(X.v) = X.\phi(v)$.

Definition 7.4. A \mathfrak{g} -module is called *irreducible* if it has precisely two \mathfrak{g} -submodules (itself and 0).

Definition 7.5. A \mathfrak{g} -module is called *completely reducible* if it is a direct sum of irreducible \mathfrak{g} -submodules.

Lemma 7.6 (Schur's Lemma). *Let $\phi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be an irreducible representation of \mathfrak{g} . Then the only endomorphisms of V commuting with $\phi(\mathfrak{g})$ are the scalars.*

Theorem 7.7 (Weyl). *Let $\phi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be a (finite-dimensional) representation of a semisimple Lie algebra. Then ϕ is completely reducible.*

Let \mathfrak{g} be a semisimple Lie algebra, \mathfrak{h} a fixed Cartan subalgebra and Σ a root system relative to \mathfrak{h} . If V is a finite-dimensional \mathfrak{g} -module, \mathfrak{h} acts diagonally on V , and V is a direct sum of its weight spaces: $V = \bigoplus_{\lambda \in \mathfrak{h}^*} V_\lambda$, where $V_\lambda = \{v \in V | h.v = \lambda(h)v \ \forall h \in \mathfrak{h}\}$. Whenever $V_\lambda \neq 0$, it is called *weight space* and λ is called *weight*.

7. Representations and Dynkin Diagrams

Example 7.8. The Lie algebra \mathfrak{g} itself is an \mathfrak{g} -module via the adjoint representation. Then, weights are the roots $\alpha \in \Sigma$ with the weight spaces \mathfrak{g}_α from the root space decomposition along with the weight space \mathfrak{h} of 0.

Exercise 7.9. Verify that \mathfrak{g}_α maps V_λ into $V_{\lambda+\alpha}$, $\lambda \in \mathfrak{h}^*$, $\alpha \in \Sigma$.

Example 7.10. Let us consider $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ with the standard basis $X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. One can verify that $[H, X] = 2X$, $[H, Y] = -2Y$, $[X, Y] = H$, and thus for $\mathfrak{sl}(2, \mathbb{C})$ -module V it follows that if $v \in V_\lambda$, then $X.v \in V_{\lambda+2}$ and $Y.v \in V_{\lambda-2}$. Assume now that V is an irreducible finite-dimensional \mathfrak{g} -module. There exists a nonzero $v_0 \in V_\lambda$ such that $V_{\lambda+2} = 0$, and $X.v_0 = 0$. Define $v_i = \frac{1}{i!} Y^i.v_0$, ($i > -1$). Then,

- (1) $H.v_i = (\lambda - 2i)v_i$;
- (2) $Y.v_i = (i + 1)v_{i+1}$;
- (3) $X.v_i = (\lambda - i - 1)v_{i-1}$.

All nonzero v_i are linearly independent. Let m be an integer such that $v_m \neq 0$ and $v_{m'} = 0$ for $m < m'$. The subspace of V with basis v_0, v_1, \dots, v_m is a \mathfrak{g} -submodule, so it must be equal to V , since the latter is irreducible. Note, that from (3) it follows that $m = \lambda$, and $m = \dim(V) - 1$, because each weight space is of dimension one. Thus, there exist only one irreducible $\mathfrak{sl}(2, \mathbb{C})$ -module of each possible dimension.

It turns out, that for any complex semisimple Lie algebra \mathfrak{g} is built from copies of $\mathfrak{sl}(2, \mathbb{C})$. For any root $\alpha \in \Sigma$, and any $X_\alpha \in \mathfrak{g}_\alpha$, there exist $Y_{-\alpha} \in \mathfrak{g}_{-\alpha}$ such that X_α, Y_α and $[X_\alpha, Y_\alpha]$ span a subalgebra of \mathfrak{g} isomorphic to $\mathfrak{sl}(2, \mathbb{C})$.

Definition 7.11. Let \mathfrak{h}_0 be the real form of \mathfrak{h} . An element $\lambda \in \mathfrak{h}_0^*$ is *algebraically integral* if

$$\frac{2(\lambda, \alpha)}{(\alpha, \alpha)}$$

are integers for all roots α .

Exercise 7.12. The weight of any finite-dimensional representation is algebraically integral.

The *fundamental weights* w_1, \dots, w_l are defined in a way that they form basis of \mathfrak{h}_0^* such that $\frac{2(w_i, \alpha_j)}{(\alpha_j, \alpha_j)} = \delta_{ij}$, where α_i are simple roots. An element λ is algebraically integral if and only if it is an integral combination of the fundamental weights. Thus, the set of all algebraically integral weights in \mathfrak{h}_0^* form the *weight lattice* for \mathfrak{g} .

Suppose that the Lie algebra \mathfrak{g} is the Lie algebra of a Lie group G . Then, $\lambda \in \mathfrak{h}_0^*$ is *analytically integral* if for each t in \mathfrak{h} such that $\exp(t) = e$ in G we have $(\lambda, t) \in 2\pi i\mathbb{Z}$. If a representation of \mathfrak{g} arises from representation of G , then the weights of the representation are analytically integral. For semisimple G the set of analytically integral weights form a sub-lattice of the lattice of algebraically integral weights. If G is simply connected then both lattices coincide.

There is a partial order on the space of weights. We say that μ is *higher* than λ ($\mu \geq \lambda$), if $\mu - \lambda$ is expressible as a linear combination of positive roots with non-negative coefficients.

An integral element λ is called *dominant* if it is a non-negative integer combination of the fundamental weights. Note, that it is not the same as being higher than 0.

Definition 7.13. A weight λ of a representation V of \mathfrak{g} is called *highest weight* if it is higher than every other weight of V .

Theorem 7.14. (1) Every irreducible finite-dimensional representation has a highest weight.

(2) The highest weight is dominant and algebraically integral element.

(3) Two irreducible representations with the same highest weight are isomorphic.

(4) Every dominant, algebraically integral element is the highest weight of an irreducible representation.

7.2. Dynkin diagrams and related classifications

Let Σ be a root system of dimension l , with ordered basis of simple roots $\{\alpha_1, \dots, \alpha_l\}$. For two roots $\alpha, \beta \in \Sigma$, let $\langle \beta, \alpha \rangle := \frac{2(\beta, \alpha)}{(\alpha, \alpha)}$.

Definition 7.15. The root system Σ is called *irreducible* if it cannot be partitioned into the union of two orthogonal proper subsets.

Theorem 7.16. The root system of a simple Lie algebra is irreducible.

The following statement reduces the problem of characterizing semisimple Lie algebras by their root systems to the problem of characterizing simple ones by their irreducible root systems.

Theorem 7.17. Let \mathfrak{g} be a semisimple Lie algebra with \mathfrak{h} and Σ as defined before. If $\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_t$ is the decomposition of \mathfrak{g} into simple ideals, then $\mathfrak{h}_i = \mathfrak{h} \cap \mathfrak{g}_i$ is a Cartan subalgebra of \mathfrak{g}_i with relative irreducible root subsystem Σ_i of Σ in such a way that $\Sigma = \Sigma_1 \cup \dots \cup \Sigma_t$ is the decomposition of Σ into its irreducible components.

Irreducible root systems can be classified by so called *Dynkin diagrams*.

Since $\langle \beta, \alpha \rangle$ must be an integer (called *Cartan integer*), there are only a few possible values: $0, \pm 1, \pm 2, \pm 3$. Moreover, at most two distinct root lengths may occur in an irreducible system Σ , which are referred to as *long* and *short* roots. The matrix $(\langle \alpha_i, \alpha_j \rangle)_{i,j=1,\dots,l}$ is called the *Cartan matrix* of Σ . Define the *Coxeter graph* of Σ to be a graph on l vertices $\alpha_1, \dots, \alpha_l$, where the α_i is joined to α_j ($i \neq j$) by $\langle \alpha_i, \alpha_j \rangle \langle \alpha_j, \alpha_i \rangle$ edges. Whenever a double or triple edge occurs in the Coxeter graph of Σ , we add an arrow pointing to the shorter of the two roots. The resulting object is called a *Dynkin diagram*. Since for simple roots $\langle \alpha, \beta \rangle \leq 0$, all Cartan integers can be recovered from the Dynkin diagram. Note that the Coxeter graph is connected if and only if its corresponding root system Σ is irreducible.

Theorem 7.18. If Σ is an irreducible root system of rank l , its Dynkin diagram is one of the following (l vertices in each case): A_l ($l \geq 1$), B_l ($l \geq 2$), C_l ($l \geq 3$), D_l ($l \geq 4$), E_6, E_7, E_8, F_4, G_2 , see Table 7.1.

Theorem 7.19. For each Dynkin diagram of type A-G, there exists an irreducible root system having the given diagram.

7. Representations and Dynkin Diagrams




Table 7.1.: List of Dynkin diagrams

Classification of real Lie algebras differs from the case of Lie algebras over algebraically closed fields.

Definition 7.20. The *complexification* of a real Lie algebra \mathfrak{g} is obtained from \mathfrak{g} by extending the field of scalars from real to complex. Elements of $\mathfrak{g}^{\mathbb{C}}$ can be considered as pairs (u, v) , $u, v \in \mathfrak{g}$ (or $u + iv$) with the following operations:

- (1) $(u_1, v_1) + (u_2, v_2) = (u_1 + u_2, v_1 + v_2)$,
- (2) $(\alpha + i\beta)(u, v) = (\alpha u - \beta v, \alpha v + \beta u)$ for any real α, β ,
- (3) $[(u_1, v_1), (u_2, v_2)] = ([u_1, u_2] - [v_1, v_2], [v_1, u_2] + [u_1, v_2])$.

If \mathfrak{g} is a finite-dimensional real Lie algebra, its complexification is either simple or a product of a simple complex Lie algebra and its conjugate. Thus, real simple Lie algebras can be classified by the classification of complex Lie algebras and some additional information, which can be done by *Satake diagrams*, which generalize Dynkin diagrams.

8. The Indefinite Special Orthogonal Group

by Jacques Audibert

8.1. The Lie group and its Lie algebra

Denote $I_{p,q} = \begin{pmatrix} I_p & 0 \\ 0 & -I_q \end{pmatrix}$.

Theorem 8.1. For every $B \in \text{GL}(n, \mathbb{R})$ symmetric there exists $p, q \geq 0$ and $P \in \text{GL}(n, \mathbb{R})$ such that ${}^tPBP = I_{p,q}$.

For every $B \in \text{GL}(n, \mathbb{R})$ symmetric define $\text{SO}(B) = \{M \in \text{SL}(n, \mathbb{R}) : {}^tMBM = B\}$.

Remark 8.2. If ${}^tPBP = I_{p,q}$ then $\text{SO}(B) = P \text{SO}(I_{p,q}) P^{-1}$.

Example 8.3. Let $0 \leq p \leq q$. Denote by

$$Q_{p,q} = \begin{pmatrix} 0 & 0 & W_p \\ 0 & -I_{q-p} & 0 \\ {}^tW_p & 0 & 0 \end{pmatrix} \in \text{GL}(p+q, \mathbb{R})$$

where

$$W_p = \begin{pmatrix} & & & (-1)^{p-1} \\ & & \cdots & \\ & -1 & & \\ 1 & & & \end{pmatrix} \in \text{GL}(p, \mathbb{R})$$

Denote

$$P = \begin{pmatrix} \frac{(-1)^{p-1}}{\sqrt{2}} & & & & & & & & \frac{(-1)^p}{\sqrt{2}} \\ & \cdots & & & & & & & \cdots \\ & & -\frac{1}{\sqrt{2}} & & & & & & \frac{1}{\sqrt{2}} \\ & & & \frac{1}{\sqrt{2}} & & & -\frac{1}{\sqrt{2}} & & \\ & & & & -I_{q-p} & & & & \\ & & & & & \frac{1}{\sqrt{2}} & & & \\ & & & & & & \frac{1}{\sqrt{2}} & & \\ & & & & & & & \frac{1}{\sqrt{2}} & \\ & \cdots & & & & & & & \cdots \\ \frac{1}{\sqrt{2}} & & & & & & & & \frac{1}{\sqrt{2}} \end{pmatrix} \quad (8.1)$$

Then ${}^tPQ_{p,q}P = I_{p,q}$ thus $\text{SO}(Q_{p,q}) = P \text{SO}(I_{p,q}) P^{-1}$.

Proposition 8.4. • $\text{SO}(Q_{p,q})$ is a Lie group.

• If $p = 0$, then $\text{SO}(Q_{0,q})$ is connected and compact.

8. The Indefinite Special Orthogonal Group

- If $p > 0$, then $\text{SO}(Q_{p,q})$ has two connected components and is not compact.

Denote $\text{SO}_0(Q_{p,q})$ the connected component of $\text{SO}(Q_{p,q})$ containing the identity.

Theorem 8.5. $\text{SO}_0(Q_{p,q})$ is homeomorphic to $\text{SO}(I_p) \times \text{SO}(I_q) \times \mathbb{R}^{pq}$.

Exercise 8.6. Determine the fundamental group of $\text{SO}_0(Q_{p,q})$.

Proposition 8.7. The Lie algebra of $\text{SO}(Q_{p,q})$ is

$$\mathfrak{so}(Q_{p,q}) = \{M \in \text{Mat}(p+q, \mathbb{R}) : {}^t M Q_{p,q} + Q_{p,q} M = 0, \text{Tr}(M) = 0\} =$$

$$\left\{ \begin{pmatrix} M_{11} & M_{12} & M_{13} \\ M_{21} & M_{22} & {}^t M_{12} W_p \\ M_{31} & {}^t W_p {}^t M_{21} & -{}^t W_p {}^t M_{11} W_p \end{pmatrix} : {}^t M_{22} = -M_{22}, M_{13} = -W_p {}^t M_{13} W_p, M_{31} = -W_p {}^t M_{31} W_p \right\}$$

with

$$M_{ij} = -{}^t W_p {}^t M_{ij} W_p \iff M_{ij} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1p-1} & 0 \\ a_{21} & a_{22} & \cdots & 0 & a_{1p-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{p-11} & 0 & \cdots & (-1)^{p-2} a_{22} & (-1)^{p-1} a_{12} \\ 0 & a_{p-11} & \cdots & (-1)^{p-1} a_{21} & (-1)^p a_{11} \end{pmatrix}.$$

The Lie bracket is $[X, Y] = XY - YX$.

We thus see that $\dim \mathfrak{so}(Q_{p,q}) = \frac{(p+q)(p+q-1)}{2}$.

Proposition 8.8. Its Killing form is $B(X, Y) = (p+q-2) \text{Tr}(XY)$ ($p+q \geq 3$).

Since, for $p+q \geq 3$, $B(X, Y)$ is non-degenerate, $\mathfrak{so}(Q_{p,q})$ is semisimple.

Remark 8.9. Let $p+q \geq 3$. Then $\mathfrak{so}(Q_{p,q})$ is simple unless

- $p = q = 2$ because $\mathfrak{so}(Q_{2,2}) = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R})$,
- $p = 0$ and $q = 4$ because $\mathfrak{so}(Q_{0,4}) = \mathfrak{su}(I_2) \oplus \mathfrak{su}(I_2)$.

Exercise 8.10. Prove that $\text{SO}_0(I_{2,2}) \cong \text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R}) / \{\pm(I_2, I_2)\}$ and $\text{SO}_0(I_4) \cong \text{SU}(I_2) \times \text{SU}(I_2) / \{\pm(I_2, I_2)\}$.

Example 8.11. $\text{SO}(Q_{2,3})$ has Lie algebra

$$\mathfrak{so}(Q_{2,3}) = \left\{ \begin{pmatrix} a_{12} & a_{12} & a_{13} & a_{14} & 0 \\ a_{21} & a_{22} & a_{23} & 0 & a_{14} \\ a_{31} & a_{32} & 0 & a_{23} & -a_{13} \\ a_{41} & 0 & a_{32} & -a_{22} & a_{12} \\ 0 & a_{41} & -a_{31} & a_{21} & -a_{11} \end{pmatrix} : a_{ij} \in \mathbb{R} \right\}.$$

8.2. The restricted root system

Let $0 \leq p \leq q$ with $p + q \geq 3$.

Proposition 8.12. *The map $\theta: \mathfrak{so}(Q_{p,q}) \rightarrow \mathfrak{so}(Q_{p,q})$, $M \mapsto -{}^tM$ is a Cartan involution.*

Proof. Since $\mathfrak{so}(Q_{p,q})$ is close under transpose, θ is a Lie involution of $\mathfrak{so}(Q_{p,q})$. For any $X \in \mathfrak{so}(Q_{p,q})$ non-zero $-B(X, \theta(X)) = (p + q - 2) \text{Tr}(X {}^tX) > 0$ \square

Thus $\mathfrak{so}(Q_{p,q}) = \mathfrak{k} \oplus \mathfrak{p}$ where $\mathfrak{k} = \{M \in \mathfrak{so}(Q_{p,q}) \mid -{}^tM = M\}$ and $\mathfrak{p} = \{M \in \mathfrak{so}(Q_{p,q}) \mid {}^tM = M\}$. This is the Cartan decomposition.

Denote

$$\mathfrak{a} = \left\{ D_\lambda = \begin{pmatrix} \lambda_1 & & & & & \\ & \ddots & & & & \\ & & \lambda_p & & & \\ & & & 0 & & \\ & & & & -\lambda_p & \\ & & & & & \ddots \\ & & & & & & -\lambda_1 \end{pmatrix} : \lambda_i \in \mathbb{R} \right\}$$

This is a maximal abelian subspace of \mathfrak{p} as the following computations show.

Denote

$$(E_{kl})_{ij} = \begin{cases} 1 & \text{if } i = k \text{ and } j = l, \\ 0 & \text{otherwise.} \end{cases}$$

Then we compute the following brackets:

$$\left[D_\lambda, \begin{pmatrix} E_{ij} & & & \\ & 0 & & \\ & & (-1)^{i+j+1} E_{p-j+1, p-i+1} & \\ & & & \end{pmatrix} \right] = (\lambda_i - \lambda_j) \begin{pmatrix} E_{ij} & & & \\ & 0 & & \\ & & (-1)^{i+j+1} E_{p-j+1, p-i+1} & \\ & & & \end{pmatrix} \quad (8.2)$$

$$\left[D_\lambda, \begin{pmatrix} 0 & & \\ & E_{ij} & \\ & & 0 \end{pmatrix} \right] = 0 \quad (8.3)$$

$$\left[D_\lambda, \begin{pmatrix} 0 & E_{ij} & & \\ & 0 & (-1)^{p-i} E_{j, p-i+1} & \\ & & & \\ & & & 0 \end{pmatrix} \right] = \lambda_i \begin{pmatrix} 0 & E_{ij} & & \\ & 0 & (-1)^{p-i} E_{j, p-i+1} & \\ & & & \\ & & & 0 \end{pmatrix} \quad (8.4)$$

$$\left[D_\lambda, \begin{pmatrix} 0 & & & \\ E_{ij} & & & \\ & 0 & & \\ & & (-1)^{p-j} E_{p-j+1, i} & 0 \end{pmatrix} \right] = -\lambda_j \begin{pmatrix} 0 & & & \\ E_{ij} & & & \\ & 0 & & \\ & & (-1)^{p-j} E_{p-j+1, i} & 0 \end{pmatrix} \quad (8.5)$$

$$\left[D_\lambda, \begin{pmatrix} & E_{ij} + (-1)^{p+i+j} E_{p-j+1, p-i+1} & & \\ & 0 & & \\ & & & \\ 0 & & & \end{pmatrix} \right] = (\lambda_i + \lambda_{p-j+1}) \begin{pmatrix} & E_{ij} + (-1)^{p+i+j} E_{p-j+1, p-i+1} & & \\ & 0 & & \\ & & & \\ 0 & & & \end{pmatrix} \quad (8.6)$$

$$\left[D_\lambda, \begin{pmatrix} & & & 0 \\ & & & \\ & & & \\ E_{ij} + (-1)^{p+i+j} E_{p-j+1, p-i+1} & & & 0 \end{pmatrix} \right] = (-\lambda_{p-i+1} - \lambda_j) \begin{pmatrix} & & & 0 \\ & & & \\ & & & \\ E_{ij} + (-1)^{p+i+j} E_{p-j+1, p-i+1} & & & 0 \end{pmatrix} \quad (8.7)$$

8. The Indefinite Special Orthogonal Group

Remark 8.13. The rank of $\mathfrak{so}(Q_{p,q})$ is the dimension of \mathfrak{a} which is p .

We see that

$$Z_{\mathfrak{k}}(\mathfrak{a}) = \{X \in \mathfrak{k} : [A, X] = 0 \ \forall A \in \mathfrak{a}\} = \left\{ \begin{pmatrix} 0 & & & \\ & M_{22} & & \\ & & 0 & \\ & & & 0 \end{pmatrix} : M_{22}^\top = -M_{22} \right\}.$$

Definition 8.14. A real semisimple Lie algebra is said to be *split* if $Z_{\mathfrak{k}}(\mathfrak{a}) = \{0\}$.

Thus $\mathfrak{so}(Q_{p,q})$ is split if and only if $p = q$ or $p = q - 1$.

Denote $\varepsilon_i : \mathfrak{a} \rightarrow \mathbb{R}, D_\lambda \mapsto \lambda_i$ for every i .

Proposition 8.15. The set of restricted roots of $\mathfrak{so}(Q_{p,q})$ relative to \mathfrak{a} is

- if $p \neq q$, then $\Sigma = \{\varepsilon_i \ (1 \leq i \leq p), -\varepsilon_i \ (1 \leq i \leq p), \varepsilon_i + \varepsilon_j \ (1 \leq i < j \leq p), -\varepsilon_i - \varepsilon_j \ (1 \leq i < j \leq p), \varepsilon_i - \varepsilon_j \ (1 \leq i \neq j \leq p)\}$,
- if $p = q$, then $\Sigma = \{\varepsilon_i + \varepsilon_j \ (1 \leq i < j \leq p), -\varepsilon_i - \varepsilon_j \ (1 \leq i < j \leq p), \varepsilon_i - \varepsilon_j \ (1 \leq i \neq j \leq p)\}$,

and the root space decomposition is

$$\mathfrak{so}(Q_{p,q}) = \mathfrak{a} \oplus Z_{\mathfrak{k}}(\mathfrak{a}) \oplus \bigoplus_{1 \leq i \leq p} \mathfrak{g}_{\varepsilon_i} \oplus \bigoplus_{1 \leq i \leq p} \mathfrak{g}_{-\varepsilon_i} \oplus \bigoplus_{1 \leq i \neq j \leq p} \mathfrak{g}_{\varepsilon_i - \varepsilon_j} \oplus \bigoplus_{1 \leq i < j \leq p} \mathfrak{g}_{\varepsilon_i + \varepsilon_j} \oplus \bigoplus_{1 \leq i < j \leq p} \mathfrak{g}_{-\varepsilon_i - \varepsilon_j}$$

where $\mathfrak{g}_\lambda = \{X \in \mathfrak{so}(Q_{p,q}) : [A, X] = \lambda(A)X \ \forall A \in \mathfrak{a}\}$ with $\dim(\mathfrak{g}_{\varepsilon_i}) = q - p$, $\dim(\mathfrak{g}_{-\varepsilon_i}) = q - p$, $\dim(\mathfrak{g}_{\varepsilon_i - \varepsilon_j}) = 1$, $\dim(\mathfrak{g}_{\varepsilon_i + \varepsilon_j}) = 1$ and $\dim(\mathfrak{g}_{-\varepsilon_i - \varepsilon_j}) = 1$.

Proof. This is a consequence of the previous bracket computations. \square

Proposition 8.16. If $p \neq q$, then $\Delta = \{\varepsilon_1 - \varepsilon_2, \varepsilon_2 - \varepsilon_3, \dots, \varepsilon_{p-1} - \varepsilon_p, \varepsilon_p\}$ is a set of simple roots. If $p = q$ then $\Delta = \{\varepsilon_1 - \varepsilon_2, \varepsilon_2 - \varepsilon_3, \dots, \varepsilon_{p-1} - \varepsilon_p, \varepsilon_{p-1} + \varepsilon_p\}$ is a set of simple roots.

Proof. We only prove the case $p \neq q$. We have to express each root as a combination of the simple ones involving only non-negative or non-positive coefficients. We see that $\varepsilon_i = \varepsilon_i - \varepsilon_{i+1} + \varepsilon_{i+1} - \varepsilon_{i+2} + \dots + \varepsilon_{p-1} - \varepsilon_p + \varepsilon_p$. Also for $i < j$ $\varepsilon_i + \varepsilon_j = \varepsilon_i - \varepsilon_{i+1} + \varepsilon_{i+1} - \varepsilon_{i+2} + \dots + \varepsilon_{j-1} - \varepsilon_j + 2\varepsilon_j - 2\varepsilon_{j+1} + 2\varepsilon_{j+1} - 2\varepsilon_{j+2} + \dots + 2\varepsilon_{p-1} - 2\varepsilon_p + 2\varepsilon_p$ and $\varepsilon_i - \varepsilon_j = \varepsilon_i - \varepsilon_{i+1} + \varepsilon_{i+1} - \varepsilon_{i+2} + \dots + \varepsilon_{j-1} - \varepsilon_j$. We deduce the other equations by multiplying those by -1 . \square

Exercise 8.17. Check the case $p = q$.

The Dynkin diagram of $\mathfrak{so}(Q_{p,q})$ is B_p if $p \neq q$ and D_p if $p = q$.

Example 8.18. Case of $\mathfrak{so}(Q_{2,3})$:

$$\mathfrak{a} = \left\{ \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & 0 & \\ & & & -\lambda_2 \\ & & & & -\lambda_1 \end{pmatrix} : \lambda_i \in \mathbb{R} \right\}$$

so $\mathfrak{so}(Q_{2,3})$ has rank 2. We have $Z_{\mathfrak{k}}(\mathfrak{a}) = \{0\}$ so $\mathfrak{so}(Q_{2,3})$ is split. Then $\Sigma = \{\varepsilon_1, \varepsilon_2, -\varepsilon_1, -\varepsilon_2, \varepsilon_1 + \varepsilon_2, -\varepsilon_1 - \varepsilon_2, \varepsilon_1 - \varepsilon_2, \varepsilon_2 - \varepsilon_1\}$ and all root spaces have dimension 1. As a set of simple roots we can pick $\Delta = \{\varepsilon_1 - \varepsilon_2, \varepsilon_2\}$ and the Dynkin diagram is thus B_2 .

9. Examples and Exercises - Part 1

by Alex Moriani

9.1. Closed linear groups

In this session we will talk about linear Lie groups, that is, closed Lie subgroups of $\mathrm{GL}(n, \mathbb{R})$. But first, let us just recall how to see the Lie algebra of a linear Lie group as a group of matrices. The group $\mathrm{GL}(n, \mathbb{R})$ is open in $\mathrm{Mat}(n, \mathbb{R})$ (which is a real vector space), and so it is a manifold of dimension n^2 .

Exercise 9.1. *Prove that the general linear group $\mathrm{GL}(n, \mathbb{R})$ is a Lie group. Prove that its Lie algebra is isomorphic to $\mathrm{Mat}(n, \mathbb{R})$ endowed with the usual bracket on matrices $[A, B] = AB - BA$.*

Solution. $\mathrm{GL}(n, \mathbb{R})$ is an open subset of $\mathrm{Mat}(n, \mathbb{R})$ (a real vector space), so a smooth manifold. Multiplication and inverse are polynomial in the coefficients so are smooth and $\mathrm{GL}(n, \mathbb{R})$ is a Lie group. The Lie algebra of a Lie group is defined as being the set of left invariant vector fields, which is in bijection with the tangent space at the identity. For $\mathrm{GL}(n, \mathbb{R})$, the Lie algebra may so be identified with the vector space $\mathrm{Mat}(n, \mathbb{R})$. It will be the same for every closed subgroup of $\mathrm{GL}(n, \mathbb{R})$, its Lie algebra will be a subspace of $\mathrm{Mat}(n, \mathbb{R})$.

Definition 9.1. The linear Lie algebra of a closed Lie subgroup G of $\mathrm{GL}(n, \mathbb{R})$ is its tangent space at I_n seen as a subspace of $\mathrm{Mat}(n, \mathbb{R})$:

$$\mathfrak{g} = \{\gamma'(t) \mid \gamma: \mathbb{R} \rightarrow G \text{ smooth, } \gamma(0) = I_n\},$$

endowed with the usual bracket on matrices $[X, Y] = XY - YX$.

Exercise 9.2. *Show that it is a Lie algebra (vector space, skew-symmetry and Jacobi identity).*

Solution. If $X, Y \in \mathrm{Mat}(n, \mathbb{R})$ corresponding to $x, y: \mathbb{R} \rightarrow \mathrm{GL}(n, \mathbb{R})$, xy is smooth and its derivative at $t = 0$ gives $(xy)'(0) = x'(0) + y'(0) = X + Y$, so $X + Y \in \mathfrak{g}$. Looking at kx , for $k \in \mathbb{R}$, we see that $kX \in \mathfrak{g}$. So \mathfrak{g} is a vector space. Computations show that $[\cdot, \cdot]$ is bilinear skew-symmetric and verify the Jacobi identity. So \mathfrak{g} is a Lie algebra.

Exercise 9.3. *Show for the case of $\mathrm{GL}(n, \mathbb{R})$, its Lie algebra $\mathfrak{gl}(n, \mathbb{R})$ and its linear Lie algebra \mathfrak{g} are isomorphic.*

Hint. They are isomorphic as vector spaces. Recall that the bracket of the Lie algebra $\mathfrak{gl}(n, \mathbb{R})$ is given by the bracket of vector fields ($\mathfrak{gl}(n, \mathbb{R}) = \{\text{left invariant vector fields}\}$).

$$\begin{aligned} \Phi: \mathfrak{g} &\rightarrow \mathfrak{gl}(n, \mathbb{R}) \\ X &\mapsto X^L, \end{aligned}$$

9. Exercises - Part 1

where X^L is the left invariant vector field with $(X^L)_{I_n} = X$. We want to show that this map is a Lie algebra isomorphism. It is well-defined, one-one, onto. We need to check that it respects the Lie bracket, i.e. we have to verify that for every $X, Y \in \mathfrak{g}$, $[X^L, Y^L] = [X, Y]^L = (XY - YX)^L$, and to do that we check that they act in the same way as derivations on functions. It is differential calculus. If this is true, the two vector fields are the same, and so is their value at the identity, i.e. $[X^L, Y^L]_{I_n} = [X, Y] = XY - YX$.

Remark: recall that the exponential map for Lie groups gives the exponential of matrices in a linear group. To find a particular curve with derivative X one can choose $t \mapsto \exp(tX)$. Here is another way of thinking of the linear Lie algebra of a closed linear subgroup G :

$$\mathfrak{g} = \{X \in \text{Mat}(n, \mathbb{R}) \mid \exp(tX) \in G \forall t \in \mathbb{R}\}.$$

9.2. Exercises

Exercise 9.4. Show that the classical groups

- $\text{SL}(n, \mathbb{R}) = \{A \in \text{GL}(n, \mathbb{R}) : \det(A) = 1\}$,
- $\text{Sp}(2n, \mathbb{R}) = \{A \in \text{GL}(2n, \mathbb{R}) : {}^tAJ_{n,n}A = J_{n,n}\} \cap \text{SL}(2n, \mathbb{R})$,
- $\text{SO}(p, q) = \{A \in \text{GL}(p+q, \mathbb{R}) : {}^tAI_{p,q}A = I_{p,q}\} \cap \text{SL}(p+q, \mathbb{R})$,

are closed linear groups, and compute their linear Lie algebras.

Hint. Use the inverse function theorem.

Solution.

- $\text{SL}(n, \mathbb{R}) = \det^{-1}(1)$, $(d_{I_n} \det)(X) = \text{tr}(X)$, so $\mathfrak{sl}(n, \mathbb{R}) = ((d_{I_n} \det))^{-1}(0) = \{X \in \mathfrak{gl}(n, \mathbb{R}) : \text{tr}(X) = 0\}$,
- $\text{Sp}(2n, \mathbb{R}) = f^{-1}(J_{n,n})$, where

$$f: \text{SL}(2n, \mathbb{R}) \rightarrow \text{Mat}(2n, \mathbb{R}), \quad A \mapsto {}^tAJ_{n,n}A.$$

Then $(d_{I_{2n}} f)(X) = {}^tXJ_{n,n} + J_{n,n}X$, so $\mathfrak{sp}(2n, \mathbb{R}) = \{X \in \mathfrak{gl}(2n, \mathbb{R}) : {}^tXJ_{n,n} + J_{n,n}X = 0\}$,

- $\text{SO}(p, q) = g^{-1}(I_{p,q})$, where

$$g: \text{SL}(p+q, \mathbb{R}) \rightarrow \text{Mat}(p+q, \mathbb{R}), \quad A \mapsto {}^tAI_{p,q}A.$$

Thus $\mathfrak{so}(p, q) = \{X \in \mathfrak{gl}(p+q, \mathbb{R}) : {}^tXI_{p,q} + I_{p,q}X = 0\}$.

Exercise 9.5. Compute the Killing form on $\mathfrak{gl}(n, \mathbb{R})$. Compute the Killing form on $\mathfrak{sl}(3, \mathbb{R})$ and $\mathfrak{sp}(4, \mathbb{R})$.

Solution. Taking the basis (E_{ij}) (matrices with zeros, and only one 1 on the i -th row and j -th column), and a lot of patience, one can compute the killing form on $\mathfrak{gl}(n, \mathbb{R})$ to obtain that $B(X, Y) = 2n \text{Tr}(XY) - 2 \text{Tr}(X) \text{Tr}(Y)$. Since $\mathfrak{sl}(n, \mathbb{R})$ is an ideal of $\mathfrak{gl}(n, \mathbb{R})$, its Killing form is the restriction of that of $\mathfrak{gl}(n, \mathbb{R})$, i.e. $B(X, Y) = 2n \text{Tr}(XY)$. For $\mathfrak{sl}(3, \mathbb{R})$, decompose it into skew and symmetric part, so it becomes $\mathfrak{sl}(3, \mathbb{R}) = \mathfrak{so}(3, \mathbb{R}) \oplus \mathfrak{p}$. Take the basis

$$\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

and find the matrix of the Killing form being equal to

$$\begin{pmatrix} -12 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -12 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -12 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 12 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 12 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 12 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 12 & 6 \\ 0 & 0 & 0 & 0 & 0 & 0 & 6 & 12 \end{pmatrix},$$

which is non-degenerate of signature $(5, 3)$.

For $\mathfrak{sp}(2n, \mathbb{R})$, a computation as that of $\mathfrak{gl}(n, \mathbb{R})$ gives us $B(X, Y) = (2n + 2) \text{Tr}(XY)$, which is non-degenerate of signature $(n^2, n^2 + n)$ in $\mathfrak{sp}(2n, \mathbb{R})$.

Exercise 9.6. Show that the Lie algebras above are semisimple Lie algebras, except for $\mathfrak{gl}(n, \mathbb{R})$, which is only reductive (i.e. direct sum of an abelian Lie algebra and a semisimple one).

Remark 9.2. We see that the Killing form is non-degenerate for $\mathfrak{sl}(n, \mathbb{R})$, $\mathfrak{sp}(2n, \mathbb{R})$ and $\mathfrak{so}(p, q)$.

Recall the Cartan criterion: a Lie algebra is semisimple if and only if its Killing form is non-degenerate. Write $\mathfrak{gl}(n, \mathbb{R})$ as $\mathfrak{sl}(n, \mathbb{R}) \oplus \mathbb{R}I_n$.

Exercise 9.7. Find a Cartan decomposition for $\mathfrak{sl}(n, \mathbb{R})$, $\mathfrak{sp}(2n, \mathbb{R})$, $\mathfrak{so}(p, q)$. Find the associated symmetric space of the Lie group $SL(n, \mathbb{R})$ and its maximal flats.

Solution. We can take the skew and symmetric part in each case (that are eigenspaces of the involution $X \mapsto -{}^tX$). This gives the following Cartan decompositions:

- $\mathfrak{sl}(n, \mathbb{R}) = \mathfrak{so}(n, \mathbb{R}) \oplus \mathfrak{p}$ with \mathfrak{p} being the symmetric matrices with trace 0,
- $\mathfrak{sp}(2n, \mathbb{R}) = \mathfrak{k} \oplus \mathfrak{q}$,
- $\mathfrak{so}(p, q) = \mathfrak{so}(p, \mathbb{R}) \oplus \mathfrak{so}(q, \mathbb{R}) \oplus \mathfrak{r}$.

Let us detail the cases $\mathfrak{sl}(n, \mathbb{R})$ and $\mathfrak{sp}(2n, \mathbb{R})$. In $\mathfrak{sl}(n, \mathbb{R})$, the skew part gives all the skew-symmetric matrices, i.e. the Lie algebra of $SO(n, \mathbb{R})$, so a maximal compact of $SL(n, \mathbb{R})$ is $SO(n, \mathbb{R})$. The symmetric part of $\mathfrak{sl}(n, \mathbb{R})$ are the symmetric matrices with trace zero. In \mathfrak{p} , define \mathfrak{a} as the set of diagonal matrices of \mathfrak{p} , i.e. diagonal matrices of trace zero. The space \mathfrak{a} is abelian, and is maximal for this property. Let us say that if you take a symmetric matrix with zero trace, then its exponential will give a positive definite matrix with determinant one (symmetric implies diagonalizable, and then apply exponential to give another symmetric matrix with eigenvalues $\exp(l_i) > 0$, and recall that $\exp \circ \text{Tr} = \det \circ \exp$). So the exponential of \mathfrak{p} will give us $\text{Sym}_+^1(n, \mathbb{R})$ the set of positive definite symmetric matrices of determinant one. The Cartan decomposition can be written as

$$SL(n, \mathbb{R}) \cong SO(n, \mathbb{R}) \times \text{Sym}_+^1(n, \mathbb{R}),$$

9. Exercises - Part 1

which is nothing else than the polar decomposition. The symmetric space is then $X = \mathrm{SL}(n, \mathbb{R})/\mathrm{SO}(n, \mathbb{R}) \cong \mathrm{Sym}_+^1(n, \mathbb{R})$. A maximal flat of X is then given by integrating the subspace $\mathfrak{a} \subset \mathfrak{p}$, and this gives diagonal matrices with positive entries and determinant one.

In $\mathfrak{sp}(2n, \mathbb{R})$, we can decompose the matrices as

$$X = \begin{pmatrix} M_1 & M_2 \\ M_3 & -{}^tM_1 \end{pmatrix},$$

with $M_1 \in \mathfrak{sl}(n, \mathbb{R})$ and M_2, M_3 symmetric. Then we see that the Lie subalgebra \mathfrak{k} is made of matrices $\begin{pmatrix} M_1 & -M_2 \\ M_2 & M_1 \end{pmatrix}$, with M_1 skew- and M_2 symmetric. At the Lie group level, the maximal compact K associated to the Lie sub-algebra \mathfrak{k} can be thought of in a convenient way. We recall the definition

$$\mathrm{U}(n) = \{A \in \mathrm{GL}(n, \mathbb{C}) : A^*A = I_n\},$$

where A^* stands for ${}^t\bar{A}$ (conjugate transpose). Recall also that complex matrices can be seen as real matrices of twice the size by writing

$$A = \mathrm{Re}(A) + i\mathrm{Im}(A) \mapsto \begin{pmatrix} \mathrm{Re}(A) & -\mathrm{Im}(A) \\ \mathrm{Im}(A) & \mathrm{Re}(A) \end{pmatrix}.$$

Looking at $\mathrm{U}(n)$ as a group of real matrices, and doing the computation ${}^tAJ_{n,n}A$ for $A \in \mathrm{U}(n)$, we find $J_{n,n}$, i.e $\mathrm{U}(n) \subset \mathrm{Sp}(2n, \mathbb{R})$. Moreover, computing the Lie algebra of $\mathrm{U}(n)$ seen in $\mathrm{GL}(2n, \mathbb{R})$ we find exactly \mathfrak{k} . Then just remark that $\mathrm{U}(n)$ is connected so we can say that a maximal compact of $\mathrm{Sp}(2n, \mathbb{R})$ is $\mathrm{U}(n)$. The symmetric space of $\mathrm{Sp}(2n, \mathbb{R})$ will be studied in the following worksheet.

Exercise 9.8. Compute the restricted root spaces, the Weyl chambers (and Weyl group), the KAK decomposition and the parabolic subgroups of $\mathrm{SL}(3, \mathbb{R})$.

For $\mathrm{SL}(3, \mathbb{R})$, recall that

$$\mathfrak{a} = \left\{ \begin{pmatrix} \lambda & & \\ & \mu & \\ & & \nu \end{pmatrix} : \lambda, \mu, \nu \in \mathbb{R}, \lambda + \mu + \nu = 0 \right\}.$$

The space \mathfrak{a}^* is generated by f_i , for $1 \leq i \leq 3$, where $f_i \left(\begin{pmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \lambda_3 \end{pmatrix} \right) = \lambda_i$. Let us take

$A = \begin{pmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \lambda_3 \end{pmatrix} \in \mathfrak{a}$, and $X = (x_{ij}) \in \mathfrak{sl}(3, \mathbb{R}) = \mathfrak{g}$. We have

$$[A, X] = \begin{pmatrix} 0 & (\lambda_1 - \lambda_2)x_{12} & (\lambda_1 - \lambda_3)x_{13} \\ (\lambda_2 - \lambda_1)x_{21} & 0 & (\lambda_2 - \lambda_3)x_{23} \\ (\lambda_3 - \lambda_1)x_{31} & (\lambda_3 - \lambda_2)x_{32} & 0 \end{pmatrix},$$

so the roots are $\Sigma = \{\lambda_i - \lambda_j : i \neq j\}$ and the root spaces associated are $\mathfrak{g}_{(ij)} = \{X \in \mathfrak{g} \mid x_{kl} = 0 \forall (kl) \neq (ij)\}$ of dimension one. Here $\mathfrak{g}_0 = \mathfrak{a}$. An example of a Lie algebra with $\mathfrak{a} \neq \mathfrak{g}_0$ is $\mathfrak{sl}(n, \mathbb{C})$, where all works like the real case (with Hermitian instead of symmetric) but $\mathfrak{g}_0 = \mathfrak{a} \oplus i\mathfrak{a}$.

Let us compute the Weyl chambers. Let us take the root $\alpha = \lambda_1 - \lambda_2$ and compute $\ker(\alpha) =$

$\left\{ \begin{pmatrix} d & & \\ & d & \\ & & -2d \end{pmatrix} : d \in \mathbb{R} \right\}$. We can represent the space \mathfrak{a} as a 2-plane, namely the hyperplane $x + y + z = 0$ in \mathbb{R}^3 , and draw the walls we obtain.

10. Examples and Exercises - Part 2

by Colin Davalo

10.1. Hermitian symmetric spaces, in general

Several equivalent definitions of Hermitian symmetric spaces exist. Here is a first definition.

Definition 10.1. A symmetric space (X, g) is called of *Hermitian type* if it admits a complex structure $J: TX \rightarrow TX$ that is invariant by the identity component of G and such that $g(J\cdot, J\cdot) = g(\cdot, \cdot)$.

In particular g becomes a Hermitian metric, and $\omega(\cdot, \cdot) = g(\cdot, J\cdot)$ defines a non-degenerate 2-form on X .

Exercise 10.1. Any differential form on a symmetric space X that is invariant under the identity component of the isometry group is closed. In particular the previous ω is a symplectic form.

Hint. Consider $x \in X$ and the symmetry σ_x at x . Then if ω is an invariant k -form, then $\sigma_x^*\omega$ is also an invariant k -form, equal to $(-1)^k\omega$.

A characterisation of symmetric spaces of Hermitian type is the following: let X whose identity component of the isometry group is G . Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition associated with X . Then X is of Hermitian type if and only if \mathfrak{k} has a non-zero center.

There exist Hermitian symmetric spaces of compact and non-compact type, but we are more interested in the Hermitian symmetric spaces of non-compact type.

10.2. An interesting example

We study one example of an Hermitian symmetric space. Through this example, we try to understand the main features of Hermitian symmetric spaces.

Let $n \geq 1$ be an integer, and let $\mathrm{Sp}(2n, \mathbb{R})$ be the subgroup of $\mathrm{GL}(2n, \mathbb{R})$ of elements preserving the symplectic form ω , which is the bilinear form defined by $\omega(X, Y) = {}^tX J_{n,n} Y$ for $X, Y \in \mathbb{R}^{2n}$ with :

$$J_{n,n} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

The goal of this exercise session is to study the properties of the symmetric space associated with $\mathrm{Sp}(2n, \mathbb{R})$.

10. Exercises - Part 2

Exercise 10.2. Find a Cartan involution, and a maximal abelian subalgebra $\mathfrak{a} \subset \mathfrak{p}$ for $\mathrm{Sp}(2n, \mathbb{R})$. What is the associated restricted root system? And the Weyl group?

Hint. The Killing form is proportional to $A, B \mapsto \mathrm{Tr}(AB)$. The Weyl groups acts by isometry on \mathfrak{a}

A model for this symmetric space associated with $\mathrm{Sp}(2n, \mathbb{R})$ is the *Siegel upper half-space* \mathcal{X}_n , defined as the space of complex $n \times n$ matrices which can be written $S + iT$ with S, T real symmetric $n \times n$ matrices, and T positive definite.

Remark 10.2. When $n = 1$, the Siegel space \mathcal{X}_1 becomes the upper half-plane:

$$\{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}.$$

It is a model for the symmetric space \mathbb{H}^2 associated with $\mathrm{Sp}(2, \mathbb{R}) = \mathrm{SL}(2, \mathbb{R})$.

The symplectic groups acts on the Siegel space in a way that is similar to the action of $\mathrm{SL}(2, \mathbb{R})$ on the upper half-plane. Consider for

$$g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{R}),$$

for some $n \times n$ blocks A, B, C, D . Let $Z = S + iT \in \mathcal{X}_n$ for some symmetric real matrices S, T . We define $g \cdot Z = (AZ + B)(CZ + D)^{-1}$.

Exercise 10.3. Assuming that this is well defined, check that it is an action. Find an equivariant diffeomorphism between \mathcal{X}_n and the symmetric space $\mathrm{Sp}(2n, \mathbb{R})/\mathrm{U}(n)$.

Hint. Show that the stabilizer of $iI_n \in \mathcal{X}_n$ is isomorphic to $\mathrm{U}(n)$.

The action of any element $\mathrm{Sp}(2n, \mathbb{R})$ on \mathcal{X}_n is holomorphic. Hence the symmetric space associated to $\mathrm{Sp}(2n, \mathbb{R})$ inherits a complex structure.

Exercise 10.4. Check that the Riemannian metric actually defines an Hermitian metric for this complex structure.

A Hermitian symmetric space is of *tube type* if it is biholomorphic to a domain of the form $V + i\Omega$ with V a real vector space and Ω a convex cone.

It is handy to realize the symmetric space as a bounded domain of a complex vector space. For every symmetric space of Hermitian type, a general construction called the *Harish-Chandra embedding*. For the group $\mathrm{Sp}(2n, \mathbb{R})$, this domain is \mathcal{D}_n , the space of complex symmetric $n \times n$ matrices Z such that $I_n - {}^t Z \bar{Z}$ is positive definite.

Exercise 10.5. Find a biholomorphism from \mathcal{X}_n to \mathcal{D}_n , inspired from a biholomorphism from the upper half-plane to the unit disk, that can be extended continuously to the boundary of \mathcal{X}_n .

Recall the maximal principle: let f be a continuous function from the closure $\overline{\Omega}$ of a bounded domain Ω of a complex vector space into \mathbb{C} that is holomorphic on Ω . Then:

$$\max_{x \in \overline{\Omega}} |f(x)| = \max_{x \in \partial\Omega} |f(x)|.$$

However this inequality can be improved sometimes. We say that a closed subset $F \subset \partial\Omega$ is a *boundary* if for all function f continuous on $\overline{\Omega}$ and holomorphic on Ω :

$$\max_{x \in \overline{\Omega}} |f(x)| = \max_{x \in F} |f(x)|.$$

The intersection of all boundaries of Ω is called the *Shilov boundary* of Ω .

Exercise 10.6. Show that the Shilov boundary of $D \times D \subset \mathbb{C}^2$ where $D \subset \mathbb{C}$ is the unit disk is the set of pairs $(z_1, z_2) \in \mathbb{C}^2$ such that $|z_1| = |z_2| = 1$.

Show that $\mathrm{Sp}(2n, \mathbb{R})$ acts on $\partial\mathcal{D}_n$, and that any $Z \in \partial\mathcal{D}_n$ such that $Z\overline{Z} = I_n$ is in the same orbit as I_n .

Show that the Shilov boundary of \mathcal{D}_n is the set of symmetric complex matrices Z such that $Z\overline{Z} = I_n$. (harder)

The Shilov boundary of \mathcal{D}_n is invariant by the action of $\mathrm{Sp}(2n, \mathbb{R})$. Let Q_n be the stabilizer of a Lagrangian in \mathbb{R}^{2n} , i.e. of a n -dimensional subspace on which ω is degenerate. The space \mathcal{L}_n of Lagrangians can be seen as the flag manifold $\mathrm{Sp}(2n, \mathbb{R})/Q_n$.

Exercise 10.7. Check that Q_n is a parabolic subgroup. To what set of root is it associated? Show that the Shilov boundary of \mathcal{D}_n is diffeomorphic in a $\mathrm{Sp}(2n, \mathbb{R})$ -equivariant way to \mathcal{L}_n .

Part III.
Known Notions of Positivity

Romeo Troubat
Université de Strasbourg

Xian Dai
Ruprecht-Karls-Universität Heidelberg

Jingyi Xue
National University of Singapore

Raphael Appenzeller
ETH Zürich

Francesco Fournier-Facio
ETH Zürich

Samuel Bronstein
ENS Paris

Ilia Smilga
Institut des Hautes Études Scientifiques

11. Totally Positive Matrices

by Romeo Troubat

In the section, our aim will be to introduce a notion of positivity in $\mathfrak{gl}(n, \mathbb{R})$. First, we will discuss the link between triples of points in S^1 and matrices in $\mathrm{SL}(2, \mathbb{R})$ which will give us a notion of positivity for $\mathrm{SL}(2, \mathbb{R})$. We will then generalize this notion for matrices in $\mathfrak{gl}(n, \mathbb{R})$ using *totally positive matrices*.

11.1. Positivity for the special linear group

The tangent bundle of S^1 is equivalent to the trivial bundle $S^1 \times \mathbb{R}$, thus the choice of a half line in \mathbb{R} gives a causal structure to the space S^1 . If we choose $\mathbb{R}^{>0} \subset \mathbb{R}$, we can for instance say that clockwise rotations in S^1 are positive whereas counter clockwise rotations are negative.

Definition 11.1. A triple of points (x, y, z) in S^1 is said to be *positive* if the points are pairwise distinct and if one has to meet the point y when going from x to z following the positive rotation.

The group $\mathrm{SL}(2, \mathbb{R})$ acts on $S^1 \simeq \mathbb{RP}^1$. Up to an action by $\mathrm{SL}(2, \mathbb{R})$, we can assume that $x = \mathbb{R}e_2$ and $z = \mathbb{R}e_1$ where $e_1 = (0, 1)$ and $e_2 = (1, 0)$. Let us consider the subgroup

$$U := \left\{ g \in \mathrm{SL}(2, \mathbb{R}), g = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \right\}.$$

One could check that U fixes z and acts transitively on $S^1 \setminus \{z\}$. It can be identified with \mathbb{R} using the application

$$t \in \mathbb{R} \longmapsto \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \in U,$$

and the choice of a half-line $\mathbb{R}^{>0} \subset \mathbb{R}$ gives us a subsemigroup $U^{>0} \subset U$ defined by

$$U^{>0} = \left\{ \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, t > 0 \right\}.$$

Any point on S^1 different from z can be written in a unique way as $y = u_y x$. If $y = t_y e_1 + e_2$, then u_y is equal to $\begin{pmatrix} 1 & t_y \\ 0 & 1 \end{pmatrix}$. It is then easy to verify that the triple (x, y, z) is positive if and only if $t_y > 0$, i.e if and only if $u_y \in U^{>0}$.

In the same way, let us define the group

$$O := \left\{ g \in \mathrm{SL}(2, \mathbb{R}), g = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix} \right\}$$

11. Totally Positive Matrices

and its subsemigroup

$$O^{>0} = \left\{ \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}, t > 0 \right\}.$$

Finally, we call $A = \left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \lambda \neq 0 \right\}$ the set of diagonal matrices in $\mathrm{SL}(2, \mathbb{R})$ and $A^\circ = \left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \lambda > 0 \right\}$ the connected component of the identity in A .

Definition 11.2. We call $\mathrm{SL}(2, \mathbb{R})^{>0} = O^{>0}A^\circ U^{>0}$ the set of *totally positive matrices* in $\mathrm{SL}(2, \mathbb{R})$.

Proposition 11.3. *The set of totally positive matrices is the set of matrices in $\mathrm{SL}(2, \mathbb{R})$ whose coefficients are all positive. In particular, it is a subsemigroup of $\mathrm{SL}(2, \mathbb{R})$.*

Proof. It is easy to compute that

$$\begin{pmatrix} 1 & 0 \\ s & 1 \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \lambda & \lambda t \\ \lambda s & \lambda st + \lambda^{-1} \end{pmatrix}.$$

By choosing the parameters λ, s, t correctly, we can put any positive number we want in the top left, top right and bottom left spots in the matrix. The bottom right coefficient is then determined by the fact that the determinant has to be 1. Thus, $\mathrm{SL}(2, \mathbb{R})^{>0} = O^{>0}A^\circ U^{>0}$.

□

11.2. Totally positive matrices

We will now try to generalize the previous notion of positivity to matrices in $\mathfrak{gl}(n, \mathbb{R})$.

Definition 11.4. A matrix in $\mathfrak{gl}(n, \mathbb{R})$ is said to be *totally positive* if all of its minors are positive. We call $\mathfrak{gl}(n, \mathbb{R})^{>0}$ the set of totally positive matrices in $\mathfrak{gl}(n, \mathbb{R})$. We call U (resp. O) the set of upper triangular (resp. lower triangular) matrices with 1 on all diagonal entries and A the set of diagonal matrices in $\mathfrak{gl}(n, \mathbb{R})$.

Proposition 11.5. *The set of totally positive matrices forms a subsemigroup of $\mathfrak{gl}(n, \mathbb{R})$.*

Proof. Let A and B be two totally positive matrices and let I and J be two subsets of $\{1, \dots, n\}$ of cardinal k . We write $A|_I$ the matrix we obtain by extracting the lines in I from A and $A|_{I,J}$ the one we obtain by extracting the lines from I and the columns from J . We have

$$(AB)|_{I,J}(e_1 \wedge \dots \wedge e_k) = A|_I B|_J(e_1 \wedge \dots \wedge e_k).$$

The matrix $B|_J$ is an application from \mathbb{R}^k to \mathbb{R}^n . By projecting $B|_J(e_1 \wedge \dots \wedge e_k)$ onto $e_{i_1} \wedge \dots \wedge e_{i_k}$, we get the determinant of $p \circ B|_J$, where p is the projection on $\langle e_{i_1}, \dots, e_{i_k} \rangle$. Thus, we have

$$B|_J(e_1 \wedge \dots \wedge e_k) = \sum_{|K|=k} \det(B|_{K,J}) e_{i_1} \wedge \dots \wedge e_{i_k},$$

which gives us

$$\begin{aligned} (AB)|_{I,J}(e_1 \wedge \dots \wedge e_k) &= \sum_{|K|=k} \det(B|_{K,J}) A_I(e_{i_1} \wedge \dots \wedge e_{i_k}) \\ &= \left(\sum_{|K|=k} \det(A|_{I,K}) \det(B|_{K,J}) \right) e_1 \wedge \dots \wedge e_k, \end{aligned}$$

thus, $\det((AB)|_{I,J}) = \sum_{|K|=k} \det(A|_{I,K}) \det(B|_{K,J})$. This is known as the *Cauchy-Binet formula*. Since all of A and B 's minors are positive, the product AB is a totally positive matrix. \square

Definition 11.6. We call $U^{>0}$ (resp. $O^{>0}$) the set of matrices in U (resp. O) which have all positive minors, except for those which have to be equal to 0 by virtue of being in U (resp. O). Finally, we call A° the connected component of the identity in A .

We give an explicit parametrisation of the sets $U^{>0}$ and $O^{>0}$. The group U is generated by the elementary matrices

$$u_i(t) = I_n + tE_{i,i+1}, \quad i = 1, \dots, n-1.$$

We call $U^{\geq 0}$ the group generated by the elements $u_i(t)$ for $i = 1, \dots, n-1$ and $t > 0$. For the case $n = 2$, we have shown that $U^{\geq 0} = U^{>0}$, but that is no longer the case for $n \geq 3$ since the matrices $u_i(t)$, $t > 0$, are not contained in $U^{>0}$ as too many of their minors are equal to zero.

To get the appropriate parametrisation, let us study the group S_n of permutations on the set $\{1, \dots, n\}$. Each of its elements can be decomposed as a product of adjacent transpositions $(i, i+1)$ and the length of a permutation can be defined as the smallest number of adjacent transposition in one of its decomposition. For each group S_n , there exists a permutation ω_0 of maximum length,

$$\omega_0 = (n, n-1)(n-1, n-2) \dots (2, 1)(n, n-1)(n-1, n-2) \dots (3, 2) \dots (n, n-1)(n-1, n-2)(n, n-1).$$

For $n = 4$, we have $\omega_0 = (4, 3)(3, 2)(2, 1)(4, 3)(3, 2)(4, 3)$. Let $\omega_0 = \sigma_{i_1} \dots \sigma_{i_k}$ be a decomposition of $\omega_0 \in S_n$ and let us define the map

$$F_{\sigma_{i_1} \dots \sigma_{i_k}} : (t_1, \dots, t_k) \in \mathbb{R}^k \longmapsto u_{i_1}(t_1) \dots u_{i_k}(t_k).$$

Proposition 11.7. *The map $F_{\sigma_{i_1} \dots \sigma_{i_k}}|_{(\mathbb{R}^+)^k}$ is a bijection unto $U^{>0}$. It provides a parametrisation of $U^{>0}$. There is a symmetric result for $O^{>0}$.*

11. Totally Positive Matrices

For $n = 3$, this gives us the parametrisation

$$u_1(a)u_2(b)u_1(c) = \begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & a+c & ab \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$$

In the case $n = 2$, we defined the set of totally positive matrices using the decomposition $\mathrm{SL}(2, \mathbb{R})^{>0} = O^{>0}A^\circ U^{>0}$. Let us show a similar result for $n \geq 3$.

Proposition 11.8. *For $n \geq 2$, the set of totally positive matrices can be decomposed as $\mathfrak{gl}(n, \mathbb{R})^{>0} = O^{>0}A^\circ U^{>0}$.*

Proof. Let $A = (a_{i,j})$ be a totally positive matrix. In particular, all of its coefficients are positive. We are going to reduce our matrix using operations on lines and columns. First, let us subtract $\frac{a_{i,1}}{a_{i-1,1}}L_{i-1}$ to L_i for $i = 2, \dots, n$. This gives us the matrix

$$A = \left(I_n + \frac{a_{n,1}}{a_{n-1,1}}E_{n,n-1} \right) \cdots \left(I_n + \frac{a_{2,1}}{a_{1,1}}E_{2,1} \right) \begin{pmatrix} 1 & a_{1,j} \\ 0 & a_{2,j} - \frac{a_{2,1}}{a_{1,1}} \\ \vdots & \vdots \\ 0 & a_{n,j} - \frac{a_{n,1}}{a_{n-1,1}} \end{pmatrix}$$

Let us write $B = \begin{pmatrix} a_{1,j} \\ a_{2,j} - \frac{a_{2,1}}{a_{1,1}} \\ \vdots \\ a_{n,j} - \frac{a_{n,1}}{a_{n-1,1}} \end{pmatrix}$. We admit that the matrix A is totally positive if and only if

B is totally positive [Whi52]. By repeating the same process, we can reach the decomposition

$$A = \left(I_n + \frac{a_{n,1}}{a_{n-1,1}}E_{n,n-1} \right) \cdots \left(I_n + \frac{a_{n-1,n-1}}{a_{n-2,n-1}}E_{n-1,n-2} \right) \left(I_n + \frac{a_{n,n}}{a_{n-1,n}}E_{n,n-1} \right) C,$$

where C is an upper triangular matrix. Using the parametrisation we have shown, we can see that the product of transvections is equal to a matrix in $O^{>0}$. We now only have to repeat the same operations on the columns to obtain the decomposition $\mathfrak{gl}(n, \mathbb{R})^{>0} = O^{>0}A^\circ U^{>0}$. \square

12. Lusztig's Total Positivity

by Xian Dai

Lusztig's total positivity generalizes total positivity for $n \times n$ matrices in $\mathrm{GL}(n, \mathbb{R})$ to split real Lie groups. Our goal in this section is to explain this general notion of total positivity.

12.1. Some motivation for total positivity

We recall a $n \times n$ matrix is said to be (*totally*) *positive* if all of its minors are positive (i.e. in $\mathbb{R}^{>0}$). The set of all totally positive $n \times n$ matrices form a subset $\mathrm{GL}(n, \mathbb{R})^{>0} \subset \mathrm{GL}(n, \mathbb{R})$.

Let $V \cong \mathbb{R}^n$. Another characterization of total positivity in $\mathrm{GL}(n, \mathbb{R})$ due to I.J. Schoenberg [Sch30] is as follows.

Proposition 12.1. *A matrix $g \in \mathrm{GL}(V)$ is positive if and only if the coefficients of k -th exterior power $\Lambda^k g: \Lambda^k V \rightarrow \Lambda^k V$ are positive for $k = 1, 2, \dots, n$.*

Proof. Suppose a totally ordered basis of V is e_1, e_2, \dots, e_n , then for any $k \in [1, n]$, the k -th exterior power $\Lambda^k V$ has a basis $(e_{r_1} \wedge e_{r_2} \cdots \wedge e_{r_k})$ indexed by the sequences $r_1 < r_2 < \cdots < r_k$ in $[1, n]$ and $\Lambda^k g: \Lambda^k V \rightarrow \Lambda^k V$ are positive for $k = 1, 2, \dots, n$ is given by

$$\Lambda^k g(e_{r_1} \wedge e_{r_2} \cdots \wedge e_{r_k}) = g e_{r_1} \wedge g e_{r_2} \cdots \wedge g e_{r_k}.$$

The conclusion follows once we notice that coefficients of above matrices are exactly minors of g .

□

One source of the idea of total positivity appear in the work of Gantmacher and Krein in 1935. The following Theorem can be viewed as one of the initial motivation of total positivity.

Theorem 12.2 (Gantmacher-Krein). *If $g \in \mathrm{GL}(n, \mathbb{R})$ is totally positive, then g has distinct real positive eigenvalues.*

Proof. Suppose $g \in \mathrm{GL}(n, \mathbb{R})^{>0}$. We denote c_1, \dots, c_n to be eigenvalues of g arranged so that $|c_1| \geq |c_2| \geq \cdots \geq |c_n|$. Then the eigenvalues of $\Lambda^k g$ are $c_{r_1} c_{r_2} \cdots c_{r_k}$ for $r_1 < r_2 < \cdots < r_k$ and $r_i \in [1, n]$. For instance, the first two eigenvalues of $\Lambda^k g$ in decreasing order of absolute values are $c_1 c_2 \cdots c_k$ and $c_1 c_2 \cdots c_{k-1} c_{k+1}$. Since $\Lambda^k g$ has positive entries by Proposition 12.1. By Perron-Frobenius Theorem, we must have

$$c_1 c_2 \cdots c_k > 0$$

Taking $k = 1$ yields $c_1 > 0$. By induction on k , we obtain $c_1 > c_2 > \cdots > c_n > 0$.

□

12. Lusztig's Total Positivity

12.2. Lusztig's total positivity for matrices

We start from a presentation of Lusztig's total positivity in a simple case which is $\mathrm{GL}(n, \mathbb{R})$. Totally positive matrices in $\mathrm{GL}(n, \mathbb{R})$ satisfy a decomposition theorem and form in fact a semigroup.

Let U be the group of upper triangular matrices with ones on the diagonal, O be the group of lower triangular matrices with ones on the diagonal, and A the group of diagonal matrices. The group U is generated by elementary matrices

$$u_i(t) = I_n + tE_{i,i+1}, \quad i = 1, \dots, n-1,$$

where I_n denotes the identity matrix and $E_{i,i+1}$ the matrix with the single entry 1 in the i -th row and $(i+1)$ -th column.

Let furthermore $U^{>0} \subset U$ and $O^{>0} \subset O$ be the subsets of totally positive unipotent matrices, i.e. those matrices of U , where all minors are positive, except those which have to be zero by the condition of being an element of U , similarly for O . We can parametrize the set $U^{>0}$ using the symmetric group S_n on n letters. We denote by σ_i , $i = 1, \dots, n-1$, the transposition $(i, i+1)$ and by ω_0 the longest element of the symmetric group, which send $(1, 2, \dots, n)$ to $(n, \dots, 2, 1)$. Let $k = \frac{n(n-1)}{2}$. For every way to write $\omega_0 = \sigma_{i_1}\sigma_{i_2}\cdots\sigma_{i_k}$, we can define a map

$$\begin{aligned} F_{\sigma_{i_1}\sigma_{i_2}\cdots\sigma_{i_k}} : \mathbb{R}^k &\rightarrow U \\ (t_1, \dots, t_k) &\mapsto u_{i_1}(t_1)u_{i_2}(t_2)\cdots u_{i_k}(t_k). \end{aligned}$$

An element is in $U^{>0}$ if and only if it is of the form $u_{i_1}(t_1)u_{i_2}(t_2)\cdots u_{i_k}(t_k)$ with $t_i \in \mathbb{R}_+$. Therefore the map $F_{\sigma_{i_1}\sigma_{i_2}\cdots\sigma_{i_k}}|_{(\mathbb{R}_{>0})^k}$ is a bijection onto $U^{>0}$ and provide a parametrization of $U^{>0}$ by \mathbb{R}_+^k . The same works for $O^{>0}$.

The following decomposition theorem is due to A. Whitney [Whi52]:

$$\mathrm{GL}(n, \mathbb{R})^{>0} = O^{>0}A^0U^{>0},$$

where A^0 is the connected component of the identity in A , i.e. the diagonal matrices all of whose entries are positive.

12.3. Lusztig's total positivity for split real Lie groups

In this subsection, we want to explain the generalized notion of total positivity of arbitrary split real reductive Lie group introduced by Lusztig [Lus94]. Moreover, the explicit parametrization of $U^{>0}$ (resp. $O^{>0}$) can also be generalized to arbitrary split real reductive Lie group.

Let G be a split semisimple algebraic group over \mathbb{R} . Let \mathfrak{g} be the Lie algebra of G . To describe total positivity in G , we start from some discussion of some important ingredients involved in defining positivity in this general setting.

12.3.1. Root space decomposition and Chevalley generators

Fix a Cartan subalgebra \mathfrak{h} of \mathfrak{g} . We denote by Σ the set of all roots and Σ^+ a choice of positive roots (resp. Σ^- negative roots) and $\Delta \subset \Sigma^+$ the set of simple roots. The Lie algebra \mathfrak{g} admits the root space decomposition,

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}_\alpha.$$

Denote h_α as the coroot of $\alpha \in \Sigma$, we say a family $(X_\alpha)_{\alpha \in \Sigma}$ is a *Chevalley basis* for $(\mathfrak{g}, \mathfrak{h})$ (see [Bou75, Chapter VIII, Section 2.4, Definition 3] or [Hum72, Chapter VII, Section 25]) if

- (1) $X_\alpha \in \mathfrak{g}_\alpha$ for all $\alpha \in \Sigma$.
- (2) $[X_\alpha, X_{-\alpha}] = -h_\alpha$ for all $\alpha \in \Sigma$.
- (3) the linear map from \mathfrak{g} to \mathfrak{g} which is equal to -1 on the Cartan subalgebra \mathfrak{a} and takes X_α to $X_{-\alpha}$ for all $\alpha \in \Sigma$ is an automorphism of \mathfrak{g} .

Take $e_\alpha = X_\alpha$ and $f_\alpha = X_{-\alpha}$ for $\alpha \in \Delta$ so that $(e_\alpha, f_\alpha, h_\alpha)_{\alpha \in \Delta}$ form a *Chevalley generators* of \mathfrak{g} . We define

$$x_\alpha(t) = \exp(te_\alpha), \quad y_\alpha(t) = \exp(tf_\alpha)$$

In analogy to the last subsection, the elements e_α (resp. f_α) play the roles of $E_{i,i+1}$ (resp. $E_{i+1,i}$). Also $x_\alpha(t)$ (resp. $y_\alpha(t)$) are similar to $u_i(t)$ (resp. $v_i(t)$) in $\mathrm{GL}(n, \mathbb{R})$ case.

12.3.2. Longest element in Weyl group

We denote by W the Weyl group of G generated by s_α for $\alpha \in \Delta$. W together with $(s_\alpha)_{\alpha \in \Delta}$ is a Coxeter group.

Let $l: W \rightarrow \mathbb{N}$ be the standard length function. For $\omega \in W$, we let Δ_ω be the set of all sequences $(\alpha_1, \alpha_2, \dots, \alpha_p)$ in Δ so that $p = l(\omega)$ and

$$s_{\alpha_1} s_{\alpha_2} \cdots s_{\alpha_p} = \omega.$$

We let ω_0 be the element such that $p_0 = l(\omega_0)$ is maximum. To make a link with the theory in the last subsection, one notices that the symmetric group S_n is the Weyl group for the case $\mathrm{GL}(n, \mathbb{R})$.

 12.3.3. Parametrization of U^\pm and total positivity

With a fixed ω_0 so that $p_0 = l(\omega_0)$ reaches maximum, we can define maps

$$\begin{aligned} \Phi_{\omega_0}^\pm : \mathbb{R}_{\geq 0}^{p_0} &\rightarrow U^\pm \\ \Phi_{\omega_0}^+(a_1, a_2, \dots, a_{p_0}) &:= x_{\alpha_1}(a_1) x_{\alpha_2}(a_2) \cdots x_{\alpha_{p_0}}(a_{p_0}), \\ \Phi_{\omega_0}^-(a_1, a_2, \dots, a_{p_0}) &:= y_{\alpha_1}(a_1) y_{\alpha_2}(a_2) \cdots y_{\alpha_{p_0}}(a_{p_0}). \end{aligned}$$

Here U^\pm are the unipotent radicals of the Borel subgroups (minimal parabolic subgroups) B^\pm corresponding to Σ^\pm respectively. It turns out that the definition of $\Phi_{\omega_0}^\pm$ do not depend on a particular choice of $(\alpha_1, \alpha_2, \dots, \alpha_{p_0})$.

12. Lusztig's Total Positivity

We then define

$$U_{\omega_0, \geq 0}^\pm := \Phi_{\omega_0}^\pm(\mathbb{R}_{\geq 0}^p) \subset U^\pm,$$

and

$$U_{\omega_0, > 0}^\pm := \Phi_{\omega_0}^\pm(\mathbb{R}_{> 0}^p) \subset U^\pm.$$

The sets $U_{\geq 0}^\pm = U_{\omega_0, \geq 0}^\pm$ and $U_{> 0}^\pm = U_{\omega_0, > 0}^\pm$ are semigroups of G .

Proposition 12.3. *The maps $\Phi_{\omega_0}^\pm$ are homeomorphisms between $(\mathbb{R}^{> 0})^p$ and $U_{> 0}^\pm$. Therefore $U_{> 0}^\pm$ are cells.*

Now consider

$$\begin{aligned} G_{\geq 0} &= U_{\geq 0}^+ T_{> 0} U_{\geq 0}^- = U_{\geq 0}^- T_{> 0} U_{\geq 0}^+, \\ G_{> 0} &= U_{> 0}^+ T_{> 0} U_{> 0}^- = U_{> 0}^- T_{> 0} U_{> 0}^+, \end{aligned}$$

where T is the subgroup of G with Lie algebra \mathfrak{a} (sometimes called the maximal torus of G). We let T^0 be the connected component of T containing the identity. $G_{\geq 0}$ and $G_{> 0}$ are semigroups in G . More precisely, $G_{\geq 0}$ is a semigroup with the identity element e of G but $G_{> 0}$ is a semigroup without the identity element e .

As one could easily guess, an element g in G is called (*totally*) *positive* if $g \in G_{> 0}$. We have $G_{> 0}$ is also a cell because of the following,

Theorem 12.4. *The map $U_{> 0}^+ \times T_{> 0} \times U_{> 0}^- \rightarrow G$, given by multiplication in G , is a homeomorphism onto the subset $G_{> 0} \subset G$.*

This can be viewed as a generalisation of the Whitney decomposition mentioned in the last subsection.

13. Positivity of Lie Groups of Hermitian Type

by Jingyi Xue

13.1. Examples of Hermitian type Lie groups

13.1.1. $\mathrm{Sp}(2n, \mathbb{R})$

Again we first look at the symplectic group $\mathrm{Sp}(2n, \mathbb{R})$, which is the subgroup of $\mathrm{GL}(2n, \mathbb{R})$ of elements preserving the symplectic form ω , which is the bilinear form defined by $\omega(X, Y) = {}^tX J_{n,n} Y$ for $X, Y \in \mathbb{R}^{2n}$ with :

$$J_{n,n} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

Recall from the previous sections, $\mathrm{U}(n)$ embeds as a maximal compact subgroup. Results of this example come from Bump's book [Bum13, Chap. 28].

Proposition 13.1. *If $Z = X + iY \in \mathcal{X}_n$ (the Siegel upper half-space) and $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{R})$, then $CZ + D$ is invertible. Define*

$$g(Z) = (AZ + B)(CZ + D)^{-1}$$

Then $g(Z) \in \mathcal{X}_n$, and this defines an action of $\mathrm{Sp}(2n, \mathbb{R})$ on \mathcal{X}_n . The action is transitive, and the stabilizer of $iI_n \in \mathcal{X}_n$ is $\mathrm{U}(n)$. If W is the imaginary part of $g(Z)$ then

$$W = \left({}^t\bar{Z}C + {}^tD \right)^{-1} Y (CZ + D)^{-1}.$$

Proof. Using the conditions for g being in $\mathrm{Sp}(2n, \mathbb{R})$, one easily checks that

$$\frac{1}{2i} \left(\left({}^t\bar{Z}C + {}^tD \right) (AZ + B) - \left({}^t\bar{Z}A + {}^tB \right) (CZ + D) \right) = \frac{1}{2i} (Z - \bar{Z}) = Y,$$

From this it follows that $CZ + D$ is invertible since if it had a nonzero nullvector v , then we would have ${}^t\bar{v}Yv = 0$, which is impossible since $Y \gg 0$. To check that $g(Z)$ is symmetric, $g(Z) = {}^t\bar{g}(Z)$ is equivalent to

$$(AZ + B) ({}^t\bar{Z}C + {}^tD) = ({}^t\bar{Z}A + {}^tB) (CZ + D)$$

which is easily confirmed. Next the imaginary part W of $g(Z)$ is positive definite since

$$W = \frac{1}{2i} (g(Z) - \overline{g(Z)}) = \frac{1}{2i} \left((AZ + B)(CZ + D)^{-1} - \left({}^t\bar{Z}C + {}^tD \right)^{-1} \left({}^t\bar{Z}A + {}^tB \right) \right)$$

Simplifying this gives the desired expression; and W is Hermitian (real) and that $W \gg 0$. It is easy to check that $g(g'(Z)) = (gg')(Z)$. To show that this action is transitive, note that if $Z = X + iY \in \mathcal{X}_n$, then

$$\begin{pmatrix} I_n & -X \\ 0 & I_n \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{R}),$$

13. Positivity of Lie Groups of Hermitian Type

and this matrix takes Z to iY . Now if $h \in \mathrm{GL}(n, \mathbb{R})$, then $\begin{pmatrix} h & 0 \\ 0 & {}^t h^{-1} \end{pmatrix}$ takes iY to iY' , where $Y' = hY {}^t h$. Since $Y \gg 0$, we may choose h so that $Y' = I_n$. This shows that any element in \mathcal{X}_n may be moved to iI_n , and the action is transitive. To check that $\mathrm{U}(n)$ is the stabilizer of iI_n , let

$$A(iI_n) + B = (C(iI_n) + D)(iI_n)$$

which implies $A = D, B = -C$. □

Therefore $\mathrm{Sp}(2n, \mathbb{R})/\mathrm{U}(n) \cong \mathcal{X}_n$ is a tube type Hermitian symmetric space since $\mathrm{U}(n)$ has center S^1 .

Just like the classical Cayley transform $c(z) = \frac{z-i}{z+i}$ which maps \mathbb{H}^2 to the unit disc \mathbb{D}^2 , the generalized Cayley transform is applicable to Hermitian symmetric spaces. It was shown by Cartan and Harish-Chandra that any non-compact type Hermitian symmetric space is biholomorphic to a bounded domain in a complex vector space. Piatetski-Shapiro [PS69] gave unbounded realizations. Korányi and Wolf [WK65, KW65] gave a completely general theory relating bounded symmetric domains to unbounded ones by means of the Cayley transform.

Consider now the Cayley transform $c \in \mathrm{Sp}(2n) = \mathrm{Sp}(2n, \mathbb{C}) \cap \mathrm{U}(2n)$ for $\mathrm{Sp}(2n, \mathbb{R})/\mathrm{U}(n)$:

$$c = \frac{1}{\sqrt{2i}} \begin{pmatrix} I_n & -iI_n \\ I_n & iI_n \end{pmatrix}, \quad c^{-1} = \frac{1}{\sqrt{2i}} \begin{pmatrix} iI_n & iI_n \\ -I_n & I_n \end{pmatrix}$$

To embed \mathcal{X}_n in its compact dual $G_c/K = \mathrm{Sp}(2n)/\mathrm{U}(n)$, the first step is to interpret G_c/K as an analog of the Riemann sphere, a space on which the actions of both groups G and G_c may be realized as linear fractional transformations.

Define the *Siegel parabolic subgroup* $H^{\mathbb{C}}W^{\mathbb{C}}$ (see below) of $G^{\mathbb{C}} = \mathrm{Sp}(2n, \mathbb{C})$:

$$P = \left\{ \begin{pmatrix} h & 0 \\ 0 & {}^t h^{-1} \end{pmatrix} \begin{pmatrix} I & X \\ 0 & I \end{pmatrix} : h \in \mathrm{GL}(n, \mathbb{C}), X \in \mathrm{Mat}(n, \mathbb{C}), X = {}^t X \right\}.$$

Followed from Iwasawa decomposition, it is easy to verify the following:

Proposition 13.2.

$$P \mathrm{Sp}(2n) = \mathrm{Sp}(2n, \mathbb{C}), \quad P \cap \mathrm{Sp}(2n) = cKc^{-1} = \left\{ \begin{pmatrix} g & 0 \\ 0 & {}^t g^{-1} \end{pmatrix} : g \in \mathrm{U}(n) \right\}.$$

Define $\mathfrak{X}_n = G_{\mathbb{C}}/P$ with dense open subset

$$\mathfrak{X}_n^{\circ} = \left\{ gP : g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2n, \mathbb{C}), \det C \neq 0 \right\} \subset \mathfrak{X}_n.$$

Here notice that if g is in $\mathrm{Sp}(2n, \mathbb{C})$ of block form with C invertible, then $AC^{-1} = {}^t C^{-1} \cdot {}^t C A \cdot C^{-1}$ is symmetric since ${}^t C A$ is symmetric by definition. g, g' define same coset if and only if there exists some $h \in \mathrm{GL}(n, \mathbb{C})$ such that $A' = Ah, C' = Ch$ if and only if $A'(C')^{-1} = AC^{-1}$. From this observation, we may define a bijection σ from $\mathrm{Sym}(n, \mathbb{C})$ to \mathfrak{X}_n° by $\sigma(Z) = \begin{pmatrix} Z & -I \\ I & 0 \end{pmatrix} P$,

which can be rewritten as $\begin{pmatrix} A & B \\ C & D \end{pmatrix} P$ if and only if $AC^{-1} = Z$.

With the above notations, we have

$$g(\sigma(Z)) = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} Z & -I \\ I & \end{pmatrix} P = \begin{pmatrix} AZ + B & -A \\ CZ + D & -C \end{pmatrix} P = \sigma((AZ + B)(CZ + D)^{-1})$$

if $CZ + D$ is invertible.

Therefore we may identify $\text{Sym}(n, \mathbb{C})$ with \mathfrak{X}_n° , and the action of $\text{Sp}(2n, \mathbb{C})$ is by linear fractional transformations.

We may also identify \mathfrak{X}_n with G_c/K by means of the compositions of bijections

$$G_c/K \xrightarrow{\text{conjugation}} G_c/cKc^{-1} \xrightarrow{\text{Proposition 13.2}} G^{\mathbb{C}}/P = \mathfrak{X}_n$$

Now we may apply c to embed \mathcal{X}_n into its compact dual \mathfrak{X}_n :

Proposition 13.3.

$$c(\mathcal{X}_n) = \mathfrak{D}_n := \{W \in \mathfrak{X}_n^\circ : I - \overline{W}W > 0\} = \{W \in \text{Mat}(n, \mathbb{C}) : W = {}^t\overline{W}, I - W^*W > 0\}.$$

The group $c\text{Sp}(2n, \mathbb{R})c^{-1}$, acting on \mathfrak{D}_n by linear fractional transformations, consists of all symplectic matrices of the form

$$\begin{pmatrix} A & B \\ \overline{B} & \overline{A} \end{pmatrix}.$$

Proof. This is a direct computation:

$$\begin{aligned} \text{Im}(c^{-1}(W)) &= \text{Im}(-i(W - I)(W + I)^{-1}) \\ &= -\frac{1}{2}((W - I)(W + I)^{-1} + (\overline{W} - I)(\overline{W} + I)^{-1}) \\ &= -\frac{1}{2}((W - I)(W + I)^{-1} + (\overline{W} + I)^{-1}(\overline{W} - I)). \end{aligned}$$

This is positive definite if and only if $(\overline{W} + I)\text{Im}(c^{-1}(W))(W + I) > 0$; where the latter is

$$-\frac{1}{2}((\overline{W} + I)(W - I) + (\overline{W} - I)(W + I)) = I - \overline{W}W.$$

For $g \in \text{Sp}(2n, \mathbb{C})$, $c^{-1}gc = \overline{c^{-1}gc}$ gives us the desired form of elements in $c\text{Sp}(2n, \mathbb{R})c^{-1}$. \square

Proposition 13.4. (1) The closure of \mathfrak{D}_n is contained within \mathfrak{X}_n° . The boundary of \mathfrak{D}_n consists of all complex symmetric matrices W such that $I - \overline{W}W$ is positive semidefinite but such that $\det(I - \overline{W}W) = 0$.

- (2) If W and W' are points of the closure of \mathfrak{D}_n in \mathfrak{X}_n° that are congruent modulo cGc^{-1} , then the ranks of $I - \overline{W}W$ and $I - \overline{W}'W'$ are equal.
- (3) Let W be in the closure of \mathfrak{D}_n , and let r be the rank of $I - \overline{W}W$. Then there exists $g \in cGc^{-1}$ such that $g(W)$ has the form

$$\begin{pmatrix} W_r & 0 \\ 0 & I_{n-r} \end{pmatrix}, \quad W_r \in \mathfrak{D}_r.$$

13. Positivity of Lie Groups of Hermitian Type

Proof. The diagonal entries in $\overline{W}W$ are the squares of the lengths of the rows of the symmetric matrix W . If $I - \overline{W}W$ is positive definite, these must be less than 1. So \mathfrak{D}_n is a bounded domain within the set \mathfrak{R}_n° . The rest of (1) is clear.

For (2), if $g \in cGc^{-1}$, see Proposition 13.3 of the form, $W' = g(W)$,

$$I - \overline{W}'W' = I - (\overline{W}^t B + {}^t A)^{-1} (\overline{W}^t \overline{A} + {}^t \overline{B}) (AW + B)(\overline{B}W + \overline{A})^{-1}.$$

Now the rank of $I - \overline{W}'W'$ is the same as the rank of

$$\begin{aligned} (\overline{W}^t B + {}^t A) (I - \overline{W}'W') (\overline{B}W + \overline{A}) &= (\overline{W}^t B + {}^t A)(\overline{B}W + \overline{A}) - (\overline{W}^t \overline{A} + {}^t \overline{B})(AW + B) \\ &= I - \overline{W}W. \end{aligned}$$

For (3), observe that up to an action by $\begin{pmatrix} u & 0 \\ 0 & \bar{u} \end{pmatrix}$ for some $u \in U(n)$, we may assume that W does not have 1 as an eigenvalue. Now $c^{-1}(W) = i(I+W)(I-W)^{-1} = X+iY \in \overline{\mathfrak{X}}_n$. Since $Y \geq 0$, there exists some $k \in O(n)$ such that $D = kYk^{-1}$ is diagonal with nonnegative eigenvalues. Consider $\gamma = \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix} \begin{pmatrix} I & -X \\ 0 & I \end{pmatrix} \in \text{Sp}(2n, \mathbb{R})$, $\gamma(X+iY) = iD$, then $c\gamma c^{-1}(W) = W' = (D-I)(D+I)^{-1}$. The diagonal entries of $I - \overline{W}'W'$ are 0 if and only if the corresponding entries of D are 0. So after exchanging order of orthogonal basis of k , we will write $W' = \text{diag}(w_1, \dots, w_r, -1, \dots, -1)$, where $-1 < w_i < 1$. Then $\begin{pmatrix} iI_r & 0 \\ 0 & -iI_{n-r} \end{pmatrix} \cdot W'$ will be the desired special form. \square

Now let us fix $r < n$ and consider

$$\mathfrak{B}_r = \left\{ \begin{pmatrix} W_r & 0 \\ 0 & I_{n-r} \end{pmatrix} : W_r \in \mathfrak{D}_r \right\}$$

By Proposition 13.3, the subgroup of cGc^{-1} of the form

$$\begin{pmatrix} A_r & 0 & B_r & 0 \\ 0 & I_{n-r} & 0 & 0 \\ \overline{B}_r & 0 & \overline{A}_r & 0 \\ 0 & 0 & 0 & I_{n-r} \end{pmatrix}$$

is isomorphic to $\text{Sp}(2r, \mathbb{R})$, and \mathfrak{B}_r is homogeneous with respect to this subgroup. Thus, \mathfrak{B}_r is a copy of the lower-dimensional Siegel space \mathfrak{D}_r embedded into the boundary of \mathfrak{D}_n .

Theorem 13.5. *The Shilov boundary of \mathfrak{D}_n is $\text{Sym}(n, \mathbb{C}) \cap U(n)$.*

Proof. Let $f \in \mathcal{O}(\mathfrak{D}_n) \cap \mathcal{C}(\overline{\mathfrak{D}_n})$, we show that $|f|$ must take its maximum on $\text{Sym}(n, \mathbb{C}) \cap U(n)$, which is the unique closed cGc^{-1} -orbit in $\overline{\mathfrak{D}_n}$ (union of all orbits of \mathfrak{B}_0). This is sufficient because cGc^{-1} acts transitively on $\text{Sym}(n, \mathbb{C}) \cap U(n)$ (Proposition 13.4), so the set $\text{Sym}(n, \mathbb{C}) \cap U(n)$ cannot be replaced by any strictly smaller subspace with the same maximizing property.

Suppose $x \in \overline{\mathfrak{D}_n}$ maximizes $|f|$. By Proposition 13.4, $\exists g \in cGc^{-1}$, s.t. $g \cdot x \in \mathfrak{B}_r$ for some $r < n$. If $r > 0$, then noting that $f|_{g^{-1} \cdot \mathfrak{B}_r} \in \mathcal{O}(g^{-1} \cdot \mathfrak{B}_r)$, the maximum modulus principle implies that f is constant on $g^{-1} \cdot \mathfrak{B}_r$ and hence $|f|$ takes the same maximum value on $\partial(g^{-1} \cdot \mathfrak{B}_r)$, which intersects $\text{Sym}(n, \mathbb{C}) \cap U(n)$. \square

Notice that the action of $U(n)$ on Lagrangian Grassmannians \mathcal{L} ($U(n)$ embedded in $\text{Sp}(2n, \mathbb{R})$) is transitive; the stabilizer of a point is readily verified to be $O(n)$ and we get $\mathcal{L} \simeq U(n)/O(n)$. Now consider the action of $U(n)$ on $\text{Sym}(n, \mathbb{C}) \cap U(n)$ by conjugation, the action is still transitive and the stabilizer of I_n is $O(n)$, so $\text{Sym}(n, \mathbb{C}) \cap U(n) \cong U(n)/O(n)$.

Therefore we can identify \mathcal{L} with $\text{Sym}(n, \mathbb{C}) \cap U(n)$.

13.1.2. $SU(n, n)$

This is quite similar to $\text{Sp}(2n, \mathbb{R})$. Explicitly,

$$SU(n, n) = \{g \in \text{SL}(2n, \mathbb{C}) : g^* I_{n,n} g = I_{n,n}\}, \quad \text{where} \quad I_{n,n} = \begin{pmatrix} I_n & 0 \\ 0 & -I_n \end{pmatrix}.$$

Consider domain $\mathcal{T} := \{Z \in \text{Mat}(n, \mathbb{C}) : Z^* Z < I_n\}$, then one can check that the linear fractional transformation $g(Z) = (AZ+B)(CZ+D)^{-1}$ for $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in SU(n, n)$ and $Z = X+iY \in \mathcal{T}$ is well-defined, which determines a transitive action.

The stabilizer of 0 is

$$\begin{aligned} \text{Stab}_{SU(n,n)}(0) &= \left\{ \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \in SU(n, n) \right\} \\ &= \left\{ \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} : \begin{matrix} A^* A = D^* D = I_n \\ \det(A) \det(D) = 1 \end{matrix} \right\} = S(U(n) \times U(n)) \end{aligned}$$

which is maximal compact.

The center $Z(S(U(n) \times U(n))) = \left\{ \begin{pmatrix} \lambda I_n & 0 \\ 0 & \mu I_n \end{pmatrix} : \begin{matrix} |\lambda| = |\mu| = 1 \\ (\lambda\mu)^n = 1 \end{matrix} \right\}$ is non-trivial.

If $Z \in \mathcal{T}$, then $I - Z$ is invertible, the inverse Cayley transform

$$\begin{aligned} c^{-1} : \mathcal{T} &\longrightarrow \text{Herm}(n, \mathbb{C}) + i \text{Herm}^+(n, \mathbb{C}) \\ Z &\longmapsto i \frac{I + Z}{I - Z} \end{aligned}$$

is biholomorphic (verify: $\frac{1}{2i} [c(Z) - c(Z)^*] = (I - Z^*)^{-1} [I - Z^* Z] (I - Z)^{-1} \in \text{Herm}^+(n, \mathbb{C})$), where $\text{Herm}(n, \mathbb{C})$ is the (real) vector space of Hermitian matrices and $\text{Herm}^+(n, \mathbb{C})$ is the (real) cone of positive definite ones.

Therefore $SU(n, n)/S(U(n) \times U(n)) \cong \mathcal{T} \cong \text{Herm}(n, \mathbb{C}) + i \text{Herm}^+(n, \mathbb{C})$ is a Hermitian symmetric space of tube type.

The Shilov boundary of \mathcal{T} is

$$\check{S}_{n,n} = \{Z \in M_n(\mathbb{C}) : I_n - Z^* Z = 0\}$$

which corresponds to the space of maximal isotropic subspaces

$$\text{Iso}_{n,n} = \{L \in \text{Gr}_n(\mathbb{C}^{2n}) : |\omega|_L = 0\} \subset \overline{\mathcal{X}_{n,n}}$$

under the Borel embedding $\mathcal{X}_{n,n} = \{L \in \text{Gr}_n(\mathbb{C}^{2n}) : |\omega|_L > 0\} \subset \text{Gr}_n(\mathbb{C}^{2n})$

Remark 13.6. A general description of Borel embedding, Harish-Chandra embedding and the Shilov boundaries in each realizations can be found in [Wie04].

13. Positivity of Lie Groups of Hermitian Type

13.1.3. $\text{SO}(n, 2)$ ($n > 2$)

Consider the domain (also known as the *Lie ball*):

$$\mathcal{D}_{IV_n} := \left\{ Z \in \mathbb{C}^n : Z^* Z < \frac{1}{2} \left(1 + |{}^t Z Z|^2 \right) < 1 \right\} = \left\{ z \in \mathbb{C}^n : \sum |z_i|^2 < \frac{1}{2} \left(1 + \left| \sum z_i^2 \right|^2 \right) < 1 \right\}$$

then one can check that linear fractional transformation

$$\text{SO}(n, 2) \times \mathcal{D}_{IV_n} \longrightarrow \mathcal{D}_{IV_n}$$

$$\left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, Z \right) \mapsto \frac{AZ + B \begin{pmatrix} \frac{1}{2}({}^t Z Z + 1) \\ \frac{i}{2}({}^t Z Z - 1) \end{pmatrix}}{(1, i) \cdot \left(CZ + D \begin{pmatrix} \frac{1}{2}({}^t Z Z + 1) \\ \frac{i}{2}({}^t Z Z - 1) \end{pmatrix} \right)}$$

is a transitive action, which can be extended continuously to $\overline{\mathcal{D}_{IV_n}}$. The stabilizer of 0 is easily verified to be $\text{SO}(n) \times \text{SO}(2)$, which has non-trivial center.

To describe a tube domain expression of \mathcal{D}_{IV_n} , first we introduce the *future tube* in \mathbb{C}^{n+1} ($n \geq 0$):

$$\tau^+(n) = \mathbb{R}^{n+1} + iV^+ = \{z \in \mathbb{C}^{n+1} : y^2 = y_0^2 - y_1^2 - \dots - y_n^2 > 0, y_0 > 0\}.$$

where the sharp convex subset

$$V^+ = V^+(n) = \{y \in \mathbb{R}^{1,n} : y^2 > 0, y_0 > 0\}.$$

is usually called the *future cone*.

The reason of its name is that for $n = 3$, $\mathbb{R}^{1,3}$ with the Lorentz inner product becomes the Minkowski spacetime. The boundary $\partial\tau^+(3)$ consists of the smooth part

$$S = \{\zeta = \xi + i\eta \in \mathbb{C}^4 : \eta^2 = 0, \eta_0 > 0\}$$

and the distinguished boundary

$$M = \{\zeta = \xi + i\eta \in \mathbb{C}^4 : \eta = 0\} \cong \mathbb{R}^4,$$

Through any point $\zeta \in S$ there passes a generator l_ζ of the cone

$$\Gamma^+ = \partial V^+ = \{\eta \in \mathbb{R}^{1,3} : \eta^2 = 0, \eta_0 \geq 0\}$$

called a *real light ray*. The complexification λ_ζ of the ray l_ζ which coincides with a complex halfplane $\lambda_\zeta = \{\xi + \alpha\eta : \alpha \in \mathbb{C}, \text{Im } \alpha > 0\}$ is called a *complex light ray*.

We will give a biholomorphism between the n -dimensional Lie ball \mathcal{D}_{IV_n} and the future tube $\tau^+(n-1)$. This biholomorphism is a composition of two mappings.

The first mapping is a realization of $\tau^+(n-1)$ as a domain on a complex quadric in $\mathbb{C}\mathbb{P}^{n+1}$. Let us introduce the new variables

$$z_1 = \frac{s_1}{s_0}, \dots, z_{n-1} = \frac{s_{n-1}}{s_0}, z_0 = \frac{s_n}{s_0}.$$

In these variables the domain $\tau(n-1) = \{z \in (\mathbb{R}^{1,n-1})^2 : y^2 > 0\}$ will transform to the domain

$$T' = \left\{ s \in \mathbb{C}^{n+2} : -|s_0|^2 - \dots - |s_{n-1}|^2 + |s_n|^2 + 2 \text{Re}(\bar{s}_0 s_{n+1}) > 0, \right. \\ \left. -s_0^2 - \dots - s_{n-1}^2 + s_n^2 + 2s_0 s_{n+1} = 0 \right\}$$

Changing the variables s_0, s_1, \dots, s_{n+1} to the variables $t_0 = s_0 - s_{n+1}, t_1 = s_1, \dots, t_{n+1} = s_{n+1}$ we can write T' in the form

$$T = \left\{ t \in \mathbb{C}^{n+2} : -|t_0|^2 - \dots - |t_{n-1}|^2 + |t_n|^2 + |t_{n+1}|^2 > 0, \right. \\ \left. -t_0^2 - \dots - t_{n-1}^2 + t_n^2 + t_{n+1}^2 = 0 \right\}$$

which is a section of the domain $\tilde{T} = \left\{ t \in \mathbb{C}^{n+2} : |t_0|^2 + \dots + |t_{n-1}|^2 < |t_n|^2 + |t_{n+1}|^2 \right\}$ by the complex quadric $\{t_0^2 + \dots + t_{n-1}^2 = t_n^2 + t_{n+1}^2\}$. The domains T and \tilde{T} are given by homogeneous relations so it is more natural to consider them as domains in $\mathbb{C}\mathbb{P}^{n+1}$. Note that the Levi form of the domain \tilde{T} being restricted to the complex tangent space of $\partial\tilde{T}$ at a point t with $t_{n+1} \neq 0$ has one negative and n positive eigenvalues. The domain T has two components distinguished by the sign of $\text{Im} \frac{t_n}{t_{n+1}}$. $\tau^+(n-1)$ is biholomorphic to the domain T_+ on the quadric in $\mathbb{C}\mathbb{P}^{n+1}$ given in homogeneous coordinates as follows

$$T_+ = \left\{ [t_0 : t_1 : \dots : t_{n+1}] : |t_0|^2 + \dots + |t_{n-1}|^2 < |t_n|^2 + |t_{n+1}|^2, \right. \\ \left. t_0^2 + \dots + t_{n-1}^2 = t_n^2 + t_{n+1}^2, \text{Im} \frac{t_n}{t_{n+1}} > 0 \right\}$$

The second mapping given by

$$w_0 = \frac{t_0}{t_n + it_{n+1}}, \dots, w_{n-1} = \frac{t_{n-1}}{t_n + it_{n+1}}$$

transforms the domain T_+ biholomorphically onto the domain \mathcal{D}_{IV_n} .

The composed mapping of $\tau^+(n-1)$ onto \mathcal{D}_{IV_n} is given by

$$w_0 = i \frac{1 + z^2}{(z + \mathbf{i})^2}, w_1 = i \frac{2z_1}{(z + \mathbf{i})^2}, \dots, w_{n-1} = i \frac{2z_{n-1}}{(z + \mathbf{i})^2}$$

where $\mathbf{i} = (i, 0, \dots, 0)$. We conclude that $\text{SO}(n, 2)/(\text{SO}(n) \times \text{SO}(2)) \cong \mathcal{D}_{IV_n} \cong \tau^+(n-1)$ is a Hermitian symmetric space of tube type.

The distinguished boundary of $\tau^+(n-1)$ transforms into the set

$$S_L = \left\{ |w_0|^2 + \dots + |w_{n-1}|^2 = 1, |w_0^2 + \dots + w_{n-1}^2| = 1 \right\}.$$

Set $w = u + iv, (z, \omega) = z_0\omega_0 + z_1\omega_1 + \dots + z_{n-1}\omega_{n-1}$. Then the intersection of S_L with the complex sphere $\Sigma_1 = \{w \in \mathbb{C}^n \mid (w, w) = 1\}$ is given by the equations $|u|^2 - |v|^2 = 1, (u, v) = 0, |u|^2 + |v|^2 = 1$. It follows that $v = 0$; hence Σ_1 intersects S_L in the $(n-1)$ -dimensional real sphere $\{u \in \mathbb{R}^n \mid |u|^2 = 1\}$. So S_L can be written as

$$\check{S} := S_L = \left\{ e^{i\theta} u \mid u \in S^{n-1} \subset \mathbb{R}^n \right\} \simeq S^1 \times S^{n-1} / \mathbb{Z}_2,$$

which is also the (distinguished) Shilov boundary of \mathcal{D}_{IV_n} , known as the *Lie sphere*.

Remark 13.7. For the convenience of readers, we record at here the classification of all irreducible Hermitian symmetric spaces, according as to whether or not they are of tube type:

tube type	nontube type
$\text{SU}(n, n)$	$\text{SU}(p, q), p > q$
$\text{Sp}(2n, \mathbb{R})$	
$\text{SO}^*(2n), n \text{ even}$	$\text{SO}^*(2n), n \text{ odd}$
$\text{SO}(n, 2)$	
$\text{E}_7(-25)$	$\text{E}_6(-14)$

where $\text{E}_7(-25)$ and $\text{E}_6(-14)$ correspond to the exceptional Hermitian symmetric spaces of complex dimension 27 and 16, respectively.

13. Positivity of Lie Groups of Hermitian Type

13.2. Maslov index and a semigroup

For $\mathrm{Sp}(2n, \mathbb{R})$, consider

$$V = \left\{ g \in \mathrm{Sp}(2n, \mathbb{R}) : g = \begin{pmatrix} I_n & 0 \\ M & I_n \end{pmatrix}, M \in \mathrm{Sym}(n, \mathbb{R}) \right\}$$

$$W = \left\{ g \in \mathrm{Sp}(2n, \mathbb{R}) : g = \begin{pmatrix} I_n & N \\ 0 & I_n \end{pmatrix}, N \in \mathrm{Sym}(n, \mathbb{R}) \right\}$$

and

$$H = \left\{ g \in \mathrm{Sp}(2n, \mathbb{R}) : g = \begin{pmatrix} A & 0 \\ 0 & {}^t A^{-1} \end{pmatrix} \right\} \cong \mathrm{GL}(n, \mathbb{R})$$

where the matrices are written with respect to a symplectic basis. Define

$$\mathrm{Sp}(2n, \mathbb{R})^{>0} := V^{>0} H^\circ W^{>0}$$

where

$$V^{>0} = \left\{ \begin{pmatrix} I_n & 0 \\ M & I_n \end{pmatrix} \in V : M \in \mathrm{Pos}(n, \mathbb{R}) \right\}, W^{>0} = \left\{ \begin{pmatrix} I_n & N \\ 0 & I_n \end{pmatrix} \in W : N \in \mathrm{Pos}(n, \mathbb{R}) \right\}$$

and H° is the identity component of H . To show $\mathrm{Sp}(2n, \mathbb{R})^{>0}$ is a subsemigroup, we only need to verify the product of an element in $W^{>0}$ with an element in $V^{>0}$ is in $V^{>0} H^\circ W^{>0}$. Notice if $N, M \in \mathrm{Pos}(n, \mathbb{R})$, then $I + NM$ is invertible since if it had a nonzero nullvector v , then $Mv = -N^{-1}v$, then ${}^t v M v = -{}^t v N^{-1} v$, a contradiction. An explicit calculation (in view of block Gaussian elimination) is

$$\begin{aligned} & \begin{pmatrix} I & N \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ M & I \end{pmatrix} = \begin{pmatrix} I + NM & N \\ M & I \end{pmatrix} \\ & = \begin{pmatrix} I & 0 \\ M(I + NM)^{-1} & I \end{pmatrix} \begin{pmatrix} I + NM & 0 \\ 0 & I - M(I + NM)^{-1}N \end{pmatrix} \begin{pmatrix} I & (I + NM)^{-1}N \\ 0 & I \end{pmatrix}. \end{aligned}$$

Remark 13.8. The definition for $\mathrm{Sp}(2n, \mathbb{R})^{>0}$ is obtained from Θ -positive structure by taking $\Theta = \{\alpha_n\}$, where the restricted roots $\alpha_i = e_i - e_{i+1}$, $\alpha_n = 2e_n$.

We will see that a triple $(L_1, L_2, L_3) \in \mathcal{L}^3$ is positive if and only if $\exists g \in \mathrm{Sp}(2n, \mathbb{R})$ such that $g(L_1, L_2, L_3) = (L_E, uL_E, L_F)$ for some $u \in W^{>0}$ if and only if (L_1, L_2, L_3) has maximal Maslov index n , where $L_E = \mathrm{span}\{e_1, \dots, e_n\}$, $L_F = \mathrm{span}\{f_1, \dots, f_n\}$ are two standard Lagrangians, $\{f_1, \dots, f_n, e_1, \dots, e_n\}$ is a symplectic basis with respect to ω .

Now we generalize this construction to any Hermitian Lie group G which is of tube type.

First we briefly introduce the notion of Jordan algebras and their relationship with symmetric cones. For details, a good reference is [FK94].

Definition 13.9. A *Jordan algebra* is a vector space V with a bilinear (not necessarily associative) product $V \times V \rightarrow V, (x, y) \mapsto xy$ with

$$yx = xy$$

$$x(x^2y) = x^2(xy).$$

We always assume that V has an identity element e .

An *idempotent* in V is an element $c \in V$ satisfying $c^2 = c$. Two idempotents c and d are said to be *orthogonal* if $cd = 0$. Since then

$$\langle c, d \rangle = \langle c^2, d \rangle = \langle c, cd \rangle = \langle c, 0 \rangle = 0,$$

orthogonal idempotents are orthogonal with respect to the inner product.

Remarks 13.10. (1) $\forall x \in V$, we usually denote by $L(x) : V \rightarrow V$ the multiplication by x .

(2) Let c be an idempotent in a Jordan algebra. It is not hard to verify that the only possible eigenvalues of $L(c)$ are $0, \frac{1}{2}$ and 1 . (verify the identity $2L(c)^3 - 3L(c)^2 + L(c) = 0$)

A Jordan algebra over \mathbb{R} is said to be *Euclidean* if there exists a positive definite symmetric bilinear form $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ which is associative; i.e. $\langle xu, y \rangle = \langle u, xy \rangle$ for all $x, y, u \in V$.

An idempotent is called *primitive* if it is non-zero and cannot be written as sum of two (necessarily orthogonal) non-zero idempotents. A complete system of orthogonal primitive idempotents or a *Jordan frame*, is a set of primitive idempotents c_1, \dots, c_m satisfying

$$c_i c_j = 0, \quad \forall i \neq j, \quad \sum_{i=1}^m c_i = e.$$

Let V be a finite-dimensional Euclidean Jordan algebra over \mathbb{R} , we define the rank of V as

$$\text{rk}(V) := \max\{\deg(x) \mid x \in V\},$$

where $\deg(x)$ is the degree of the unique minimal polynomial of x . An element $x \in V$ is called *regular* if $\deg(x) = \text{rk}(V)$.

Proposition 13.11 (Characteristic polynomial). *The set of regular elements is open and dense in V . There exist polynomials a_1, \dots, a_r on V such that the minimal polynomial of every regular element $x \in V$ in the variable λ is given by*

$$f(\lambda, x) = \lambda^r - a_1(x)\lambda^{r-1} + \dots + (-1)^r a_r(x).$$

The polynomials a_i are unique and homogeneous of degree i .

Another useful result is

Theorem 13.12 (Spectral decomposition of type II). *Suppose V has rank r . Then for every x in V there exists a Jordan frame c_1, \dots, c_r and real numbers $\lambda_1, \dots, \lambda_r$ such that*

$$x = \sum_{j=1}^r \lambda_j c_j.$$

The numbers λ_j (with their multiplicities) are uniquely determined by x . Moreover,

$$a_k(x) = \sum_{1 \leq i_1 \leq \dots \leq i_k \leq r} \lambda_{i_1} \dots \lambda_{i_k},$$

where a_k ($1 \leq k \leq r$) is the polynomial defined in the above proposition.

13. Positivity of Lie Groups of Hermitian Type

The reason why we choose Jordan algebras is the cones of squares give a full characterization of symmetric cones that we now describe.

Let V be a Euclidean Jordan \mathbb{R} -algebra and let \mathcal{K} be the set of all squares, i.e.,

$$\mathcal{K} := \{x^2 : x \in V\}.$$

If $x^2 \in \mathcal{K}$ and $\alpha \geq 0$ then $\alpha x^2 = (\sqrt{\alpha}x)^2 \in \mathcal{K}$. It follows that \mathcal{K} is a cone. We call \mathcal{K} the *cone of squares* of V . Furthermore, it can be shown that \mathcal{K} is a closed symmetric cone, cf. [FK94]. (i.e. homogeneous and self-dual; $\text{Aut}(\mathcal{K})$ transitively on the interior of \mathcal{K} , $\mathcal{K} = \mathcal{K}^*$)

Remark 13.13. *Since \mathcal{K} is closed, here \mathcal{K}^* is the dual closed cone $\mathcal{K}^* := \{y \in V \mid \langle x, y \rangle \geq 0, \forall x \in \mathcal{K}\}$. If \mathcal{C} is an open convex cone in V , we say that \mathcal{C} is homogeneous if $\text{Aut}(\mathcal{C})$ transitively on \mathcal{C} ; is self-dual if $\mathcal{C} = \mathcal{C}^*$, where \mathcal{C}^* is the dual open cone $\mathcal{C}^* := \{y \in V \mid \langle x, y \rangle > 0, \forall x \in \mathcal{C} \setminus \{0\}\}$.*

Proposition 13.14. *Let \mathcal{I} be the set of invertible elements in V , then $\mathcal{K}^\circ = \{x^2 \mid x \in \mathcal{I}\}$, where \mathcal{K}° is the interior of \mathcal{K} .*

Proof. $y \in \mathcal{K}^*$ if and only if $\langle y, x^2 \rangle = \langle L(y)x, x \rangle \geq 0$ for all $x \in V$, if and only if $L(y)$ is positive semidefinite. Since $\mathcal{K} = \mathcal{K}^*$, we need to verify that $L(y)$ is positive definite if and only if $y = w^2$ for some invertible w .

Necessity: Suppose $L(y) > 0$, and $y = \sum_{i=1}^r \lambda_i c_i$ is the spectral decomposition. Since

$$\langle L(y)c_i, c_i \rangle = \lambda_i \langle c_i, c_i \rangle = \lambda_i \|c_i\|^2 > 0,$$

we know $\lambda_i > 0$. Therefore y is invertible and $y = w^2$ with $w = \sum_{i=1}^r \sqrt{\lambda_i} c_i$.

Sufficiency: Suppose $y = w^2$ for invertible $w = \sum_{i=1}^r \lambda_i c_i$, notice that we have $\langle zL(w^2), z \rangle = \langle \sum_{i=1}^r \lambda_i^2 L(c_i)z, z \rangle$. Since w is invertible, the eigenvalues of w^2 are greater than zero, i.e. $\lambda_i > 0$ for all i . Hence we only need to verify for each $z \in V \setminus \{0\}$, there is some i such that $\langle L(c_i)z, z \rangle > 0$. Suppose not, $\exists z_0 \in V \setminus \{0\}$ s.t. $\langle L(c_i)z_0, z_0 \rangle = 0$ for all i . (remark that $L(c_i) \geq 0$) Therefore

$$0 = \sum_{i=1}^r \langle L(c_i)z_0, z_0 \rangle = \sum_{i=1}^r \langle c_i z_0, z_0 \rangle = \langle e z_0, z_0 \rangle = \langle z_0, z_0 \rangle,$$

which is a contradiction since $z_0 \neq 0$. □

From this relation one can also prove that \mathcal{K}° is an open symmetric cone. We have mentioned that \mathcal{K} is a closed symmetric cone; in fact, the converse also holds. We state the following result which is a Jordan algebraic characterization of symmetric cones.

Theorem 13.15 (Theorem III.3.1 in [FK94]). *A cone is symmetric if and only if it is the cone of squares of some Euclidean Jordan algebra.*

Proposition 13.16. *Let V be an Euclidean Jordan \mathbb{R} -algebra with rank r and \mathcal{K} its cone of squares. If $x \in V$ is such that $x = \sum_{i=1}^r \lambda_i c_i$, where $(c_i)_{i=1, \dots, r}$ is a Jordan frame, then $\lambda_i \geq 0$, resp. > 0 . for $i = 1, \dots, r$ if and only if $x \in \mathcal{K}$, resp. \mathcal{K}° .*

Proof. If $\lambda_i \geq 0$ then we can write $x = y^2$, with $y = \sum_{i=1}^r \sqrt{\lambda_i} c_i$, which means that $x \in \mathcal{K}$. In case that $\lambda_i > 0$ for $i = 1, \dots, r$, we know that x is invertible, and it follows that $x \in \mathcal{K}^\circ$. Conversely, if $x \in \mathcal{K}$ we have $x = y^2$, with $y \in V$. If we denote α_i the eigenvalues of y , we can write $x = \sum_{i=1}^r \alpha_i^2 c_i$ where α_i^2 's are the eigenvalues of x , which are greater or equal than zero. If $x \in \mathcal{K}^\circ$ then y is invertible, therefore α_i^2 's are greater than zero. □

We denote \mathcal{K}° by Ω , tube domain $T_\Omega := V \oplus i\Omega = \{\sum \mu_i c_i \mid \text{Im } \mu_i > 0\}$. Via the Cayley transform $c(z) = (z - ie)(z + ie)^{-1}$, T_Ω is biholomorphic to bounded symmetric domain in $V^\mathbb{C}$:

$$\mathcal{D} := \left\{ v = \sum \lambda_i c_i \in V^\mathbb{C} : (c_i) \text{ Jordan frame, } |\lambda_i| < 1 \right\}.$$

The Shilov boundary of T_Ω is V . (Proposition IX.5.5 in [FK94]) For the Shilov boundary \check{S} of \mathcal{D} , we have:

Proposition 13.17 (Proposition.X.2.3, Theorem.X.4.6 in [FK94]). *Let $\mathcal{D} \subset V^\mathbb{C}$ be a bounded symmetric domain in the complexification of an Euclidean Jordan algebra V . The following are equivalent:*

- (1) $z \in \check{S}$,
- (2) $z = \sum \lambda_i c_i$, where c_1, \dots, c_r is a Jordan frame and $|\lambda_i| = 1$,
- (3) $z \in \overline{c(V)}$,
- (4) $\bar{z} = z^{-1}$.

Example 13.18. For $\mathcal{X}_n = \text{Sym}(n, \mathbb{R}) + i \text{Pos}(n, \mathbb{R})$, define $X \circ Y := \frac{XY + YX}{2}$ for $X, Y \in \text{Sym}(n, \mathbb{R})$, where XY denotes the usual matrix product. The inner product is defined as $\langle X, Y \rangle = \text{Tr}(X \circ Y) = \text{Tr}(XY)$ for all $X, Y \in \text{Sym}(n, \mathbb{R})$. One can check that $(\text{Sym}(n, \mathbb{R}), \circ)$ is a Euclidean Jordan algebra. Any $X \in \text{Sym}(n, \mathbb{R})$ has spectral decomposition:

$$X = \sum_{i=1}^n \lambda_i q_i {}^t q_i$$

where $\lambda_i \in \mathbb{R}$, $i = 1, \dots, n$, are the eigenvalues of X and $q_i, i = 1, \dots, n$ are unitary eigenvectors of X . Each $q_i {}^t q_i$ is an idempotent; $\{q_1 {}^t q_1, \dots, q_n {}^t q_n\}$ is a Jordan frame.

$\text{Aut}(\text{Pos}(n, \mathbb{R})) = \text{GL}(n, \mathbb{R}) / \{\pm I_n\}$ acts on $\text{Sp}(2n, \mathbb{R})$ transitively by $g \cdot x := gx {}^t g$. For any $Y \in \text{Pos}(n, \mathbb{R})^*$, $\langle Y, \xi {}^t \xi \rangle > 0$ for all nonzero $\xi \in \mathbb{R}^n$. Namely, $\text{tr}(Y \xi {}^t \xi) = \sum_{i,j} y_{ij} \xi_i \xi_j > 0$, which implies $Y \in \text{Pos}(n, \mathbb{R})$. On the other hand, take any $X \in \text{Pos}(n, \mathbb{R})$, $Y \in \overline{\text{Pos}(n, \mathbb{R})} \setminus \{0\}$, we write $Y = \sum_{i=1}^n \mu_i q_i {}^t q_i$ for $\mu_i \geq 0, \forall i$ but not all 0. Then $\text{Tr}(XY) = \sum_i \mu_i {}^t q_i X q_i > 0$, hence $X \in \text{Pos}(n, \mathbb{R})^*$. This shows $\text{Pos}(n, \mathbb{R}) = \text{Pos}(n, \mathbb{R})^*$, therefore $\text{Pos}(n, \mathbb{R})$ is an open symmetric cone.

For the Maslov cocycle, we follow Anna Wienhard's definition in [Wie04], which is a slight variation of Clerc's definition [Cle04].

Let $T : X = G/K \rightarrow T_\Omega = V + i\Omega$ be a biholomorphism, where $\Omega \subset V$ is a symmetric convex cone in the real vector space V ; \mathcal{D} the bounded domain realization and \check{S} its Shilov boundary; r_X the rank of X . Then as mentioned in previous sections, \check{S} is a homogeneous G -space of the form G/Q , where Q is a specific parabolic subgroup (which is maximal if X is irreducible). Two points $x, y \in \check{S}$ are *transversal* if (x, y) lies in the open G -orbit in \check{S}^2 .

Let

$$\check{S}^{[3]} := \{(x_1, x_2, x_3) \in \check{S}^3 : x_i \text{ is transverse to } x_j, x_k \text{ for some } i\}$$

be the space of triples, where one point is transverse to the other two. For any triple $(x_1, x_2, x_3) \in \check{S}^{[3]}$, we may assume that $T(x_3) = \infty$ and $y_1 = T(x_1), y_2 = T(x_2) \in V$ by using the transitivity of the G -action on \check{S} , then the *Maslov cocycle* is defined to be

$$\tau(x_1, x_2, x_3) := k_+(y_2 - y_1) - k_-(y_2 - y_1),$$

where k_\pm are the numbers of positive respectively negative eigenvalues in the spectral decomposition of $(y_2 - y_1)$ with respect to a Jordan frame $(c_j)_{j=1, \dots, r_X}$.

13. Positivity of Lie Groups of Hermitian Type

Remark 13.19. Let $(x_1, x_2, x_3) \in \check{S}^{[3]}$ with $\tau(x_1, x_2, x_3) = r_X$, then (x_1, x_2, x_3) are pairwise transverse since if we assume x_3 is transverse to x_1, x_2 w.l.o.g., then $\tau = r_X$ implies eigenvalues of $y_2 - y_1$ are all positive; especially $y_2 - y_1$ is invertible.

Given any triple $x_1, x_2, x_3 \in \check{S}$, using G -action we may assume that x_1, x_2, x_3 are transverse to ∞ , hence y_1, y_2, y_3 in V . The cocycle identity allows us to define τ on \check{S}^3 as

$$\begin{aligned} \tau(x_1, x_2, x_3) &= \tau(x_1, x_2, \infty) + \tau(x_2, x_3, \infty) - \tau(x_1, x_3, \infty) \\ &= (k_+(y_2 - y_1) - k_-(y_2 - y_1)) \\ &\quad + (k_+(y_3 - y_2) - k_-(y_3 - y_2)) \\ &\quad - (k_+(y_3 - y_1) - k_-(y_3 - y_1)). \end{aligned}$$

This clearly defines a G -invariant real function on \check{S}^3 . We call a triple $(x_1, x_2, x_3) \in \check{S}^3$ *maximal* if $\tau(x_1, x_2, x_3)$ attains its maximal possible value $\text{rk}(V) = r_X$.

If we restrict ourselves to pairwise transverse triples (usually denoted by $\check{S}^{(3)}$), we will recover Clerc and Ørsted's definition of Maslov index given in [CØ01].

Under the Cayley transform c , $c(\infty) = e$, $c(0) = -e$, we also let

$$\varepsilon_k := \sum_{i=1}^k c_i - \sum_{i=k+1}^{r_X} c_i \in \check{S}$$

which is a point on \check{S} following from Proposition 13.17, where $(c_j)_{j=1, \dots, r_X}$ is a fixed Jordan frame. We have

$$\begin{aligned} c(\varepsilon_k) &= \left(\sum_{j=1}^k (1-i)c_j - \sum_{j=k+1}^{r_X} (1+i)c_j \right) \cdot \left(\sum_{j=1}^k (1+i)c_j - \sum_{j=k+1}^{r_X} (1-i)c_j \right)^{-1} \\ &= \left(\sum_{j=1}^k (1-i)c_j - \sum_{j=k+1}^{r_X} (1+i)c_j \right) \cdot \frac{1}{2} \left(\sum_{j=1}^k (1-i)c_j - \sum_{j=k+1}^{r_X} (1+i)c_j \right) = (-i)\varepsilon_k, \end{aligned}$$

hence $\tau(-e, (-i)\varepsilon_k, e) = k_+(\varepsilon_k) - k_-(\varepsilon_k) = 2k - r_X$.

Theorem 13.20 (Theorem 4.3 in [CØ01], Theorem 3.5 in [Cle04]). *There are exactly $r_X + 1$ orbits of pairwise transverse triples in \check{S}^3 under the action of G . Each $(-e, (-i)\varepsilon_j, e)$, $0 \leq j \leq r_X$, represents one orbit. The function τ takes values in $\{-r_X, -r_X + 2, \dots, r_X - 2, r_X\}$ and it classifies all these G -orbits.*

Therefore we can also define for any pairwise transverse triple (x_1, x_2, x_3) , $\tau(x_1, x_2, x_3) = 2k - r_X$, where k is the unique integer, $0 \leq k \leq r_X$, such that (x_1, x_2, x_3) is conjugate under G to the triplet $(-e, (-i)\varepsilon_k, e)$. See [CØ01].

Remarks 13.21. For tube type bounded symmetric domain \mathcal{D} , $\pi\tau$ actually coincides with the restriction of Bergmann cocycle β on $\check{S}^{(3)}$. For non-tube type \mathcal{D} , however, $\beta(\check{S}^{(3)}) = [-\pi r_X, \pi r_X]$. See [Wie04] Chapter 5 for details.

Fix $x, z \in \check{S} = G/Q$ such that $\text{Stab}_G(z) = Q$, $\text{Stab}_G(x) = Q^{opp}$. Given any $y \in G/Q$ transverse to z , there exists unique $u \in \text{UniRad}(Q)$ such that $y = ux$, where $\text{UniRad}(Q)$ is the unipotent radical of the parabolic Q .

Denote by

$$U^{>0} := \{u \in \text{UniRad}(Q) \mid \tau(x, ux, z) = r_X\},$$

$$U^{opp, >0} := \{v \in \text{UniRad}(Q^{opp}) \mid \tau(z, vz, x) = r_X\},$$

$$L^\circ := \text{identity component of } Q \cap Q^{opp}.$$

then we can define the positive subsemigroup $G^{>0} \subset G$ to be the subsemigroup generated by $U^{>0}$, $U^{opp, >0}$ and L° .

14. Positivity of Triples of Flags

by *Raphael Appenzeller, Francesco Fournier-Facio*

14.1. Flags

In this section we introduce the objects that we will be working with. We start with the flag variety, then consider the relevant (semi)groups, and finally put both together by letting the (semi)groups act on the flag variety.

14.1.1. The flag variety

Definition 14.1. A *flag* is an $(n-1)$ -tuple (F_1, \dots, F_{n-1}) , where F_i is a vector subspace of \mathbb{R}^n of dimension i , and $F_i \subset F_{i+1}$ for all $1 \leq i \leq (n-2)$.

The set of flags is denoted by $\text{Flag}(\mathbb{R}^n)$ and is called the *(full) flag variety*.

The two most basic examples of flags also turn out to be the most important ones:

Example 14.2. Let $\{e_i\}_{i=1}^n$ be the canonical basis of \mathbb{R}^n .

The *standard ascending flag* is $F = (F_1, \dots, F_{n-1})$, where $F_i = \langle e_1, \dots, e_i \rangle$.

The *standard descending flag* is $E = (E_1, \dots, E_{n-1})$, where $E_i = \langle e_n, \dots, e_{n-i+1} \rangle$.

In some sense, these two flags are *opposite* each other. One way to formalize this is the following:

Definition 14.3. Two flags F^1, F^2 are *transverse*, denoted $F^1 \pitchfork F^2$, if $F_i^1 \cap F_{n-i}^2 = \{0\}$ for every $1 \leq i < n$. Equivalently, F^1 and F^2 are transverse if $F_i^1 \oplus F_{n-i}^2 = \mathbb{R}^n$ for every $1 \leq i < n$.

Given a flag F , the set of flags transverse to F is denoted by Ω_F .

Example 14.4. The standard ascending and descending flags are transverse to each other.

The full flag variety $\text{Flag}(\mathbb{R}^n)$ comes equipped with a natural topology, induced by the inclusion in a product of Grassmannians $\text{Flag}(\mathbb{R}^n) \subset \text{Gr}(n, 1) \times \dots \times \text{Gr}(n, n-1)$.

Proposition 14.5. *Let F be the standard ascending flag. Then Ω_F is open and dense in F .*

Remark 14.6. *We will soon show that $\text{Flag}(\mathbb{R}^n)$ is a homogeneous space. Therefore the statement of the proposition holds for every flag, not only the standard ascending one.*

Proof idea. Let $F^1 \in \Omega_F$. If another flag F^2 is close enough to F^1 , then F_i^1 is close to F_i^2 , and so every element of $\mathbb{R}^n = F_i^1 + F_{n-i}$ is close to an element of $F_i^2 + F_{n-i}$, which is not possible if $\dim(F_i^2 + F_{n-i}) < n$. Thus $F^2 \in \Omega_F$, which shows that Ω_F is open.

Now let $F^1 \in \text{Flag}(\mathbb{R}^n)$ be arbitrary, then for every i we can modify F_i^1 by a small amount in a direction transverse to F_{n-i} in order to obtain a subspace F_i^2 such that $F_i^2 + F_{n-i} = \mathbb{R}^n$. It is possible to do this in a compatible way on every i by an induction argument, to put these spaces together into a flag F^2 , which is then an element of Ω_F close to F^1 . This shows that Ω_F is dense in $\text{Flag}(\mathbb{R}^n)$. \square

A further relevant notion is the following, which may be thought of as an analogue of transversality for triples of flags:

Definition 14.7. A triple of flags (F^1, F^2, F^3) is *generic* if for all $a, b, c \geq 0$ such that $a + b + c = n$, it holds $F_a^1 + F_b^2 + F_c^3 = \mathbb{R}^n$.

By choosing a, b or c to be equal to 0, we see that if (F^1, F^2, F^3) is generic, then the three flags are pairwise transverse. The converse is not true however (contrary to what is claimed in [BD14]), as the following example shows:

Example 14.8. Let F be the standard ascending, and E the standard descending flag. Let T be defined by $T_1 := \langle t_1 \rangle$ and $T_2 := \langle t_1, t_2 \rangle$, where

$$t_1 := \begin{pmatrix} y \\ z \\ 1 \end{pmatrix} \quad \text{and} \quad t_2 := \begin{pmatrix} x \\ 1 \\ 0 \end{pmatrix}$$

(it will be soon clear why we chose this notation). If $y \neq 0$ and $xz - y \neq 0$, then E, F and T are pairwise transverse. However, if $z = 0$, then the triple is not generic, since $E_1 + F_1 + T_1 = \langle e_1, e_3 \rangle$. As a concrete example, one can take $x = 1, y = 1$ and $z = 0$.

14.1.2. The (semi)groups

We will be working in the group $G := \text{GL}(n, \mathbb{R})$ and the main players will be its subgroups B of upper-triangular matrices, and U of unipotent upper-triangular matrices. We further denote by tB and tU the subgroups of lower-triangular and unipotent lower-triangular matrices, and $A := B \cap {}^tB$ the group of diagonal matrices.

Recall from the previous talk that a matrix $g \in G$ is called *totally positive* if each of its minors is positive. We denote by $G_{>0}$ the subset of totally positive matrices. Especially relevant for this talk is the subset $U_{>0}$: this is defined as the set of matrices in U such that every minor (that is not forced to be zero) is positive.

Example 14.9. Suppose that $n = 3$. Then each $u \in U$ is of the form

$$\begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix}$$

where $a, b, c \in \mathbb{R}$. Most minors are already determined: for instance the $(1, 3)$ minor is 0, and the $(1, 1)$ minor is 1. The condition that $u \in U_{>0}$ then amounts to:

$$x > 0, \quad y > 0, \quad z > 0, \quad xz - y > 0.$$

14. Positivity of Triples of Flags

We remark the following fact:

Lemma 14.10. $G_{>0}$ and $U_{>0}$ are subsemigroups.

Proof. This follows easily from the Cauchy–Binet Formula (see Proposition 11.5 and its proof). \square

14.1.3. The action

There is a natural action of G on $\text{Flag}(\mathbb{R}^n)$, given by $g \cdot (F_1, \dots, F_{n-1}) = (gF_1, \dots, gF_{n-1})$. The following result gives the important properties of this action that will be used in the definition of positivity.

Proposition 14.11. *The action of G on $\text{Flag}(\mathbb{R}^n)$ has the following properties:*

- (1) (Naturality) *The action is continuous and preserves the notions of transversality and genericity.*
- (2) (Transitivity) *The action of G on $\text{Flag}(\mathbb{R}^n)$ is transitive, and the action of G on pairs of transverse flags is transitive.*
- (3) (Stabilizers) *Let F be the standard ascending flag. Then the stabilizer of F is the Borel subgroup B , and the induced action of U on Ω_F is simply transitive.*

Proof. The first item is clear from the definitions. For transitivity, let T be a flag, and choose elements t_1, \dots, t_{n-1} such that $T_i = \langle t_1, \dots, t_i \rangle$. Then the element $g \in G$ whose i -th column is t_i sends the standard ascending flag F to T . Now the statement on transitivity on pairs of transverse flags will follow from the third item.

It is easy to see that B is the stabilizer of F . Next, we prove that U acts transitively on Ω_F . We prove this by induction on n . For $n = 1$ is clear. Now let $n > 1$ and assume that the statement is true up to $(n - 1)$. Let $T \in \Omega_F$: we need to show that there exists $u \in U$ such that $uE = T$, where E is the standard descending flag.

Let $v \in \mathbb{R}^n$ be a vector such that $T_1 = \langle v \rangle$. Since $T \pitchfork F$, it holds $T_1 + F_{n-1} = \mathbb{R}^n$, so the last coordinate of v is non-zero. Therefore, up to rescaling, we may assume that the last coordinate of v is 1. This allows to consider a matrix $u \in U$ whose last column is v . In other words $uE_1 = u\langle e_n \rangle = \langle v \rangle = T_1$. Since no condition has been imposed on the first $(n - 1)$ columns of u , by induction we may choose them so that $uE_i = T_i$ for every other i . Thus $uE = T$.

It remains to show that the action is simply transitive. Since we already know it is transitive, it suffices to show that the stabilizer in U of the standard descending flag E is trivial. Now the stabilizer in G of E is just tB , so the intersection of the two stabilizers is $B \cap {}^tB$, that is, the subgroup A of diagonal matrices. Since U intersects A trivially, we conclude. \square

14.2. Positivity of triples of flags via total positivity of groups

From the properties of the action in Proposition 14.11 we deduce that we can identify U with the set of flags transverse to F via the orbit of E :

$$\begin{aligned} U &\cong \Omega_F \\ u &\mapsto uE \end{aligned}$$

This allows us to give the following definition:

Definition 14.12. Let T be a flag. The triple (E, T, F) is called *positive (relative to E, F)* if there exists $u \in U_{>0}$ such that $T = uE$.

Remark 14.13. We follow the convention of Guichard-Wienhard [GW18]. Some authors [BD14, BD17, FG06] permute the order and say (E, F, T) is positive.

Example 14.14. We consider flags in \mathbb{R}^3 . Let

$$u = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{then} \quad uE = \left\{ \left\langle \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\rangle, \left\langle \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} \right\rangle \right\}.$$

Using the criterion from Example 14.9 it is easy to see that u is totally positive, and thus (E, uE, F) is a positive triple of flags.

We recommend to solve the following exercise now.

Exercise 14.1. Consider the following flag in \mathbb{R}^3

$$T = \left(\left\langle \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \right\rangle, \left\langle \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix} \right\rangle \right).$$

(a) Is there a $u \in U$ such that $(E, T, F) = (E, uE, F)$?

(b) Is there a $u \in U_{>0}$ such that $(E, T, F) = (E, uE, F)$?

The solution to (a) is

$$u = \begin{pmatrix} 1 & -2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}$$

which has some negative entries, thus $u \notin U_{>0}$. Remember that U acts on Ω_F simply transitively and thus the element $u \in U$ with $uE = T$ is unique, answering (b) negatively.

If (E, T, F) is positive, then $F \pitchfork T$ (because the action preserves transversality: Proposition 14.11) and the element $u \in U_{>0}$ such that $uF = T$ is unique. Moreover, E, F and T are pairwise transverse. Something stronger is true:

Lemma 14.15. Let (E, T, F) be a positive triple (relative to E, F). Then (E, T, F) is generic.

For the proof, we introduce the following notation, which will also be used in the next section:

14. Positivity of Triples of Flags

Notation 14.16. Let $g \in G$, and let $a, b, c \geq 0$ be such that $a + b + c = n$. Split g as a block matrix as follows.

$$g = \left(\begin{array}{c|c|c} (a \times a) & & \\ \hline & & (c \times c) \\ \hline & (b \times b) & \end{array} \right)$$

We denote by $g(a, b, c)$ the highlighted $(c \times c)$ block.

While this may seem a little arbitrary, it will be clear through the course of the proof how these blocks - and their minors - appear naturally when considering triples of the form (E, uE, F) .

Proof. Let $u \in U_{>0}$ be the unique element such that $T = uE$. We need to show that for every $a, b, c \geq 0$ such that $a + b + c = n$, it holds $F_a + E_b + T_c = \mathbb{R}^n$. Now we have $F_a = \langle e_1, \dots, e_a \rangle$, $E_b := \langle e_n, \dots, e_{n-b+1} \rangle$, and T_c is the span of the last c columns of the matrix u . Since u is totally positive, $u(a, b, c)$ has positive determinant. In particular, the last c columns of u span $\langle e_{a+1}, \dots, e_{n-b} \rangle$ modulo $F_a + E_b$. We conclude that $F_a + E_b + T_c = \mathbb{R}^n$. \square

Remark 14.17. Therefore the set $\{T \in \text{Flag}(\mathbb{R}^n) : (E, T, F) \text{ is positive}\}$ is a subset of $\Omega_E \cap \Omega_F$. By Proposition 14.11, it can be identified with $U_{>0}$, and in particular it is connected. In fact, Lusztig even proved that it is a connected component of $\Omega_E \cap \Omega_F$ [Lus94]. This is not very relevant for our purposes so we would not get into detail.

We can extend the definition from triples of this form to all triples by exploiting the high transitivity of the action.

Definition 14.18. A triple of flags (F^1, F^2, F^3) is (GW) -positive, if there exists $g \in G$ such that $(gF^1, gF^2, gF^3) = (E, uE, F)$ with $u \in U_{>0}$. This is the definition used in [GW18].

One should be careful to notice that while (E, T, F) is positive relative to E, F implies that it is (GW) -positive, the converse does not hold, as the following exercise shows.

Exercise 14.2. Show that (E, T, F) from Exercise 14.1 is (GW) -positive.

One can check (E, T, F) is (GW) -positive using $g = \begin{pmatrix} -1 & & \\ & 1 & \\ & & -1 \end{pmatrix}$. In fact we have the following characterisation.

Lemma 14.19. Let $u \in U$. Then (E, uE, F) is (GW) -positive if and only if there exists a diagonal matrix d such that $dud^{-1} \in U_{>0}$.

Proof. If the triple (E, uE, F) is (GW) -positive, then there exists $d \in G$ such that $(dE, duE, dF) = (E, vE, F)$, where $v \in U_{>0}$. This implies that d stabilizes both E and F , and therefore it must be a diagonal matrix. Moreover, since d stabilizes both E and F , it is also true that $dud^{-1}E = vE$. Now it is easy to see that $dud^{-1} \in U$, and so by simple transitivity (Proposition 14.11), it follows that $dud^{-1} = v \in U_{>0}$.

Conversely, if $dud^{-1} \in U_{>0}$, then $d \cdot (E, uE, F) = (dE, duE, dF) = (E, dud^{-1}E, F)$, which is positive relative to E, F . \square

Example 14.20. (Triples on the circle) In \mathbb{R}^2 , flags consist only of one subspace, and thus the flag variety can be identified with the projective line, which is homeomorphic to a circle. Note that every triple consisting of three distinct points is (GW) -positive, while only some triples are positive relative to E, F , namely those which are positively oriented in the usual sense.

14.3. Positivity via triple ratios

The notion of positivity introduced in the previous section has one disadvantage, namely that the standard ascending and descending flags seem to play a central role, even though the definition applies to all flags. In this section we introduce a set of *coordinates* that parametrizes the G -orbits on generic triples of flags, and allows to give a definition of positivity that is more intrinsic. In this section, we follow [BD14].

Let (E, F, T) be a generic triple of flags. For $0 \leq i \leq n$, fix a generator $e^{(i)} \in \Lambda^i(E_i) \cong \mathbb{R}$, which we also see as an element of $\Lambda^i(\mathbb{R}^n)$. Similarly, fix generators $f^{(i)}$ and $t^{(i)}$ of $\Lambda^i(F_i)$ and $\Lambda^i(T_i)$. Now if $a + b + c = n$, then $e^{(a)} \wedge f^{(b)} \wedge t^{(c)} \in \Lambda^n(\mathbb{R}^n) \cong \mathbb{R}$. Note that since the triple is generic, this element is never 0.

Definition 14.21. Fix an identification of $\Lambda^n(\mathbb{R}^n)$ with \mathbb{R} . Let $a, b, c \geq 1$ be such that $a + b + c = n$. Then we define the (a, b, c) -triple ratio of the triple (E, F, T) as follows:

$$T_{abc}(E, F, T) := \frac{e^{(a+1)} \wedge f^{(b)} \wedge t^{(c-1)}}{e^{(a-1)} \wedge f^{(b)} \wedge t^{(c+1)}} \cdot \frac{e^{(a)} \wedge f^{(b-1)} \wedge t^{(c+1)}}{e^{(a)} \wedge f^{(b+1)} \wedge t^{(c-1)}} \cdot \frac{e^{(a-1)} \wedge f^{(b+1)} \wedge t^{(c)}}{e^{(a+1)} \wedge f^{(b-1)} \wedge t^{(c)}}.$$

Note that each $e^{(i)}$ appears exactly once on the numerator and exactly once on the denominator (if it does appear). Therefore the definition of $T_{abc}(E, F, T)$ is independent of the choice of the generators $e^{(i)}$. The same holds of course for the choices of $f^{(i)}$ and $t^{(i)}$. It is a little less clear that is independent of the choice of the identification $\Lambda^n(\mathbb{R}^n) \cong \mathbb{R}^n$:

Lemma 14.22. For all $\gamma \in G$, it holds $T_{abc}(E, F, T) = T_{abc}(\gamma E, \gamma F, \gamma T)$. In particular, $T_{abc}(E, F, T)$ is independent of the choice of the identification $\Lambda^n(\mathbb{R}^n) \cong \mathbb{R}^n$.

Proof. We choose $\gamma \cdot e^{(i)}$ as a generator of $\Lambda^i(\gamma E_i)$, which we are allowed to do by the previous remark. Then for every $a, b, c \geq 0$ such that $a + b + c = n$ it holds

$$\gamma \cdot e^{(a)} \wedge \gamma \cdot f^{(b)} \wedge \gamma \cdot t^{(c)} = \det(\gamma) \cdot e^{(a)} \wedge f^{(b)} \wedge t^{(c)}.$$

Since there are three such expressions in the numerator and three in the denominator, the $\det(\gamma)$ cancel out, and we obtain $T_{abc}(\gamma E, \gamma F, \gamma T) = T_{abc}(E, F, T)$. \square

In fact, much more is true:

Theorem 14.23 (Fock-Goncharov [FG06]). *Two triples (E, F, T) and (E', F', T') are in the same G -orbit if and only if their triple ratios coincide. Moreover, every possible set of non-zero triple ratios is attained by some triple.*

The definition of positivity is then natural in terms of these coordinates:

Definition 14.24. The triple (E, F, T) is called *(BD)-positive* if all triple ratios are positive.

In order to facilitate the computations, we prove the following:

14. Positivity of Triples of Flags

Lemma 14.25. *Let $a, b, c \geq 0$ be such that $a + b + c = n$. Let F be the standard ascending, and E the standard descending flag, and let $T := uE$ for some $u \in U$. Then, under a suitable choice of generators,*

$$f^{(a)} \wedge e^{(b)} \wedge t^{(c)} = (-1)^{\lfloor b/2 \rfloor + \lfloor c/2 \rfloor + bc} \cdot \det u(a, b, c),$$

where $u(a, b, c)$ is as in Notation 14.16. Thus, for every $a, b, c \geq 1$ such that $a + b + c = n$, it holds:

$$T_{abc}(F, E, T) = \frac{\det u(a+1, b, c-1)}{\det u(a-1, b, c+1)} \cdot \frac{\det u(a, b-1, c+1)}{\det u(a, b+1, c-1)} \cdot \frac{\det u(a-1, b+1, c)}{\det u(a+1, b-1, c)}.$$

Proof. We choose the following generators:

$$f^{(i)} := e_1 \wedge \cdots \wedge e_i; \quad e^{(i)} := e_n \wedge \cdots \wedge e_{n-i+1}; \quad t^{(i)} = u_n \wedge \cdots \wedge u_{n-i+1};$$

where u_j denotes the j -th column of u . Then we use the formula for wedges in terms of determinants to obtain:

$$f^{(a)} \wedge e^{(b)} \wedge t^{(c)} = \det(e_1 \cdots e_a \mid e_n \cdots e_{n-b+1} \mid u_n \cdots u_{n-c+1}).$$

Flipping the last c columns changes this determinant by $(-1)^{\lfloor c/2 \rfloor}$, and gives

$$\det \begin{pmatrix} I_a & 0 & * \\ 0 & 0 & u(a, b, c) \\ 0 & J_b & * \end{pmatrix} = \det \begin{pmatrix} 0 & u(a, b, c) \\ J_b & * \end{pmatrix}$$

where J_b is the $(b \times b)$ matrix with 1 on the antidiagonal and 0 elsewhere. Next, we compute

$$\begin{aligned} \det \begin{pmatrix} 0 & u(a, b, c) \\ J_b & * \end{pmatrix} &= (-1)^{b+c+1} \det \begin{pmatrix} 0 & u(a, b, c) \\ J_{b-1} & * \end{pmatrix} = (-1)^{b+c+1} \cdot (-1)^{b+c} \cdot \det \begin{pmatrix} 0 & u(a, b, c) \\ J_{b-2} & * \end{pmatrix} \\ &= \cdots = \left(\prod_{j=c+2}^{b+c+1} (-1)^j \right) \det u(a, b, c) = (-1)^{\lfloor b/2 \rfloor} \cdot (-1)^{bc} \cdot \det u(a, b, c). \end{aligned}$$

The last equality follows from the fact that there are $\lfloor b/2 \rfloor$ odd numbers in $\{c+2, \dots, b+c+1\}$, unless b and c are both odd, in which case there are $\lfloor b/2 \rfloor = -\lceil b/2 \rceil$. We conclude that

$$f^{(a)} \wedge e^{(b)} \wedge t^{(c)} = (-1)^{\lfloor b/2 \rfloor} \cdot (-1)^{\lfloor c/2 \rfloor} \cdot (-1)^{bc} \cdot \det u(a, b, c).$$

Using this formula to compute $T_{abc}(F, E, T)$, the signs $(-1)^{\lfloor b/2 \rfloor}$ and $(-1)^{\lfloor c/2 \rfloor}$ each appear exactly once in the numerator and once in the denominator, so they all cancel out. As for the terms of the form $(-1)^{bc}$, these also cancel out, since the parity of bc does not change when each of b and c is changed by ± 2 . \square

Example 14.26. Let $n = 3$, and let consider (F, E, uE) . There is only one triple ratio to consider, namely T_{111} . By the previous formula,

$$u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow T_{111}(F, E, uE) = \frac{1}{xz-y} \cdot \frac{1}{1} \cdot \frac{y}{1} = \frac{y}{xz-y}.$$

Therefore (F, E, uE) is (BD)-positive if and only if y and $xz - y$ have the same sign, and are non-zero (the last condition ensures that (F, E, uE) is generic).

14.4. Equivalence of the two definitions

The two definitions may look different at first sight, but this is not the case:

Theorem 14.27 (Fock-Goncharov [FG06]). *A triple (E, T, F) is (GW)-positive if and only if it is (BD)-positive.*

We start by remarking the following, which accounts for the fact that the two definitions use different conventions in terms of orderings of the triples (as is apparent from Lemma 14.25):

Lemma 14.28. *The notion of (BD)-positivity is independent of the ordering of the triple.*

Proof. By [BD14, Lemma 5], it holds:

$$T_{abc}(E, F, T) = T_{bca}(F, T, E) = T_{bac}(F, E, T)^{-1};$$

this follows from elementary computations on the wedges appearing in the definition. The first equality shows that, upon applying a cyclic permutation, the set of triple ratios itself is permuted, and in particular the positivity of each triple ratio is not affected. The same holds under a transposition flipping the first two flags: the second equality shows that then the set of triple ratios is permuted and inverted, but again the positivity of each triple ratio is not affected. \square

Therefore it suffices to show that (E, T, F) is (GW)-positive if and only if (F, E, T) is (BD)-positive.

We start with the easier implication:

Proof of (GW) \Rightarrow (BD). Suppose that (E, T, F) is (GW)-positive. Note that both notions are not affected by the action of G : this holds by definition for (GW)-positivity, and follows from Lemma 14.22 for (BD)-positivity. Therefore we may assume that E is the standard descending flag, that F is the standard ascending flag, and that $T = uE$, where $u \in U_{>0}$. We now need to show that the triple (F, E, uE) is (BD)-positive.

First, it is generic, by Lemma 14.15. Moreover, each triple ratio T_{abc} is expressed in terms of minors of u , by Lemma 14.25. But u is totally positive, so each of these minors is positive. It follows that each triple ratio is positive and we conclude. \square

The other implication is more involved, and goes beyond the scope of this exposition, so we only present the proof in case $n = 3$. Recall that in this case there is only one triple ratio involved (Example 14.26), so it is surprising that from this one can recover a totally positive matrix, something which imposes four positivity conditions (Example 14.9). This apparent incompatibility of the two definitions is resolved by remembering that the definition of (GW)-positivity actually allows for more matrices than just totally positive ones: this was made explicit in Lemma 14.19, and it will be crucial in the proof.

Proof of (BD) \Rightarrow (GW) for $n = 3$. Suppose that (F, E, T) is (BD)-positive. Again, we note that both notions are not affected by the action of G , which is transitive on pair of transverse flags by Proposition 14.11, so we may assume that E is the standard descending flag and F is the standard ascending flag. Since this triple is generic, in particular T is transverse to F , and so by Proposition 14.11 there exists a unique $u \in U$ such that $T = uE$. Our goal is to show

14. Positivity of Triples of Flags

that there exists a diagonal matrix d such that $dud^{-1} \in U_{>0}$, which will conclude the proof by Lemma 14.19.

Now the condition that (F, E, uE) is (BD)-positive amounts to the positivity of the triple ratio $T_{111}(F, E, uE)$. By Example 14.26, we have:

$$T_{111}(F, E, uE) = \frac{y}{xz - y}; \quad \text{where } u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix}.$$

In other words, y and $(xz - y)$ are non-zero and have the same sign. We split into two cases.

Suppose that $y > 0$. Then $xz > y > 0$, so x and z are non-zero and have the same sign. If they are both positive, then u is totally positive by Example 14.9 and we are done. Otherwise, if x and z are both negative, we let $d := \text{diag}(1, -1, 1)$, and then

$$dud^{-1} = \begin{pmatrix} 1 & -x & y \\ 0 & 1 & -z \\ 0 & 0 & 1 \end{pmatrix}.$$

This matrix is totally positive by Example 14.9: indeed, all relevant entries are positive, and the only relevant other minor is $(-x)(-z) - y = xz - y > 0$.

Suppose instead that $y < 0$. Then $xz < y < 0$, so x and z are non-zero and have opposite signs. Let $d := (\text{sgn}(x), 1, \text{sgn}(z))$. Then

$$dud^{-1} = \begin{pmatrix} 1 & \text{sgn}(x)x & \text{sgn}(x)\text{sgn}(z)y \\ 0 & 1 & \text{sgn}(z)z \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & |x| & -y \\ 0 & 1 & |z| \\ 0 & 0 & 1 \end{pmatrix}.$$

This matrix is totally positive by Example 14.9: indeed, all relevant entries are positive, and the only relevant other minor is $|xz| - (-y) = -(xz - y) > 0$, since we know that $xz < 0$. \square

14.5. Generalizations

It is possible to generalize the notion of positivity of triples to quadruples and more generally to n -tuples. This may be done using *quadruple ratios* as in [BD14] or by checking positivity of some relevant triples, as in [GW18]. Another way to generalize positivity of triples it to consider generalized flag varieties, which we will do in this section. Yet another generalization is given by taking non-full flags.

So far we considered $G = \text{GL}(n, \mathbb{R})$ and the Borel subgroup of upper triangular matrices $B < G$. We defined a Borel subgroup to be a connected subgroup with $\text{Lie}(B) = \mathfrak{a} \oplus_{\alpha \in \Sigma^+} \mathfrak{g}_\alpha$. Note that

$$N_G(B) = \{g \in G : gBg^{-1} \subseteq B\} = B.$$

We call all subgroups of the form gBg^{-1} for $g \in G$ *Borel subgroups*. Then G acts by conjugation on the set of Borel subgroups $\{gBg^{-1}\}$. The group G also acts transitively on the flags by multiplication, and we get a G -equivariant map

$$\begin{aligned} \{\text{Borel subgroups } gBg^{-1}\} &\rightarrow \text{Flag}(\mathbb{R}^n) \\ gBg^{-1} &\mapsto gF \\ \text{Stab}_G(T) &\leftarrow T \end{aligned}$$

Thus we may identify

$$\text{Flag}(\mathbb{R}^n) \cong G/\text{Stab}_G(F) = G/N_G(B) = G/B.$$

The notion of flags in \mathbb{R}^n is tightly intertwined with $\text{GL}(n, \mathbb{R})$, and it is not obvious how to generalize flags. However Borel subgroups are available more generally, so we will use them to define our “generalized flags”. Let now G be a simple split real linear algebraic Lie group, for example $\text{SL}(n, \mathbb{R})$ or $\text{Sp}(4, \mathbb{R})$. General theory [Bor91] tells us, that there exists a Borel subgroup $B \subseteq G$, (defined as maximal among the connected solvable subgroups), which satisfies

$$N_G(B) = \{g \in G : gBg^{-1} \subseteq B\} = B.$$

We note that G acts on the set of Borel subgroups by $g.B = gBg^{-1}$. It is a theorem [Bor91, IV.11.1] that any two Borel subgroups are conjugate, therefore the action of G on the set of Borel subgroups is transitive.

Definition 14.29. The *generalized flag variety* is the set of Borel subgroups G/B .

For positivity of triples of elements in the generalized flag variety we need an opposite Borel group tB and a notion of positivity $U_{>0}$ for unipotent subgroups $U \subseteq B$. Previous talks have constructed these objects, see Chapter 12.

Definition 14.30. A triple $(F_1, F_2, F_3) \in (G/B)^3$ is *positive* if there is a $g \in G$ and $u \in U_{>0}$ such that $g(F_1, F_2, F_3) = ({}^tB, u{}^tB, B)$.

We note that for $G = \text{GL}(n, \mathbb{R})$, this is the definition of (GW)-positivity.

14.6. The flag variety for the symplectic group

We work out the case $\text{Sp}(4, \mathbb{R})$ and give an interpretation to the generalized flag variety. Recall that $\text{Sp}(4, \mathbb{R}) = \{A \in \text{GL}(4, \mathbb{R}) : AJ_{2,2}{}^tA = J_{2,2}\}$ for $J_{2,2} = \begin{pmatrix} & I_2 \\ -I_2 & \end{pmatrix}$. A standard Borel subgroup B and unipotent subgroup U are given by

$$\begin{aligned} B &= \left\{ \begin{pmatrix} X & Y \\ 0 & {}^tX^{-1} \end{pmatrix} \in \text{GL}(4, \mathbb{R}) : X = \begin{pmatrix} \star & \star \\ 0 & \star \end{pmatrix}, X{}^tY = {}^tYX \right\} \\ U &= \left\{ \begin{pmatrix} X & Y \\ 0 & {}^tX^{-1} \end{pmatrix} \in \text{GL}(4, \mathbb{R}) : X = \begin{pmatrix} 1 & \star \\ 0 & 1 \end{pmatrix} \right\} \\ U_{>0} &= \left\{ \begin{pmatrix} 1 & b+d & -bcd & bc \\ 0 & 1 & -(ad+ab+cd) & a+c \\ & & 1 & 0 \\ & & -(b+d) & 1 \end{pmatrix} \in \text{GL}(4, \mathbb{R}) \right\} \end{aligned}$$

We can interpret $\mathbb{R}^4 = \langle x_1, x_2, y_1, y_2 \rangle$ as a symplectic vector space with the symplectic 2-form ω defined by

$$\begin{aligned} \omega(x_i, y_j) &= \delta_{ij} = -\omega(y_j, x_i), \\ \omega(x_i, x_j) &= 0 = \omega(y_i, y_j). \end{aligned}$$

14. Positivity of Triples of Flags

Definition 14.31. A sub-vector space $V \subseteq \mathbb{R}^4$ is *isotropic* if $\omega(V, V) = 0$.

Example 14.32. The sub-vector space $\langle x_1 \rangle$ is isotropic. $\langle x_1, x_2 \rangle$ is maximal isotropic.

Definition 14.33. A sub-vector space that is maximal isotropic is called a *Lagrangian*.

It can be shown that the generalized flag variety $\mathrm{Sp}(4, \mathbb{R})/B$ can be identified with the isotropic complete flags in \mathbb{R}^4 with the standard symplectic form. In particular we have

$$\begin{aligned} \{\text{Borel subgroups of } \mathrm{Sp}(4, \mathbb{R})\} &\cong \{\text{isotropic complete flags in } (\mathbb{R}^4, \omega)\} \\ B &\mapsto \{\langle x_1 \rangle, \langle x_1, x_2 \rangle\} =: F \\ {}^tB &\mapsto \{\langle y_1 \rangle, \langle y_1, y_2 \rangle\} =: E. \end{aligned}$$

Note that $BE = E$ and ${}^tBF = F$. This justifies the following definition.

Definition 14.34. A *symplectic flag* is a complete isotropic flag.

15. The Maslov Index

by Samuel Bronstein, Ilia Smilga

All the definitions and theorems of this session come from Lion and Vergne's book: [LV80].

15.1. Maslov index in the symplectic group

Consider $\mathrm{Sp}(2n, \mathbb{R})$ the subgroup of $\mathrm{GL}(2n, \mathbb{R})$ of elements preserving the symplectic form ω defined by $w(X, Y) = {}^tX J_{n,n} Y$ for $X, Y \in \mathbb{R}^{2n}$ with:

$$J_{n,n} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

A basis of \mathbb{R}^{2n} in which w translates as $J_{n,n}$ is called a *symplectic basis*. In this regard, an endomorphism of \mathbb{R}^{2n} belongs to $\mathrm{Sp}(2n, \mathbb{R})$ if and only if it sends a symplectic basis onto a symplectic basis. We are interested here in the maximal isotropic subspaces, i.e. the Lagrangians:

Definition 15.1 (Lagrangian subspace). $L \subset \mathbb{R}^{2n}$ is *Lagrangian* if it is n -dimensional and $w|_{L \times L} = 0$. The set of Lagrangian subspaces will be denoted \mathcal{L} .

We refer to the previous sections for the fact that \mathcal{L} is actually the Shilov boundary of the Siegel disk, which is the symmetric space of $\mathrm{Sp}(2n, \mathbb{R})$. As such, there is a notion of positivity of triples on \mathcal{L} , of which we here bring another description. This notion of positivity being invariant by conjugation, it is natural to study the orbits of the $\mathrm{Sp}(2n, \mathbb{R})$ action on triples of Lagrangians.

Proposition 15.2. • *The set \mathcal{L} is homogeneous: $\mathcal{L} = \mathrm{Sp}(2n, \mathbb{R})/T$, where:*

$$T = \left\{ \begin{pmatrix} M & SM \\ 0 & {}^tM^{-1} \end{pmatrix} : S \in \mathrm{Sym}(n, \mathbb{R}), M \in \mathrm{GL}(n, \mathbb{R}) \right\} \quad (15.1)$$

- *The $\mathrm{Sp}(2n, \mathbb{R})$ action is transitive on the set of pairs of transverse Lagrangians, denoted $\mathcal{L}^{2,T}$. We have the identification:*

$$\mathcal{L}^{2,T} = \mathrm{Sp}(2n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \quad (15.2)$$

Proof. Of course the action by conjugation on the considered sets is an action by isometries, the main trouble is to convince oneself of the transitivity of those action: For a pair of transverse Lagrangian, concatenating two basis of the Lagrangians yields a symplectic basis of \mathbb{R}^{2n} . As we always can send a symplectic basis onto another one via an element of $\mathrm{Sp}(2n, \mathbb{R})$, it follows that the action is transitive. We let the reader compute the stabilizers. The stabilizer of the standard pair of transverse Lagrangian is the subset of T defined by $S = 0$. \square

15. The Maslov Index

More generally, one could have shown that the orbits of the conjugation action on pairs of Lagrangians are characterized by the degree of the intersection of the Lagrangians. Keep in mind that, given (L_1, L_2) a transverse pair of Lagrangians, w gives an isomorphism $L_2 \approx L_1^*$. That clarification done, let us give a first definition of the Maslov index, due to Kashiwara:

Definition 15.3 (Maslov Index). For (L_1, L_2, L_3) a triple of Lagrangians, the *Maslov index* is denoted $\tau(L_1, L_2, L_3)$ and defined as the signature of the quadratic form Q on $L_1 \times L_2 \times L_3$:

$$Q(x_1, x_2, x_3) = w(x_1, x_2) + w(x_2, x_3) + w(x_3, x_1) \quad (15.3)$$

There are several nice properties that we can easily deduce from this definition: The Maslov index is invariant by conjugation by an element of $\mathrm{Sp}(2n, \mathbb{R})$, it is an integer between $-3n$ and $3n$, it is alternating. Conveniently, we also have the following property, called the *chain rule*.

Proposition 15.4 (Chain rule). For (L_1, L_2, L_3, L_4) Lagrangians, we have:

$$\tau(L_1, L_2, L_3) = \tau(L_1, L_2, L_4) + \tau(L_2, L_3, L_4) + \tau(L_1, L_3, L_4) \quad (15.4)$$

This last property leads to think of the Maslov index as some kind of cocycle, but it is not the topic we are interested in here. For transverse triples, we have another possible definition, which is easier to handle:

Proposition 15.5. For a pairwise transverse triple of Lagrangians (L_1, L_2, L_3) , its Maslov index is the signature of the quadratic form S on L_2 defined by:

$$S(x_2) = \omega(p_{13}x_2, p_{31}x_2) \quad (15.5)$$

where p_{ij} denotes the projection on L_i with kernel L_j , well defined because of transversality.

Proof. A small computation allows us to show:

$$Q(x_1, x_2, x_3) = B(p_{13}x_2, p_{31}x_2) - B(x_1 - p_{13}x_2, x_3 - p_{31}x_2) \quad (15.6)$$

$$= S(y_2) - B(y_1, y_3) \quad (15.7)$$

with (y_1, y_2, y_3) another set of coordinates. As such, it is clear that the signature of Q equals the signature of S . \square

Remark 15.6. • This proposition is actually true as soon as $L_1 \cap L_3 = \emptyset$, we do not need that L_2 is transverse to the others. However, L_2 being transverse to L_1 and L_3 is equivalent to asking that S is non-degenerate.

- This gives a more precise bound on the Maslov index: For transverse triples, it is less or equal than n .
- We can check that $B(p_{13}x, p_{31}y)$ is symmetric on L_2 , so this is the bilinear form associated to S .

It remains to check that the Maslov index actually distinguishes the orbits of the $\mathrm{Sp}(2n, \mathbb{R})$ action on the set of transverse triples.

15.2. Maslov index in Hermitian Lie groups of tube type

Theorem 15.7. For (L_1, L_2, L_3) transverse triple, there is a symplectic basis $(p_1, \dots, p_n, q_1, \dots, q_n)$ and $0 \leq k \leq n$ such that:

$$L_1 = \langle p_1, \dots, p_n \rangle \tag{15.8}$$

$$L_2 = \langle q_1, \dots, q_n \rangle \tag{15.9}$$

$$L_3 = \langle p_1 + \varepsilon_1 q_1, \dots, p_n + \varepsilon_n q_n \rangle \tag{15.10}$$

where $\varepsilon_i = 1$ if $i \leq k$ and $\varepsilon_i = -1$ otherwise. Then the Maslov index is:

$$\tau(L_1, L_2, L_3) = n - 2k \tag{15.11}$$

Proof. the quadratic form S being non-degenerate, one can choose q_1, \dots, q_n basis of L_2 such that $S(q_i, q_j) = -\varepsilon_j \delta_{ij}$. But L_1 and L_2 being transverse, the symplectic form ω yields an isomorphism $L_1 = L_2^*$. So take the dual basis of (q_i) to be a basis of L_1 , denoted (p_i) .

Finally, fix $z_i = p_i + \varepsilon_i q_i$. Check that for all j , $B(z_i, q_j) = -\varepsilon_j \delta_{ij}$. Thus (by uniqueness of the dual basis) $z_i = -\varepsilon_i p_i$. This means that $(q_i + \varepsilon_i p_i)$ is a basis of L_3 , and so is $(p_i + \varepsilon_i q_i)$. \square

It is clear from this theorem that the maximality of the Maslov index corresponds to the maximality previously defined on $\text{Sp}(2n, \mathbb{R})$. This is the main motivation for generalizing the Maslov index to Hermitian Lie groups of tube type.

Remark 15.8 ($n=1$). For $n = 1$, the set of Lagrangians correspond to the circle seen as the boundary of the Poincaré Disk. In this setting, the Maslov index distinguishes the two orbits, on which it assigns 1 or -1 .

15.2. Maslov index in Hermitian Lie groups of tube type

To generalize this notion, one needs the notion of Euclidean Jordan algebra.

Definition 15.9. A Euclidean Jordan algebra is the data of:

- a real vector space V ,
- a scalar product on V denoted $\langle \cdot, \cdot \rangle$,
- a bilinear product $\circ : V \times V \rightarrow V$,

such that the following is verified:

- there is $e \in V$ such that for any $x \in V$, $x \circ e = e \circ x = x$,
- $x \circ y = y \circ x$,
- $x \circ (x^2 \circ y) = x^2 \circ (x \circ y)$,
- $\langle x \circ u, v \rangle = \langle u, x \circ v \rangle$.

If V is a Euclidean Jordan algebra, we also define Ω the *open cone of squares* as the interior of $\{v^2 : v \in V\}$.

A model-example for such a structure is $V = \text{Sym}(n, \mathbb{R})$ with the natural operations and the open cone of squares coincides with the set of positive definite matrices.

This is enough to define what is a Hermitian group of tube type:

15. The Maslov Index

Definition 15.10 (Hermitian group of tube type). A Hermitian Lie Group G is said to be of tube type if there is a Euclidean Jordan algebra V with open cone of squares Ω such that the homogeneous space G/K identifies with the tube domain $T_\Omega := \{X + iY : X \in V, Y \in \Omega\}$. One can rephrase that by saying that G acts transitively on T_Ω with stabilizers conjugates of K .

Example 15.11. For $G = \text{Sp}(2n, \mathbb{R})$, one can consider $V = \text{Sym}(n, \mathbb{R})$ with the action on T_Ω being:

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \cdot Z = (AZ + B)(CZ + D)^{-1} \tag{15.12}$$

Conveniently, we have a full classification of Hermitian Lie groups of tube type into four infinite families and one exceptional Lie group:

G	V
$\text{Sp}(2n, \mathbb{R})$	$\text{Sym}(n, \mathbb{R})$
$\text{SU}(n, n)$	$\text{Herm}(n, \mathbb{C})$
$\text{SO}^*(4n)$	$\text{Herm}(n, \mathbb{H})$
$E_7(-25)$	$\text{Herm}(3, \mathbb{O})$
$\text{SO}(2, n)$	$\mathbb{R} \times \mathbb{R}^{n-1}$ with $(a, v) \circ (b, w) = (ab + \langle v, w \rangle, av + bw)$

The infinite boundary on which we want to consider the G -action comes from Hermitian Lie group theory, we give a vague definition here:

Definition 15.12 (Shilov Boundary). For G Hermitian Lie group of tube type, the Shilov boundary S is a certain flag variety of G . For $G = \text{Sp}(2n, \mathbb{R})$, S corresponds to the set of Lagrangian subspaces.

Actually, the action on S is quite well understood:

Proposition 15.13. *There is a point $\infty \in S$ such that:*

- $\{s \in S | s \pitchfork \infty\} \approx V \subset \overline{T_\Omega}$
- $\text{Stab}_G(\infty) = \{g : T_\Omega \rightarrow T_\Omega : g(Z) = AZ + B \text{ with } A \in \text{GL}(V) \text{ and } B \in V\}$. In particular the G -action on the set of transverse pairs is transitive.
- $\{s \in S | s \pitchfork \infty, s \pitchfork 0\} \approx V^* = \{v \in V : \det v \neq 0\}$
- The stabilizer of a pair $(0, \infty)$ is $G(\Omega) = \{g \in \text{GL}(V) : g(\Omega) = \Omega\}$.

As a corollary, the G -orbits of transverse triples correspond to $G(\Omega)$ -orbits in V^* . The main tool that we use is the spectral decomposition for Euclidean Jordan algebras:

Proposition 15.14 (Spectral decomposition). *Let V be a Euclidean Jordan algebra. Then for any $v \in V$, there is c_1, \dots, c_r a Jordan frame, i.e. such that:*

- $\forall i$ we have $c_i^2 = c_i$ and $c_i \neq c_j + c_k$,
- $\forall i \neq j$ we have $c_i c_j = 0$,
- $\sum c_i = e$,

and there are unique eigenvalues $\lambda_1, \dots, \lambda_r$ such that $v = \sum \lambda_i c_i$.

Remark 15.15. As $\det v = \prod \lambda_i$, v is invertible if and only if $\lambda_i \neq 0$ for all $i \neq 0$. Then we can define the signature of v in the following way: $\text{sgn}(v) = |\{i : \lambda_i > 0\}| - |\{i : \lambda_i < 0\}|$.

It is a well known fact that the signature is invariant under the $G(\Omega)$ -action on V^* , and then the Maslov index is the corresponding invariant for the action on transverse triples.

Part IV.

Theta-Positivity

Clarence Kineider

ENS Rennes

Pierre-Louis Blayac

Institut des Hautes Études Scientifiques

Balthazar Fléchettes

Institut des Hautes Études Scientifiques

Dani Kaufman

Ruprecht-Karls-Universität Heidelberg

Fernando Camacho Cadena

Ruprecht-Karls-Universität Heidelberg

Merik Niemeyer

Ruprecht-Karls-Universität Heidelberg

16. Introduction to Theta-Positivity

by Clarence Kineider

Total positivity in split real Lie groups and maximality in Hermitian Lie groups of tube type have a lot in common. The notion of Θ -positivity will unify those two cases under the same definition. Moreover, we have a complete description of all semisimple Lie groups admitting a Θ -positive structure which includes two new family of groups, namely $\mathrm{SO}(p, q)$, $p \neq q$ and exceptional groups with restricted root system of type F_4 .

In this chapter we will do every computation in $\mathrm{SL}(n, \mathbb{R})$.

Let G a real semisimple Lie group with finite center and \mathfrak{g} its Lie algebra. Let K be a maximal compact subgroup of G , \mathfrak{k} its Lie algebra, and \mathfrak{k}^\perp the orthogonal of \mathfrak{k} with respect to the Killing form. Let \mathfrak{a} be a maximal abelian Cartan subspace of \mathfrak{k}^\perp , and denote by $\Sigma = \Sigma(\mathfrak{g}, \mathfrak{a})$ the system of restricted roots (see Part II for all necessary definitions). Choose $\Delta \subset \Sigma$ a set of simple roots and let Σ^+ be the set of positive roots with respect to Δ .

Example 16.1. For $G = \mathrm{SL}(n, \mathbb{R})$, $K = \mathrm{SO}(n)$ and \mathfrak{k} is the set of skew-symmetric matrices. Then \mathfrak{k}^\perp is the set of traceless symmetric matrices, and \mathfrak{a} is the set of traceless diagonal matrices. The restricted root system is $\Sigma = \{\alpha_{ij} := \varepsilon_i - \varepsilon_j : i, j \in \{1, \dots, n\}, i \neq j\}$, where $\varepsilon_i : (a_{pq})_{p, q \in \{1, \dots, n\}} \mapsto a_{ii}$. The standard choice of simple roots is $\Delta = (\alpha_{i, i+1})_{i \in \{1, \dots, n-1\}}$, and the set of positive roots associated is $\Sigma^+ = (\alpha_{ij})_{i < j}$. The weight space $\mathfrak{g}_{\alpha_{ij}}$ is the 1-dimensional subspace of $\mathfrak{sl}(n, \mathbb{R})$ spanned by E_{ij} .

Now let $\Theta \subset \Delta$ be any subset of simple roots (Θ can be empty). Let $\Sigma_\Theta^+ = \Sigma^+ \setminus \mathrm{Span}(\Delta \setminus \Theta)$, and let

$$\mathfrak{u}_\Theta = \sum_{\alpha \in \Sigma_\Theta^+} \mathfrak{g}_\alpha, \quad \mathfrak{u}_\Theta^{\mathrm{opp}} = \sum_{\alpha \in \Sigma_\Theta^+} \mathfrak{g}_{-\alpha}$$

and

$$\mathfrak{l}_\Theta = \mathfrak{g}_0 \oplus \sum_{\alpha \in \Sigma^+ \cap \mathrm{Span}(\Delta \setminus \Theta)} \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}.$$

As Lie algebra, \mathfrak{g} splits into $\mathfrak{g} = \mathfrak{u}_\Theta^{\mathrm{opp}} \oplus \mathfrak{l}_\Theta \oplus \mathfrak{u}_\Theta$. Moreover \mathfrak{l}_Θ acts on \mathfrak{u}_Θ by the Lie bracket, and we would like to understand this action and its properties. The group G acts on \mathfrak{g} by the adjoint representation, denote by P_Θ the normalizer in G of \mathfrak{u}_Θ and P_Θ^{opp} the normalizer of $\mathfrak{u}_\Theta^{\mathrm{opp}}$. Those are the standard parabolic subgroups associated with Θ , and the corresponding Levi subgroup is $L_\Theta = P_\Theta \cap P_\Theta^{\mathrm{opp}}$. Let us complete the list with $U_\Theta = \exp(\mathfrak{u}_\Theta)$ and $\mathfrak{p}_\Theta = \mathrm{Lie}(P_\Theta)$.

Proposition 16.2. \mathfrak{l}_Θ is the Lie algebra of L_Θ and $\mathfrak{p}_\Theta = \mathfrak{l}_\Theta \oplus \mathfrak{u}_\Theta$.

Example 16.3. For $G = \mathrm{SL}(n, \mathbb{R})$ let us choose $\Theta = \{\alpha_{i, i+1}, \alpha_{j, j+1}\}$ with $1 \leq i < j < n$. Then the Lie algebra $\mathfrak{sl}(n, \mathbb{R})$ admits the following block decomposition :

16. Introduction to Theta-Positivity

$$\begin{matrix}
 & & & i & i+1 & & j & j+1 & & \\
 & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\
 i & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \hline
 i+1 & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\
 & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \hline
 j & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\
 j+1 & & & * & \dots & * & * & \dots & * & * & \dots & * \\
 & & & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\
 & & & * & \dots & * & * & \dots & * & * & \dots & *
 \end{matrix}
 , \quad \text{tr} = 0.$$

In the following examples we refer to this block decomposition when speaking about matrices defined block-wise. The Levi subalgebra \mathfrak{l}_Θ is the set of traceless block-diagonal matrices, the algebra \mathfrak{u}_Θ is the set of block strictly upper triangular matrices and $\mathfrak{p}_\Theta = \mathfrak{l}_\Theta \oplus \mathfrak{u}_\Theta$ is the set of traceless block-upper triangular matrices :

$$\mathfrak{l}_\Theta = \left\{ \left(\begin{array}{c|c|c} * & & \\ \hline & * & \\ \hline & & * \end{array} \right), \text{tr} = 0 \right\}, \quad \mathfrak{u}_\Theta = \left\{ \left(\begin{array}{c|c|c} & * & * \\ \hline & & * \\ \hline & & \end{array} \right), \text{tr} = 0 \right\}, \quad \mathfrak{p}_\Theta = \left\{ \left(\begin{array}{c|c|c} * & * & * \\ \hline & * & * \\ \hline & & * \end{array} \right), \text{tr} = 0 \right\}.$$

The parabolic subgroup P_Θ is the stabilizer of the standard (partial) flag of \mathbb{R}^n with subspaces of dimensions $(0, i, j, n)$.

Since L_Θ is a reductive group, we need to know its center, the Cartan subalgebra of its Lie algebra etc. in order to understand the action of L_Θ on \mathfrak{u}_Θ . First, \mathfrak{l}_Θ splits into its center $Z(\mathfrak{l}_\Theta)$ and its semisimple part $\mathfrak{l}'_\Theta = [\mathfrak{l}_\Theta, \mathfrak{l}_\Theta]$:

$$\mathfrak{l}_\Theta = Z(\mathfrak{l}_\Theta) \oplus \mathfrak{l}'_\Theta.$$

Its center $Z(\mathfrak{l}_\Theta)$ once again splits in two parts: $\mathfrak{z}_\Theta = Z(\mathfrak{l}_\Theta) \cap \mathfrak{k}^\perp = Z(\mathfrak{l}_\Theta) \cap \mathfrak{a}$ and $Z(\mathfrak{l}_\Theta) \cap \mathfrak{k}$.

Fact 16.4. The Cartan subalgebra of \mathfrak{l}_Θ is $\mathfrak{a}_\Theta = \mathfrak{a} \cap \mathfrak{l}'_\Theta$, its restricted root system is the restriction of $\Sigma \cap \text{Span}(\Delta \setminus \Theta)$ to \mathfrak{a}_Θ , a possible choice of simple roots are the restriction of $\Delta \setminus \Theta$ to \mathfrak{a}_Θ , and the corresponding Dynkin diagram is the largest subgraph of the Dynkin diagram of G with vertices $\Delta \setminus \Theta$.

Note that since $\mathfrak{a} \subset \mathfrak{l}_\Theta \cap \mathfrak{k}^\perp$, we have

$$\mathfrak{a} = \mathfrak{a}_\Theta \oplus \mathfrak{z}_\Theta.$$

In particular, \mathfrak{z}_Θ acts on \mathfrak{u}_Θ by restriction of the action of \mathfrak{a} on \mathfrak{g} , so \mathfrak{u}_Θ decomposes into weight spaces for this action:

$$\mathfrak{u}_\Theta = \bigoplus_{\beta \in \mathfrak{z}_\Theta^*} \mathfrak{u}_\beta, \tag{16.1}$$

where

$$\mathfrak{u}_\beta = \{x \in \mathfrak{u}_\Theta : \forall z \in \mathfrak{z}_\Theta, \text{ad}(z).x = \beta(z)x\} \tag{16.2}$$

is L_Θ^0 -invariant.

Moreover if a root $\alpha \in \mathfrak{a}^*$ vanishes on \mathfrak{z}_Θ (i.e. lies in \mathfrak{a}_Θ^*) then $\alpha \in \text{Span}(\Delta \setminus \Theta)$.

The inclusion $\mathfrak{z}_\Theta \subset \mathfrak{a}$ induces a projection $i^* : \mathfrak{a}^* \rightarrow \mathfrak{z}_\Theta^*$ given by the restriction of a linear form on \mathfrak{a} to \mathfrak{z}_Θ . Thus we can write \mathfrak{u}_β as a sum of weight spaces for the action of \mathfrak{a} :

$$\mathfrak{u}_\beta = \sum_{\alpha \in \Sigma_\Theta^+, \alpha|_{\mathfrak{z}_\Theta} = \beta} \mathfrak{g}_\alpha.$$

Suppose that β is the restriction of a root $\alpha \in \Theta$. Then α is the only element of $\text{Span}(\Theta)$ which restricts to β , allowing us the following slight abuse of notation:

$$\mathfrak{u}_\alpha := \mathfrak{u}_\beta = \sum_{\gamma \in \text{Span}_\mathbb{N}(\Delta \setminus \Theta)} \mathfrak{g}_{\alpha+\gamma}. \quad (16.3)$$

Example 16.5. Using the same choice of Θ as in the previous example, the objects introduced above are the following: the center $Z(\mathfrak{l}_\Theta)$ of \mathfrak{l}_Θ is the set of matrices of the form

$$\left(\begin{array}{c|c|c} \lambda I_i & & \\ \hline & \mu I_{j-i} & \\ \hline & & \nu I_{n-j} \end{array} \right)$$

with $i\lambda + (j-i)\mu + (n-j)\nu = 0$. The semisimple part of \mathfrak{l}_Θ is then the set of block-diagonal matrices for which every block is traceless. In this situation, $Z(\mathfrak{l}_\Theta)$ is entirely contained in \mathfrak{a} , so $\mathfrak{z}_\Theta = Z(\mathfrak{l}_\Theta)$. The weight spaces \mathfrak{u}_β are the block-upper triangular matrices in \mathfrak{u}_Θ for which only one block is non-zero, and the \mathfrak{u}_β with $\beta \in \Theta$ are the one where the non-zero block is just above the diagonal. The third \mathfrak{u}_β (the one with the top right block) is associated to the restriction of the root $\alpha_{i,j+1} = \alpha_{i,i+1} + \dots + \alpha_{j,j+1}$ to \mathfrak{z}_Θ , and all the roots $\alpha_{i+1,i+2}, \dots, \alpha_{j-1,j}$ vanish on \mathfrak{z}_Θ .

In [GW18] the irreducibility of \mathfrak{u}_β is stated as a fact, but the proof is not completely straightforward and is due to Kostant.

Theorem 16.6 ([Kos10, Theorem 0.1]). *For all $\beta \in \mathfrak{z}_\Theta^*$, \mathfrak{u}_β is an irreducible representation of L_Θ^0 , and*

$$[\mathfrak{u}_\beta, \mathfrak{u}_{\beta'}] = \mathfrak{u}_{\beta+\beta'}.$$

Definition 16.7. Let V be a finite dimensional real vector space. A *cone* in V is a subset C stable by positive scalar multiplication, i.e. $\forall x \in C, \forall \lambda > 0$ we have $\lambda x \in C$. The cone C is said *sharp* if it contains no affine line.

Example 16.8. The sharpness condition ensure the cone is not too "wide", for instance the half plane $\mathbb{H} \subset \mathbb{R}^2$ is a cone, but not sharp since it contains many horizontal affine lines.

We are interested in sharp *convex* cones as they will give a good analog of $\mathbb{R}^{>0}$ in \mathbb{R} , which is our first example of sharp convex cone. A second example of sharp convex cone is the set of positive definite symmetric matrices in the space of symmetric matrices.

Since L_Θ is not necessarily connected, denote L_Θ^0 the connected component of the identity in L_Θ . We now have all the tool to give the definition of Θ -positivity.

Definition 16.9. We say that (G, Θ) admits a *Θ -positive structure* if for all $\beta \in \Theta$, the action of L_Θ^0 on \mathfrak{u}_β preserves a sharp convex cone.

16. Introduction to Theta-Positivity

We already know two notions of positivity, namely total positivity and maximality. Our goal is now to explain why those are Θ -positive structure, for a suitable choice of Θ .

Example 16.10. Let us begin with total positivity in $G = \mathrm{SL}(n, \mathbb{R})$ to simplify the notations. For $\Theta = \Delta$ (with the usual choice of Δ for $\mathrm{SL}(n, \mathbb{R})$), the Levi subgroup L_Θ is the subgroup of diagonal matrices of determinant 1, and all of the weight spaces associated to elements of Θ are one-dimensional:

$$\forall \beta = \alpha_{i,i+1} \in \Theta, \mathfrak{u}_\beta = \mathfrak{g}_\beta = \langle E_{i,i+1} \rangle.$$

In Lusztig's total positivity, see Chapter 12, the semigroup of positive elements of a split real Lie group is generated by elements of the form $u_i(t) = I_n + tE_{i,i+1} = \exp(tE_{i,i+1})$, with $t > 0$. Here the sharp convex cone in $\mathfrak{u}_\beta = \mathfrak{g}_{\alpha_{i,i+1}}$ is the cone of positive scalar multiples of $E_{i,i+1}$.

Example 16.11. Now for maximality, let us treat the case $G = \mathrm{Sp}(2n, \mathbb{R})$. For the choice $\Theta = \{\alpha_n\}$ where α_n is the root at the end of the Dynkin diagram with two arrows pointing out, there is only one space \mathfrak{u}_β , namely \mathfrak{u}_{α_n} . In this case the space \mathfrak{u}_{α_n} can be identified with the space of symmetric matrices of size n , and the sharp convex cone is the set of positive definite matrices. More details on the computations can be found in the exercise section.

17. The Classification Theorem

by Pierre-Louis Blayac, Balthazar Fléchettes

We will try to prove the following classification theorem for Θ -positive structures.

Theorem 17.1 ([GW18, thm 4.3]). *A simple Lie group G admits a Θ -positive structure if and only if (G, Θ) falls in one of the following cases:*

- (1) G is a split real Lie form, and $\Theta = \Delta$,
- (2) G is Hermitian of tube type and $\Theta = \{\alpha_r\}$,
- (3) G is locally isomorphic to $\mathrm{SO}(p, q)$ with $p < q$, and $\Theta = \{\alpha_1, \dots, \alpha_{p-1}\}$,
- (4) G is a real form of F_4 , E_6 , E_7 or E_8 whose restricted root system is of type F_4 , and $\Theta = \{\alpha_1, \alpha_2\}$.

Recall that a Θ -positive structure on G is linked to the existence of sharp convex cones invariant under the Levi subgroup L_Θ° in \mathfrak{u}_β for $\beta \in \Theta$. Conveniently, [Ben00, Prop. 4.7] gives a nice criterion for an irreducible representation of a semisimple Lie group to preserve a sharp convex cone, which translates into the following fact:

Fact 17.2. A semi-simple Lie group G admits a Θ -positive structure is positive if and only if

- (i) \mathfrak{g}_β is 1-dimensional for every $\beta \in \Theta$,
- (ii) The node in the Dynkin diagram of G corresponding to the restricted root $\beta \in \Theta$ is either disconnected from $\Delta \setminus \Theta$, or linked to $\Delta \setminus \Theta$ via a double arrow pointing toward $\Delta \setminus \Theta$.

Once this is established, it is not hard to find the list of Theorem 17.1. For example, one can use the Tables 1 and 4 in [OV94] to write an explicit list of the Lie groups admitting a Θ -positive structure, and then check in another reference the list of split real Lie forms (all rows of Table 4 of [OV94] whose column “ $\dim \mathfrak{g}_{\lambda_j}$ ” only indicates 1; one can also consult [Kna02, §C.3-4]) and Hermitian groups of tube type [BIW07, Rem. 2.1]. Another reference is [Hel01, Chapter X, Table VI]. Beware that there are mistakes in the tables of [OV94], but we believe that the information we need here is correctly displayed there. The procedure is the following. For each row of Table 4 of [OV94]:

- look at the column “ Σ ” giving the restricted root system;
- go to the corresponding row of Table 1;
- since any Dynkin diagram has at most one double arrow, there are at most two subsets of vertices satisfying Condition (ii) of Fact 17.2 (the full set of vertices and, if there is a double arrow, another subset);
- for each of these two subsets i_1, \dots, i_k , go back to Table 4 column “ $\dim \mathfrak{g}_{\lambda_j}$ ” to check Condition (i) of Fact 17.2, *i.e.* that $\mathfrak{g}_{\lambda_{i_1}}, \dots, \mathfrak{g}_{\lambda_{i_k}}$ all have dimension 1.

For instance, asking $\dim(\mathfrak{g}_{\lambda_j}) = 1$ for any j is equivalent to asking the group to be real split.

17. The Classification Theorem

Remark 17.3. If G_1 and G_2 are two semi-simple Lie groups with respective simple roots Δ_1 and Δ_2 , which are respectively Θ_1 and Θ_2 -positive (and we may allow Θ_1 or Θ_2 to be empty, considering that any group is \emptyset -positive), then $G_1 \times G_2$ is $\Theta_1 \sqcup \Theta_2$ -positive. This is clear for instance in view of Fact 17.2.

In order to prove Fact 17.2, we will need a theorem by Yves Benoist concerning representations of subsemigroups of $\mathrm{GL}(V)$, where V is a fixed real vector space of dimension d .

17.1. A theorem by Yves Benoist

Before quoting Benoist's theorem, we need to introduce some terminology:

Definition 17.4. We say a semi-group $G \subset \mathrm{GL}(V)$ is *irreducible* if any invariant non-trivial subspace is full, i.e.

$$\forall W \subset V, \quad G \cdot W = W \implies W = \{0\} \text{ or } W = V.$$

Definition 17.5. • $\gamma \in \mathrm{GL}(V)$ is *proximal* if it has only one eigenvalue of largest modulus, and if this eigenvalue is simple (and hence real). If moreover it is positive, we say that γ is *positively proximal*.

- If $\gamma \in \mathrm{GL}(V)$ is proximal, we will denote by $x_\gamma^+ \in \mathbb{P}(V)$ the projectivization of the eigenline corresponding to the eigenvalue of largest modulus. x_γ^+ is called the *attracting fixed point* of γ in $\mathbb{P}(V)$. We denote by y_γ^- the γ -invariant complementary hyperplane. Note that $\gamma^n x \rightarrow x_\gamma^+$ when $n \rightarrow \infty$ for any $x \notin y_\gamma^-$.
- A (semi-)group $\Gamma < \mathrm{GL}(V)$ is *proximal* if it has a proximal element. If moreover every proximal element of Γ is positively proximal, we say that Γ is *positively proximal*.

These definitions are of interest for the notion of limit set:

Proposition 17.6. If $\Gamma < \mathrm{GL}(V)$ is an irreducible and proximal subsemigroup, then

$$\Lambda_\Gamma := \overline{\{x_\gamma^+ : \gamma \in \Gamma \text{ proximal}\}}$$

is the smallest non-empty Γ -invariant closed subset of $\mathbb{P}(V)$. It is called the *limit set* of Γ .

Proof. The set Λ_Γ is clearly non-empty.

Let us check that it is Γ -invariant. It is clear that $\Lambda_\Gamma \subset \Gamma \Lambda_\Gamma$. Consider $\gamma, \eta \in \Gamma$ with η proximal. If Γ was a group we could just observe that $\gamma \eta \gamma^{-1} \in \Gamma$ is also proximal and $\gamma x_\eta^+ = x_{\gamma \eta \gamma^{-1}}^+ \in \Lambda_\Gamma$. When Γ is not a group the argument is more delicate. By irreducibility we can find $g \in \Gamma$ such that $g(\gamma(x_\eta^+)) \notin y_\eta^-$. Then one may check that for n large enough $\gamma \eta^n g \in \Gamma$ is proximal and that $x_{\gamma \eta^n g}^+ \rightarrow \gamma x_\eta^+$.

Consider now a non-empty closed set $F \subset \mathbb{P}(V)$ with $\Gamma F \subset F$, and let us check that $\Lambda_\Gamma \subset F$. For any $x \in F$ and $\gamma \in \Gamma$ proximal, by irreducibility there exists $g \in \Gamma$ such that $g(x) \notin y_\gamma^-$. Then $x_\gamma^+ = \lim \gamma^n g x \in F$. \square

Definition 17.7. A *convex cone* C of V is a non-trivial convex subset of V invariant under multiplication by positive scalars. It is *sharp* or *properly convex* if its closure does not contain a full line. This is consistent with Definition 16.7.

Theorem 17.8 ([Ben00, prop. 3.11]). *An irreducible semi-group $\Gamma \subset \mathrm{GL}(V)$ preserves a sharp convex cone of V if and only if it is positively proximal.*

Proof (only an idea for the converse). Suppose Γ preserves a sharp convex cone $C \subset V$.

First observe that for any proximal element $\gamma \in \Gamma$, there exists a point $p \in C$ which does not meet the repelling hyperplane of γ (C has non-empty interior by irreducibility of Γ), so that $\frac{\gamma^n p}{\|\gamma^n p\|}$ converges to a point x in $x_\gamma^+ \cap \overline{C}$. Since C is sharp, this is a γ -invariant half-line contained in x_γ^+ , so γ is positively proximal. Therefore we only need to prove that Γ is proximal.

Without loss of generality, we may assume that Γ contains all the homotheties $\{\lambda \mathrm{id}\}_{\lambda > 0}$. We will denote by $\overline{\Gamma}$ the closure of Γ in $\mathrm{End}(V)$. Consider $M := \inf_{\gamma \in \overline{\Gamma} - \{0\}} \mathrm{rank} \gamma > 0$.

Suppose $M = d$, i.e. that $\overline{\Gamma} \setminus \{0\} \subset \mathrm{GL}(V)$. The projection in $\mathbb{P}(\mathrm{End}(V))$ of $\overline{\Gamma} \setminus \{0\}$ is closed and hence compact, and equals that of $\overline{\Gamma} \cap \mathrm{SL}(V)$, which is then itself compact since $\pi: \mathrm{End}(V) \setminus \{0\} \rightarrow \mathbb{P}(\mathrm{End}(V))$ is injective on $\mathrm{SL}(V)$. Thus $\overline{\Gamma} \cap \mathrm{SL}(V)$ is a compact group (any relatively compact sequence of the form $(g^n)_{n \geq 0} \subset \mathrm{GL}(V)$ has g^{-1} as an accumulation point); let μ be its Haar measure. Then $\int g(x) d\mu(g)$ is non-zero and $\overline{\Gamma} \cap \mathrm{SL}(V)$ -invariant for any $x \in C \setminus \{0\}$, and hence spans a Γ -invariant half-line. Since Γ is irreducible, we must have $d = 1$, and Γ is proximal.

Suppose $M < d$ and consider $\gamma \in \overline{\Gamma} \setminus \{0\}$ with rank M . By irreducibility of Γ there exists $g \in G$ such that $g \mathrm{im}(\gamma) \not\subset \ker(\gamma)$. Let $\eta := g\gamma$, so that $\ker(\eta) \cap \mathrm{im}(\eta) = \emptyset$ (otherwise $\mathrm{rk}(\eta^2) < \mathrm{rk}(\eta) = \mathrm{rk}(\gamma) = M$), and hence $\mathrm{im}(\eta) \oplus \ker(\eta) = V$.

Consider now the semigroup $\eta \overline{\Gamma} \eta \subset \overline{\Gamma} \subset \mathrm{End}(V)$. It stabilizes $\mathrm{im}(\gamma)$, and acts trivially on $\ker(\gamma)$, so it naturally identifies with a semi group $\Gamma' \subset \mathrm{GL}(\mathrm{im} \gamma)$ (γ restricts to a bijection on $\mathrm{im}(\gamma)$), which preserves the sharp convex cone $C' = \overline{C} \cap \mathrm{im}(\gamma) \setminus \{0\}$. That $\eta \overline{\Gamma} \eta \subset \overline{\Gamma}$ implies that $\dim(\mathrm{im}(\eta)) \geq M' := \inf_{g \in \Gamma \setminus \{0\}} \mathrm{rk}(g) \geq M = \mathrm{rk}(\eta)$. Thus, the first case above yields $\mathrm{rank} \eta = M' = 1$, so that η and hence $\overline{\Gamma}$ is proximal. Since being proximal is an open condition, we conclude that Γ is proximal.

To prove the converse, one uses the irreducibility and positive proximality of Γ to lift in a Γ -invariant way the limit set $\Lambda_\Gamma \subset \mathbb{P}(V)$ to a set of half-lines in V whose convex hull is a sharp convex cone. However this requires a lot of work (several intermediate results) so we refer to Benoist's article for a complete proof. \square

17.2. Proof of the classification theorem

We now proceed to show fact 17.2 using theorem 17.8. For this purpose, we fix a connected reductive Lie group G (it corresponds to the Levi subgroup L_Θ° in theorem 17.1), and an irreducible representation $\rho: G \rightarrow \mathrm{GL}(V)$.

Recall the following facts:

- the weights $\lambda \in \mathfrak{a}^*$ of ρ lie in the *weight lattice*

$$P := \left\{ \lambda \in \mathfrak{a}^* : \frac{2\langle \lambda, \alpha \rangle}{\langle \alpha, \alpha \rangle} \in \mathbb{Z}, \forall \alpha \in \Delta \right\},$$

- two weights differ by an element of $\mathrm{Span}_{\mathbb{Z}} \Delta$,

17. The Classification Theorem

- the set of weights of ρ is (partially) ordered as follow:

$$\begin{aligned}\mu \geq \lambda &\iff \mu - \lambda \in \text{Span}_{\mathbb{N}} \Delta \\ &\implies \mu(X_0) \geq \lambda(X_0)\end{aligned}$$

where $X_0 \in \mathfrak{a}$ is a (fixed) vector with $\alpha(X_0) > 0$ for any $\alpha \in \Delta$.

It is a classical result of the theory of representations of reductive groups that, by irreducibility, there exists a unique highest weight $\omega \in P$.

Proposition 17.9. $\rho(G)$ is proximal if and only if the highest weight space V_ω of ρ is of dimension 1. Moreover, in this case, $\rho(G)$ is positively proximal if and only if $\omega \in 2P$, i.e.

$$\forall \alpha \in \Delta, \quad \frac{\langle \omega, \alpha \rangle}{\langle \alpha, \alpha \rangle} \in \mathbb{Z}.$$

Proof. Suppose V_ω is of dimension 1. Since ω is the highest weight, we have

$$\omega(X_0) > \lambda(X_0) \tag{17.1}$$

for all $\lambda \neq \omega$ weight of ρ . But the eigenvalues of $\rho(e^{X_0})$ are by definition the numbers of the form $e^{\lambda(X_0)}$ where λ is a weight of ρ , with corresponding eigenspace the weight space V_λ . Thus, (17.1) and the hypothesis $\dim V_\omega = 1$ means that $\rho(e^{X_0})$ is proximal.

Suppose that $\rho(G)$ is proximal, and pick $g \in G$ such that $\rho(g)$ is proximal. Consider the Jordan decomposition $g = kan$: the elements k, a, n pairwise commute, k is conjugate to $k' \in K$, while a is conjugate to $\exp(X) \in \exp(\overline{\mathfrak{a}^+})$, and n is conjugate to $\exp(N) \in \exp \mathfrak{u}$ (where $\mathfrak{u} = \sum_{\alpha \in \sigma_+} \mathfrak{g}_\alpha$).

The element $\rho(k') \in \rho(K) \subset O(V, q)$ (for some scalar product q) only has eigenvalues of modulus 1, and so does $\rho(k)$. Consider a basis $e_1, \dots, e_d \in V$ with $e_i \in V_{\lambda_i}$ for some weight λ_i , and with an ordering such that $\lambda_i > \lambda_j \implies i < j$. Observe that $d\rho(\mathfrak{g}_\alpha) \cdot V_\lambda \subset V_{\lambda+\alpha}$ for all $\alpha \in \sigma_+$ and $\lambda \in P$, so that $\rho(\exp(N)) = \exp(d\rho(N))$ is upper triangular with ones on the diagonal for the basis e_1, \dots, e_d . The moduli of the eigenvalues of $\rho(g)$ are hence given by those of $\rho(a)$, which is therefore proximal.

The eigenvalues of $\rho(a)$ are the numbers of the form $e^{\lambda(X)}$ where λ is a weight of ρ . Since ω is the highest weight, $e^{\omega(X)}$ is the highest eigenvalue of $\rho(a)$. Since $\rho(a)$ is proximal, we then have $\dim V_\omega = 1$.

We only prove that if $G = \text{SL}(n, \mathbb{R})$ and $\rho(G)$ is positively proximal then $\omega \in 2P$ (for different G , the idea is similar using $\mathfrak{sl}(2, \mathbb{R})$ -triplets).

Recall that the simple roots of G are $\Delta = \{\alpha_1, \dots, \alpha_{n-1}\}$ where $\alpha_i(\text{diag}(\lambda_1, \dots, \lambda_n)) = \lambda_{i+1} - \lambda_i$.

We want to show that for all $i \in \{1, \dots, n-1\}$,

$$\frac{\langle \omega, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle} \in \mathbb{Z}.$$

Write (where the two -1 's appear at positions i and $i+1$)

$$k_i = \text{diag}(1, \dots, 1, -1, -1, 1, \dots, 1) = e^{i\pi(E_{ii} - E_{i+1, i+1})} = e^{i\pi H_{\alpha_i}}.$$

Since $\rho(k_i)$ and $\rho(e^{X_0})$ are both diagonal matrices, $\rho(k_i)$ has only ± 1 as diagonal values, and $\rho(e^{X_0})$ is proximal, $\rho(k_i e^{X_0})$ is also proximal, so it is positively proximal. In particular, the eigenvalue of $\rho(k_i)$ on V_ω is 1.

Pick $v \in V_\omega \setminus \{0\}$. One has (after extending $d\rho : \mathfrak{g} \rightarrow \mathfrak{sl}_d \mathbb{R}$ to the complexification $d\rho^{\mathbb{C}} : \mathfrak{g}^{\mathbb{C}} \rightarrow \mathfrak{sl}(d, \mathbb{C})$),

$$v = \rho(k_i)v = \rho(e^{i\pi H_{\alpha_i}})v = e^{i\pi\omega(H_{\alpha_i})}v,$$

so that $\omega(H_{\alpha_i})$ is an even integer.

On the other hand, $H_{\alpha_i} := \frac{2t_{\alpha_i}}{\langle \alpha_i, \alpha_i \rangle}$ by definition, where $t_{\alpha_i} \in \mathfrak{g}$ is such that $\beta(t_{\alpha_i}) = \langle \beta, \alpha_i \rangle$ for all $\beta \in \mathfrak{g}^*$. Hence, for every $i \in \{1, \dots, n-1\}$ we have

$$\frac{\langle \omega, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle} = \frac{\omega(t_{\alpha_i})}{\langle \alpha_i, \alpha_i \rangle} = \frac{1}{2}\omega(H_{\alpha_i}) \in \mathbb{Z}. \quad \square$$

Proof of Fact 17.2. Note that given an irreducible representation ρ of a reductive group with simple root system Δ , there exists a unique lowest weight of ρ relatively to the choice of Δ , and it is the highest weight relatively to the opposite simple root system $-\Delta$. Therefore we may replace the term “highest” by “lowest” in Proposition 17.9.

Let us fix $\beta \in \Theta$. Denote by $\rho : L_\Theta^0 \rightarrow \mathrm{GL}(\mathfrak{u}_\beta)$ the irreducible representation given by (16.2), (16.3) and Theorem 16.6.

By Theorem 17.8, it is enough to show that $\rho(L_\Theta^0)$ is positively proximal if and only if Conditions (i) and (ii) hold.

By Proposition 17.9 and the observation at the beginning of this proof, it is enough to check that the lowest weight space of ρ has dimension 1 if and only if Condition (i) holds, and that the lowest weight divided by 2 lies in the weight lattice of L_Θ^0 if and only if Condition (ii) holds.

By Fact 16.4, the restricted roots of L_Θ^0 have the form $\alpha|_{\mathfrak{a}_\Theta}$ with $\alpha \in \Sigma \cap \mathrm{Span}(\Delta \setminus \Theta)$, and we can choose $\{\alpha|_{\mathfrak{a}_\Theta} : \alpha \in \Delta \setminus \Theta\}$ as simple roots.

By (16.2), the weights have the form $\beta|_{\mathfrak{a}_\Theta} + \alpha|_{\mathfrak{a}_\Theta}$ where $\alpha \in \Sigma \cap \mathrm{Span}_\mathbb{N}(\Delta \setminus \Theta)$.

Observe that $\beta|_{\mathfrak{a}_\Theta}$ is by definition the lowest weight, with weight space \mathfrak{g}_β . Thus the lowest weight space of ρ has dimension 1 if and only if Condition (i) holds.

Denote by $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ and $\langle \cdot, \cdot \rangle_{\mathfrak{l}_\Theta}$ the Killing forms of respectively \mathfrak{g} and \mathfrak{l}_Θ . One may check that for all $\omega \in \mathfrak{a}^*$ and $\alpha \in \mathrm{Span}(\Delta \setminus \Theta)$ we have

$$\frac{\langle \omega, \alpha \rangle_{\mathfrak{g}}}{\langle \alpha, \alpha \rangle_{\mathfrak{g}}} = \frac{\langle \omega|_{\mathfrak{a}_\Theta}, \alpha|_{\mathfrak{a}_\Theta} \rangle_{\mathfrak{l}_\Theta}}{\langle \alpha|_{\mathfrak{a}_\Theta}, \alpha|_{\mathfrak{a}_\Theta} \rangle_{\mathfrak{l}_\Theta}} \quad (17.2)$$

Therefore $\beta|_{\mathfrak{a}_\Theta}/2$ lies in the weight lattice of L_Θ^0 if and only if for any $\alpha \in \Delta \setminus \Theta$ we have

$$2 \frac{\langle \beta, \alpha \rangle_{\mathfrak{g}}}{\langle \alpha, \alpha \rangle_{\mathfrak{g}}} \in 2\mathbb{Z}. \quad (17.3)$$

The quantity in (17.3) is worth

- 0 if there is no arrow between β and α in the Dynkin diagram of G ,
- 1 if there is a simple arrow (whatever the direction) or if there is a double or triple arrow pointing toward β ,
- 2 if there is a double arrow pointing toward α ,
- 3 if there is a triple arrow pointing toward α .

Thus (17.3) holds for all $\beta \in \Theta$ and $\alpha \in \Delta \setminus \Theta$ if and only if the only arrows in the Dynkin diagram of G between Θ and $\Delta \setminus \Theta$ are double and point toward $\Delta \setminus \Theta$, which concludes the proof. \square

18. Analysis of Groups With a Theta-Positive Structure

by Dani Kaufman

18.1. Examples of Theta-positive structures

Other than the case where G is a split real form and $\Theta = \Delta$ there are 3 new families of examples of Θ -positive structures. We will call these three cases A , B and G respectively, for reasons which will be clear later. We will explore the properties of each of these examples in more detail.

The first property to establish in each case is the structure of the positive cones in \mathfrak{u}_β . In each case, one finds that there is exactly one space, \mathfrak{u}_β , which has dimension strictly greater than 1 corresponding to the case that β is the root linked to $\Delta \setminus \Theta$ by a double arrow. We will call this root space \mathfrak{u}_β and all other one-dimensional root spaces \mathfrak{u}_{α_i} .

What is the structure of \mathfrak{u}_β in each case?

- For type A , G is a Hermitian Lie group of tube type, meaning that its Hermitian symmetric space is of the form $V + i\Omega$. The vector space \mathfrak{u}_β is isomorphic to V and the positive cone is given by Ω
- For type B , G is locally isomorphic to $SO(p, q)$. The space \mathfrak{u}_β is isomorphic to $\mathbb{R}^{1, q-p+1}$ as a normed vector space and the positive cone is given by a choice of one side of the double cone consisting of vectors in \mathfrak{u}_β with positive norm.
- For type G , the space \mathfrak{u}_β is given by the exceptional Jordan algebra of 3×3 matrices over $\mathbb{R}, \mathbb{C}, \mathbb{H}$ or \mathbb{O} and the positive cones are given by the positive definite matrices in each algebra.


18.2. The Theta-Weyl group

We define for each Θ -positive structure a subgroup of the Weyl group of G called the Θ -Weyl group, denoted $W(\Theta)$.

Let $W(\theta)$ be the subgroup generated by w_{α_i} and $w_\beta = w_{\alpha_{|\Theta|}}$ where w_{α_i} are the Weyl group generators associated to the roots $\alpha_i \in W(G)$ and w_β is the element of $W(G)$ given by the longest element of the Weyl group of $\Delta \setminus \Theta \cup \{\beta\}$

The Θ -Weyl group is isomorphic in each case to the Weyl group of a root system of type A_1, B_n, G_2 for each of the three families of non trivial Θ -positive structures, justifying the use of these names for these families. We denote the Dynkin diagram for these new root systems Δ_Θ .

The Dynkin diagrams of Δ and Δ_Θ for each family are shown in Figure 18.1.


 Figure 18.1.: Δ and Δ_Θ for each family of groups with Theta-positive structures

Remark 18.1. *The analysis of positivity in Hermitian Lie groups of tube type in Section 4.3 suggests that we should treat these groups as groups of 2×2 matrices with entries in some noncommutative ring. This agrees with the fact that Δ_Θ is an A_1 root system, which is the same root system as $\mathrm{SL}(2, \mathbb{R})$. Thus we should think of groups with a Θ -positive structure as some kind of group of type Δ_Θ with noncommutative entries.*

18.3. A Theta-positive semigroup

In analogy with Lusztig's total positivity, we can define a Θ -positive semigroup, $U_\Theta^{>0}$. To do this, we first define

$$x_{\alpha_i} : \mathfrak{u}_{\alpha_i} \rightarrow U_{\alpha_i}, v \rightarrow \exp(v)$$

Recall that the Θ -positive structure on G gives a positive cone $c_{\alpha_i}^\circ \in \mathfrak{u}_{\alpha_i}$. Let $\omega_\circ \in W(\Theta)$ be the longest element in the Θ -Weyl group. Writing ω_\circ as a product of generators $\omega_{\alpha_{i_k}}$ we define a map

$$\begin{aligned} \Phi_{\omega_\circ}^+ : \prod (c_{\alpha_{i_k}}^\circ)_{\geq 0} &\rightarrow U_\Theta^{>0} \\ \Phi_{\omega_\circ}^+(a_1, a_2, \dots, a_p) &:= x_{\alpha_{i_1}}(a_1)x_{\alpha_{i_2}}(a_2) \cdots x_{\alpha_{i_p}}(a_p). \end{aligned}$$

Theorem 18.2 (Theorem 4.5 in [GW18]). *The image $U_\Theta^{>0}$ is independent of the choice of reduced expression of ω_\circ .*

Proof. We only give the basic ideas of a proof. Since two choices of reduced expression differ by braid relations, we only need to check that the image is preserved after each type of braid relation. Each braid relation is either of the form

- (1) $\omega_{\alpha_i}\omega_{\alpha_{i+1}}\omega_{\alpha_i} = \omega_{\alpha_{i+1}}\omega_{\alpha_i}\omega_{\alpha_{i+1}}$
- (2) $\omega_\alpha\omega_\beta\omega_\alpha\omega_\beta = \omega_\beta\omega_\alpha\omega_\beta\omega_\alpha$
- (3) $\omega_\alpha\omega_\beta\omega_\alpha\omega_\beta\omega_\alpha\omega_\beta = \omega_\beta\omega_\alpha\omega_\beta\omega_\alpha\omega_\beta\omega_\alpha$

where for the last two relations α is the root in Θ adjacent to β . The first relation is between one-dimensional root spaces and is thus the same as the total positivity case. The second two relations can be established by a brute force computation in the B_2 and G_2 cases. \square

With the notion of the semigroup $U_\Theta^{>0}$ we may define the theta positive semigroup $G_\Theta^{>0}$ as the semigroup generated by $U_\Theta^{opp>0}, L_\Theta^\circ, U_\Theta^{>0}$ in a totally analogous way to the case of total positivity.

18.4. Theta-positive triples of flags

Continuing the analogies with total positivity, we can define the notion of a positive triple of elements of G/P_Θ . Let E_Θ, F_Θ be two opposite standard flags in G/P_Θ i.e. we have that the stabilizer subgroups of E and F are P_Θ^{opp} and P_Θ respectively.

Definition 18.3. Let S_Θ be a flag transverse to both E_Θ and F_Θ . We call a triple $(E_\Theta, S_\Theta, F_\Theta)$ a Θ -positive triple of flags if $S_\Theta = uE_\Theta$ for $u \in U_\Theta^{>0}$

Theorem 18.4 (Theorem 4.7 [GW18]). *The set $\{S_\Theta \in G/P_\Theta \mid (E_\Theta, S_\Theta, F_\Theta) \text{ is } \Theta\text{-positive}\}$ is a connected component of the collection of flags transverse to both E_Θ and F_Θ .*

19. Positivity in Higher Teichmüller Theory

by *Fernando Camacho Cadena*

Throughout this section G will denote a real simple Lie group, and Σ will be a compact closed surface of genus at least 2 with fundamental group $\pi_1(\Sigma)$.

We recall that a *higher Teichmüller space* is a subset of $\chi(S, G)$ which is a union of connected components consisting entirely of discrete and faithful representations. The goal of this section is to see how higher Teichmüller spaces arise from representations into different types of Lie groups.

The following were the known instances of higher Teichmüller spaces:

- When G is a split real simple group, as for example $\mathrm{PSL}(n, \mathbb{R})$, the space of Hitchin representations is a higher Teichmüller space [FG06, Lab06, Gui08].
- When G is a Hermitian Lie group of tube type, for example $\mathrm{Sp}(2n, \mathbb{R})$, the space of maximal representations is a higher Teichmüller space [BIW10].

The main goal of this section is the following theorem, which states that there are new higher Teichmüller spaces.

Theorem 19.1 ([GLW21, Theorem A]). *Let G be a simple Lie group admitting a Θ -positive structure. Then there exists a connected component of the representation variety consisting entirely of discrete and faithful representations.*

From the classification of Lie groups admitting a Θ -positive structure, the above theorem gives the existence of new higher Teichmüller spaces. Namely subsets of the representation varieties of groups isomorphic to $\mathrm{SO}(p, q)$ and Lie groups of exceptional type (which had not been known about before!).

To show the above theorem, Guichard, Labourie and Wienhard generalize the following common property shared by Hitchin and maximal representations.

Theorem 19.2 ([FG06, Lab06, Gui08, BIW10]). *Let G be either a simple Lie group which is either real split, or Hermitian of tube type. Let P_Θ be the parabolic subgroup of G corresponding to the Θ -positive structure (when G is split real, Θ is the set of simple roots, and when G is Hermitian of tube type, $\Theta = \{\alpha_r\}$, see [GW18]). Then a representation $\rho: \pi_1(\Sigma) \rightarrow G$ is Hitchin, or respectively maximal, if and only if there exists a ρ -equivariant continuous map*

$$\xi: \partial\pi_1(\Sigma) \rightarrow G/P_\Theta,$$

sending positive triples in $\partial\pi_1(\Sigma)$ to positive triples in G/P_Θ .

As we saw in previous sections, there is also a notion of positive of tuples in (generalized) flag varieties G/P_Θ . The above result then motivates the following definition.

Definition 19.3 ([GLW21, Definition 5.1]). Let G be a Lie group admitting a Θ -positive structure. A representation $\rho: \pi_1(\Sigma) \rightarrow G$ is said to be Θ -**positive** if there exists a ρ -equivariant continuous map

$$\xi: \partial\pi_1(\Sigma) \rightarrow G/P_\Theta$$

that sends positive triples in $\partial\pi_1(\Sigma)$ to positive triples in G/P_Θ .

In fact, the higher Teichmüller spaces from theorem 19.1 consist entirely of positive representations. In this section we do not present the full argument, but focus on the following

Theorem 19.4 ([GLW21, Theorem B]). *Let G be a simple Lie group admitting a Θ -positive structure, and $\rho: \pi_1(\Sigma) \rightarrow G$ a Θ -positive representation. Then ρ is also a Θ -Anosov representation.*

We will give a definition of a Θ -Anosov representation in section 19.3. We nevertheless state here two essential properties of Anosov representations, due to Labourie, and Guichard-Wienhard.

Theorem 19.5 ([Lab06, GW12]). *The space of Θ -Anosov representations is open in $\text{Hom}(\pi_1(\Sigma), G)$, and every Θ -Anosov representation is discrete and faithful.*

One can then deduce the following result.

Corollary 19.6 ([GLW21]). *The set of Θ -positive representations is an open subset of $\text{Hom}(\pi_1(\Sigma), G)$.*

Remark 19.7. *In the case when G is locally isomorphic to $\text{SO}(p, q)$, Beyrer and Pozzetti showed in [BP21] that the subset of Θ -positive representations is closed. Together with the fact from [GLW21] that the set of Θ -positive representations is open and that Θ -Anosov representations are discrete and faithful, this implies that the subset of Θ -positive representations is itself a higher Teichmüller space.*

The method in [GLW21] to prove theorem 19.1 is to find a union of connected components in the representation variety consisting entirely of Θ -positive representations. Then by corollary 19.6, all these representations are discrete and faithful. Finding a union of connected components involves methods from Higgs bundle theory as well as previous work by [BCGP⁺21]. Here we will only present the ideas for the proof of theorem 19.4.

We begin with some preliminaries on diamonds, which are structures on the flag varieties associated to the Θ -positive structure. We then move on to some definitions and facts about Anosov representations and finish with the idea of the proof of theorem 19.4.

19.1. Diamonds

For the rest of this section, we will assume that G is simple and admits a Θ -positive structure, denote by $U^{>0}$ the Θ -positive subsemigroup of G , and by F_Θ the flag variety G/P_Θ . We recall here some instances of this.

Examples 19.8. • When $G = \text{SL}(2, \mathbb{R})$,

$$U^{>0} = \left\{ \begin{pmatrix} 1 & t \\ & 1 \end{pmatrix} : t > 0 \right\}.$$

- When $G = \mathrm{SL}(3, \mathbb{R})$,

$$U^{>0} = \left\{ \begin{pmatrix} 1 & a+c & ab \\ & 1 & b \\ & & 1 \end{pmatrix} : a, b, c > 0 \right\}.$$

- When $G = \mathrm{Sp}(2n, \mathbb{R})$,

$$U^{>0} = \left\{ \begin{pmatrix} I_n & N \\ & I_n \end{pmatrix} : N \text{ is positive definite and symmetric} \right\}$$

To define positivity in the flag variety, we fix two standard flags. Namely, we let $F^\circ \in F_\Theta$ be the flag stabilized by P_Θ , and E° be the flag stabilized by P_Θ^{opp} . Recall now that if $T \in F_\Theta$ is a third flag, the triple (F°, T, E°) is *positive* if there exists an element $u_T \in U^{>0}$ such that $T = u_T E^\circ$. We say that a general triple of flags (F, T, E) is positive if there exists $g \in G$ such that

$$gF^\circ = F, \quad gE^\circ = E, \quad \text{and} \quad T = guE^\circ,$$

for some $u \in U^{>0}$. In particular this implies that for two flags F and E which are transverse, letting $g \in G$ such that $gF^\circ = F$ and $gE^\circ = E$, the subset

$$\{(F, gug^{-1}E, E) : u \in U^{>0}\}$$

consists entirely of positive triples. Here is an example of what such a set looks like.

Example 19.9. In the case of $G = \mathrm{SL}(3, \mathbb{R})$, Θ is the set of all positive roots, and we can choose

$$P_\Theta = \left\{ \begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & * \end{pmatrix} \right\},$$

making F_Θ the space $\mathrm{Flag}(\mathbb{R}^3)$ of full flags in \mathbb{R}^3 . The standard flag F° corresponds to the flag $\{\mathrm{Span}(e_1), \mathrm{Span}(e_1, e_2)\}$, and the opposite flag is $\{\mathrm{Span}(e_3), \mathrm{Span}(e_3, e_2)\}$. where e_1, e_2, e_3 are the standard basis vectors for \mathbb{R}^3 . We can visualize a full flag in \mathbb{RP}^2 as a point together with a (projective) line through that point. To picture the set $\{(F^\circ, uE^\circ, E^\circ) : u \in U^{>0}\}$, we look at a coordinate patch in \mathbb{RP}^2 :

$$\begin{aligned} \{[x : y : z] \in \mathbb{RP}^2 : x \neq 0\} &\rightarrow \mathbb{R}^2 \\ [x : y : z] &\mapsto \left[1 : \frac{y}{x} : \frac{z}{x} \right]. \end{aligned}$$

Using the explicit form of $U^{>0}$ for $\mathrm{SL}(3, \mathbb{R})$, one can see that the orbit of $\mathrm{Span}(e_1)$ is

$$\left\{ \left[1 : \frac{1}{a} : \frac{1}{ab} \right] : a, b > 0 \right\},$$

and the orbit of $\mathrm{Span}(e_2)$ is

$$\left\{ \left[1 : \frac{1}{a+c} : \frac{1}{0} \right] : a, c > 0 \right\},$$

In coordinates, the orbit of $\mathrm{Span}(e_1)$ is the (open) upper right quadrant. One can do a similar computation for the orbit of the plane $\mathrm{Span}(e_1, e_2)$ by identifying planes in \mathbb{R}^3 to orthogonal lines (equivalently by identifying $(\mathbb{RP}^2)^*$ with \mathbb{RP}^2), and then see that the orbit (in coordinates) will also be a quadrant. Now imagine ∞ in the coordinates patch in for \mathbb{RP}^2 as a single point at infinity, thereby "closing up" the upper right quadrant into a "diamond" with the standard flags as extremities.

19. Positivity in Higher Teichmüller Theory

From this vague picture, we make the following definition.

Definition 19.10. Given two transverse flags $F, E \in F_\Theta$, a **diamond** is a subset of F_Θ of the form

$$\{guE^o : u \in U^{>0}\}$$

for some $g \in G$. The **extremities** of the diamond are the flags gF^o and gE^o .

As noted above, for any T in a diamond with extremities F and E , we have that (F, T, E) is a positive triple.

Remark 19.11. *The terminology of diamond was actually coined by Labourie and Toullisse in [LT20] for the case when $G = \mathrm{SO}(2, n)$ with $n \geq 3$. There are charts on the flag variety associated to G which map to Minkowski space. In appropriate charts, a diamond is the intersection of the future and past light cone of the extremities.*

Example 19.12. When $G = \mathrm{SL}(2, \mathbb{R})$, we have the following identification

$$\mathrm{SL}(2, \mathbb{R})/P \cong S^1,$$

where P is the group of upper triangular matrices. Here, diamonds are line segments on the circle as in figure 19.1. We note here that the flag variety of $\mathrm{SL}(2, \mathbb{R})$ and the one for $\mathrm{PSL}(2, \mathbb{R})$ can be further identified with the visual boundary of \mathbb{H}^2 , its symmetric space.

There is another characterization of diamonds in terms of transverse flags. To state the result, let $\Omega_F \subset F_\Theta$ be the subset of the flag variety consisting of flags transverse to F .

Proposition 19.13 ([GLW21, Proposition 2.5]). *A diamond with extremities $E, F \in F_\Theta$ (recall here that E and F must be transverse) is a connected component of $\Omega_F \cap \Omega_E$.*

In particular, we can specify a diamond by giving two transverse flags x and z , and a point y in the interior of $\Omega_F \cap \Omega_E$. We will denote such a diamond by $V_y(x, z)$.

We will now see how diamonds are related to positive representations

Proposition 19.14 ([GLW21, Proposition 2.7]). *Assume $\xi: S^1 \rightarrow F_\Theta$ sends positive triples in S^1 to positive triples in F_Θ . Then for any oriented positive tuple (x_1, \dots, x_n) in S^1 , there exist diamonds V_{ij} such that*

- (1) $\xi(x_j) \in V_{ik}$ if (x_i, x_j, x_k) is positive, and
- (2) $V_{ij} \subset V_{km}$ if (x_k, x_i, x_j, x_m) is positive.




Figure 19.1.: The circle with the clockwise orientation for positivity, and diamonds defined by x_1, x_4 in green, and x_2, x_3 in magenta.

In figure 19.1, we think of assigning to the green and magenta arcs diamonds in the flag variety. Assertion (2) in proposition 19.14 says that the containment of diamonds in the flag variety is exactly the containment of the arcs (diamonds) on the circle.

As we will see later on, we will need metrics on (subsets) of the flag varieties. For the rest of the section, we describe how to put metrics on diamonds, and state certain contraction properties.

19.2. Metrics on diamonds

We start with a particular class of positive triples.

Definition 19.15. Let $H < G$ be a subgroup isomorphic to $\mathrm{PSL}(2, \mathbb{R})$. A *positive circle* is a closed H -orbit in F_Θ parameterized by a $\mathrm{PSL}(2, \mathbb{R})$ -equivariant map

$$\mathbb{RP}^1 \rightarrow F_\Theta.$$

Here we think of $\mathbb{RP}^1 \cong S^1$ as the flag variety of $\mathrm{PSL}(2, \mathbb{R})$ (which can also be identified with $\partial_\infty \mathbb{H}^2$).

Example 19.16. Let $G = \mathrm{PSL}(d, \mathbb{R})$, then the flag variety F_Θ can be identified with G/P , where P is the group of upper triangular matrices. Thus the flag variety consists of honest full flags. We recall that a representation $\rho: \pi_1(\Sigma) \rightarrow G$ is Fuchsian (and in particular Hitchin) if it factors through the irreducible representation $\iota: \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PSL}(d, \mathbb{R})$. We then obtain a ρ -equivariant continuous boundary map $\xi: \partial\pi_1(\Sigma) \rightarrow F_\Theta$. Then setting $H = \iota(\mathrm{PSL}(2, \mathbb{R}))$, we have that the image of the boundary map ξ is a positive circle.

Now we define a *tripod* to be a triple of pairwise distinct points lying on a positive circle. Even though the definition of a positive circle seems quite restrictive, and there is no mention of positivity, we have the following.

Proposition 19.17 ([GLW21, Proposition 2.9, Proposition 2.10]). (1) *Given two transverse points $a, b \in F_\Theta$, there exists a positive circle passing through a and b .*

(2) *The arc on a positive circle from a to b is completely contained in a diamond with extremities a and b .*

19. Positivity in Higher Teichmüller Theory

(3) *The equivariant map associated to the positive circle is positive, i.e. it sends positive triples to positive triples.*

The first part of the proposition implies that given any diamond $V_y(x, z) \subset F_\Theta$, there is a positive circle going through x and z . However, it is important to note that there might not be a positive circle passing through x, y , and z . Moreover, the third part gives that a tripod is positive.

To put a metric on a diamond, we look for a way to parameterize them. Recall that from [GW18, Theorem 4.5], the positive subsemigroup $U^{>0}$ can be parameterized by an open cone $C \subset \mathfrak{g}^N$ (where N depends on Θ). That is, there is a diffeomorphism

$$F: C \rightarrow U^{>0}.$$

Let (x, y, z) be a tripod and choose some positive circle passing through these points as well as an isomorphism $\sigma: \rightarrow U_z$, where U_z is the unipotent radical of the stabilizer of z in G . Then the map

$$\begin{aligned} C &\rightarrow F_\Theta \\ u &\mapsto \sigma \circ F(u).x \end{aligned}$$

parametrizes the diamond $V_y(x, z)$. There is a special way of choosing the isomorphism σ such that $h \mapsto y$ under the above parametrization, and where $h \in C$ is a preferred unipotent associated to the positive circle (see section 4.1 of [GLW21] for details).

Pulling back the Euclidean metric on the cone C , we get a complete Riemannian metric $g_{(x,y,z)}$ on the diamond $V_y(x, z)$.

Warning 19.18. Assume we have two tripods (x, y, z) and (x, y', z) so that the diamonds $V_y(x, z)$ and $V_{y'}(x, z)$ agree and y and y' lie on the same positive circle through x and z . If $y \neq y'$, then the metrics $g_{(x,y,z)}$ and $g_{(x,y',z)}$ need not be the same.

Given an arbitrary positive triple (x, y, z) (not necessarily a tripod), it is still possible to put a Riemannian metric $g_{(x,y,z)}$ on the diamond $V_y(x, z)$. The metric is defined by taking the average of the metrics coming from tripods that are "as close as possible" to the triple (x, y, z) . We stress again that the metric depends on the triple (x, y, z) as opposed to only on the diamond defined by the triple.

The next proposition is about how metrics for diamonds contained in each other are related.

Proposition 19.19 ([GLW21, Proposition 4.11]). *Let (x_m, y_m, z_m) be a sequence of positive triples such that*

- $V_{y_m}(x_m, z_m) \subset V_{y_{m+1}}(x_{m+1}, z_{m+1})$, and
- $\bigcap_{m \in \mathbb{N}} V_{y_m}(x_m, z_m) = \{\bullet\}$.

Then

$$g_{(x_0, y_0, z_0)} \leq k_m g_{(x_m, y_m, z_m)},$$

with $k_m \rightarrow 0$ as $m \rightarrow \infty$.

We do not reproduce the proof here, but remark that the vague intuition behind this statement is as follows. The metric on diamonds is defined by the pullback to the cone C . Thus, if a diamond V is contained in another diamond W , the metric in W has to be larger.

19.3. Basics of Anosov representations

In this section, we give a definition of Anosov representations which will be useful for us, but do not give the full details.

Begin by choosing an auxiliary hyperbolic metric on Σ and with this there is an identification

$$T^1\mathbb{H}^2/\pi_1(\Sigma) \cong T^1\Sigma,$$

where $T^1\mathbb{H}^2$ is the unit tangent bundle of the hyperbolic plane, and $T^1\Sigma$ is the unit tangent bundle to Σ . Then we identify $T^1\mathbb{H}^2$ with the set of positive triples in $\partial\mathbb{H}^2$ as follows. Positivity in $\partial\mathbb{H}^2$ is inherited from the identification with the circle S^1 . Let (x, y, z) be a positive triple and consider the oriented geodesic c from x to z in \mathbb{H}^2 . Then there is a unique geodesic with y as an endpoint intersecting c orthogonally. Thus y determines a point on the geodesic, and with that, a vector in the unit tangent bundle. Note that the triple (z, y, x) is also positive and gives the negative of the vector determined by (x, y, z) . Moreover, the geodesic flow will be denoted by $(x, y, z) \mapsto (x, y_s, z)$, where s is the time parameter. See figure 19.2.




Figure 19.2.: Parametrization of unit tangent bundle with positive triples and the clockwise orientation on the circle. Here the violet geodesic with x as the forward endpoint, and z as the backward endpoint. The blue geodesics meet it orthogonally to determine a vector in $T^1\mathbb{H}^2$. In the picture, $s_2 < s_1$.

The following is an equivalent definition of Anosov representations given by Guichard and Wienhard in [GW12, Proposition 2.7].

Definition 19.20. A representation $\rho: \pi_1(\Sigma) \rightarrow G$ is Θ -Anosov if the following conditions hold.

- (1) There exists a continuous ρ -equivariant map $\xi: \partial\pi_1(T^1\Sigma) \rightarrow F_\Theta$ sending distinct points $x, y \in \partial\pi_1(T^1\Sigma)$ to transverse points in F_Θ .
- (2) There exists a continuous $\pi_1(\Sigma)$ -equivariant family of norms on

$$\{(T_{\xi(z)}F_\Theta, (x, y, z)) : (x, y, z) \in T^1\mathbb{H}^2\}$$

satisfying the following. There exist positive constants A and a such that for all $t > 0$, and $(x, y, z) \in T^1\mathbb{H}^2$

$$\|\cdot\|_{(x, y, z)} \leq Ae^{-at} \|\cdot\|_{(x, y_{-s}, z)}, \quad \text{for all } s \geq 0.$$

To be precise, the definition should include a second boundary map to the flag variety F/P_{Θ}^{opp} , but we only work with one boundary map. Moreover, the continuity for the family of norms in condition (2) is with respect to taking the trivial bundle $T^1\mathbb{H}^2 \times F_{\Theta}$.

19.4. Proof sketch of Theorem 19.4

Part (1) of the definition of Anosov representations is immediate from the definition of positive representations. For part (2), let $\xi: \partial\mathbb{H}^2 \rightarrow F_{\Theta}$ be the boundary map associated to the positive representation. The goal is to define diamonds with $\xi(z)$ in its interior, so that we can assign the norm on $T_{\xi(z)}F_{\Theta}$ coming from the Riemannian metric on the diamond.

We begin by assigning to $(x, y, z) \in T^1\mathbb{H}^2$ a fourth element $w(x, y, z) \in \partial\mathbb{H}^2$ satisfying

$$\text{cr}_{\mathbb{RP}^1}(x, y, z, w(x, y, z)) = -1,$$

where $\text{cr}_{\mathbb{RP}^1}$ is a cross ratio on $\mathbb{RP}^1 \cong \partial\mathbb{H}^2$. In the literature, such a quadruple is said to be harmonic. One can easily check that the quadruple $(x, y, z, w(x, y, z))$ is a positive quadruple, see figure 19.3. For example, taking the usual cross ratio on \mathbb{RP}^1 , the quadruple $(0, 1, \infty, -1)$ (in coordinates defined by normalizing the first coordinate of the line) is harmonic. We then assign to the triple (x, y, z) the diamond

$$\mathcal{Y}_{(x,y,z)} := V_{\xi(z)}(\xi(y), \xi(w(x, y, z))) \subset F_{\Theta}.$$

The point $\xi(z)$ lies in the interior of the above diamond. This means that we can denote by $\|\cdot\|_{(x,y,z)}$ the norm on $T_{\xi(z)}F_{\Theta}$ coming from the Riemannian metric $g_{(\xi(y), \xi(z), \xi(w(x, y, z)))}$. To see that the metrics are expanding along the geodesic flow, we use proposition 19.19. The conditions that need to be satisfied are

- (1) $\mathcal{Y}_{(x,y-s,z)} \subset \mathcal{Y}_{(x,y,z)}$ for all $s \geq 0$, and
- (2) $\mathcal{Y}_{(x,y-s,z)} \rightarrow \{\xi(z)\}$ as $s \rightarrow +\infty$.

For condition (1), one can check with a small computation that if (x, y_1, y_0, z) is a positive quadruple, then $(x, y_1, y_0, z, w(x, y_0, z), w(x, y_1, z))$ is a positive tuple as well, see figure 19.3. Since the boundary map is positive, it follows from proposition 19.14 that indeed $\mathcal{Y}_{(x,y-s,z)} \subset \mathcal{Y}_{(x,y,z)}$ for all $s \geq 0$.




Figure 19.3.: Harmonic quadruples on S^1 .

For condition (2), observe (from another calculation) that

$$\lim_{y \rightarrow z} w(x, y, z) = z.$$

Then using the continuity of the boundary map, and that $y_{-s} \rightarrow z$ as $s \rightarrow \infty$, we get condition (2). Now we can apply proposition 19.19 to get that

$$\|\cdot\|_{(x,y,z)} \leq k_s \|\cdot\|_{(x,y_{-s},z)}$$

with $k_s \rightarrow 0$ as $s \rightarrow \infty$. One can then finish the proof by using a compactness argument to get the constants A and a in the definition of Anosov representations.

20. Exercises

by Merik Niemeyer

Our goal in this exercise is to find Θ -positive structures on both $\mathrm{SO}(2, 3)$ and $\mathrm{Sp}(4, \mathbb{R})$.

20.1. The indefinite special orthogonal group

Let us first introduce the setting. A lot of this you have seen already in the talk on $\mathrm{SO}(p, q)$.

Define a quadratic form on \mathbb{R}^5 by $x \mapsto {}^t x Q x$ with

$$Q = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Then $G := \mathrm{SO}(Q) = \{X \in \mathrm{SL}(5, \mathbb{R}) : {}^t X Q X = Q\} \cong \mathrm{SO}(2, 3)$. Recall that matrices in the Lie algebra $\mathfrak{g} = \mathfrak{so}(2, 3)$ are of the form

$$\left(\begin{array}{cc|c|cc} a_{11} & a_{12} & v & b & 0 \\ a_{21} & a_{22} & w & 0 & b \\ \hline r & s & 0 & w & -v \\ \hline c & 0 & s & -a_{22} & a_{12} \\ 0 & c & -r & a_{21} & -a_{11} \end{array} \right).$$

with real entries. You have also seen already that a Cartan subspace is given by

$$\mathfrak{h} = \{\mathrm{diag}(\lambda, \mu, 0, -\mu, -\lambda) : \lambda, \mu \in \mathbb{R}\} = \underbrace{\mathbb{R} \mathrm{diag}(1, 0, 0, 0, -1)}_{=: h_1} \oplus \underbrace{\mathbb{R} \mathrm{diag}(0, 1, 0, -1, 0)}_{=: h_2},$$

i.e. the diagonal matrices in \mathfrak{g} . Define the matrices

$$e_1 := \begin{pmatrix} 0 & 1 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \\ \hline 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 1 \\ 0 & 0 & | & 0 & 0 & 0 \end{pmatrix}, \quad e_2 := \begin{pmatrix} 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 1 & 0 & 0 \\ \hline 0 & 0 & | & 0 & 1 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \end{pmatrix},$$

$$e_3 := \begin{pmatrix} 0 & 0 & | & 1 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \\ \hline 0 & 0 & | & 0 & 0 & -1 \\ 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \end{pmatrix}, \quad e_4 := \begin{pmatrix} 0 & 0 & | & 0 & 1 & 0 \\ 0 & 0 & | & 0 & 0 & 1 \\ \hline 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \\ 0 & 0 & | & 0 & 0 & 0 \end{pmatrix},$$

and $f_i := {}^t e_i$. Together with h_1 and h_2 these span the Lie algebra.

Remark 20.1. Note that \mathfrak{g} is generated by $h_1, h_2, e_1, e_2, f_1, f_2$.

Define two functions $\alpha_1, \alpha_2 \in \mathfrak{h}^*$ by

$$\begin{aligned}\alpha_1(\text{diag}(\lambda, \mu, 0, -\mu, -\lambda)) &:= \lambda - \mu, \\ \alpha_2(\text{diag}(\lambda, \mu, 0, -\mu, -\lambda)) &:= \mu.\end{aligned}$$

Exercise 20.1. Compute the root space decomposition of \mathfrak{g} and prove that \mathfrak{g} has a B_2 root system.

In case you do not want to spend time on this repetition, here are the positive root spaces:

$$\mathfrak{g}_{\alpha_1} = \mathbb{R}e_1, \quad \mathfrak{g}_{\alpha_2} = \mathbb{R}e_2, \quad \mathfrak{g}_{\alpha_1+\alpha_2} = \mathbb{R}e_3, \quad \mathfrak{g}_{\alpha_1+2\alpha_2} = \mathbb{R}e_4.$$

The corresponding negative root spaces are spanned by the f_i and $\mathfrak{g}_0 = \mathfrak{h}$ since G is split. Also, we see that $\Delta = \{\alpha_1, \alpha_2\}$ is a set of simple roots.

20.1.1. The totally positive structure

The first Θ -positive structure we want to investigate is the totally positive structure, i.e. pick $\Theta = \Delta$. However, we will look at this from the viewpoint of Θ -positivity.

Exercise 20.2. For this choice of Θ compute $\mathfrak{u}_\Theta, \mathfrak{u}_\Theta^{\text{opp}}$ and \mathfrak{l}_Θ . Find the center \mathfrak{z}_Θ of \mathfrak{l}_Θ and the weight spaces \mathfrak{u}_α for the (adjoint) action of this on \mathfrak{u}_Θ .

Remark 20.2. Note that the weight spaces coincide with the root spaces and that all of these are 1-dimensional.

In a second step, we wish to prove that G actually admits a Θ -positive structure for this choice of Θ , i.e. we need to find invariant cones in the weight spaces:

Exercise 20.3. By exponentiating, find L_Θ° (the identity component of the Levi subgroup) and compute the action on \mathfrak{u}_α for $\alpha \in \Theta$. Prove that there exists a sharp convex cone in each weight space, which is invariant under this action.

With the Θ -positive structure in place, we will compute the Θ -positive semigroup.

Exercise 20.4. Observe that the Weyl group $W(\Theta)$ is just the full Weyl group W . Use the longest word to give a parametrization $U_\Theta^{>0}$.

Hint 20.1. The longest word in the Weyl group of a B_2 root system is $\sigma_1\sigma_2\sigma_1\sigma_2 = \sigma_2\sigma_1\sigma_2\sigma_1$. It might not be worth it to explicitly compute a matrix form of an element in $U_\Theta^{>0}$.

Now you could compute $U_\Theta^{\text{opp}, >0}$ in much the same way and finally get $G_\Theta^{>0}$.

Remark 20.3. Note that this Θ -positive semigroup coincides with Lusztig's totally positive semigroup.

Exercise 20.5. As a non-example, show that $\text{SO}(2, q)$ for $q > 3$ does not admit a Θ -positive structure if we chose $\Theta = \Delta$, the set of simple (restricted) roots.

20. Exercises

Another Θ -positive structure

Now we will define a second Θ -positive structure on G by choosing $\Theta = \{\alpha_1\}$.

Remark 20.4. Recall the classification of Lie groups admitting a Θ -positive structure. As G is both split real and (locally isomorphic to) $\mathrm{SO}(p, q)$, we find two different Θ -positive structures on G .

We can proceed in much the same way as before.

Exercise 20.6. Prove that G admits a Θ -positive structure by explicitly calculating the L_Θ° -invariant cone in \mathfrak{u}_{α_1} .

Hint 20.2. You should find that elements of L_Θ° are of the form

$$\begin{pmatrix} a & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & a^{-1} \end{pmatrix}$$

where $a \in \mathbb{R}_{>0}$ and $M \in \mathrm{SO}(J)^\circ$ (here we denote by J the center 3×3 -block in Q) and that elements of $\mathfrak{u}_{\alpha_1} = \mathfrak{u}_\Theta$ are of the form

$$\begin{pmatrix} 0 & {}^t v & 0 \\ 0 & 0 & Jv \\ 0 & 0 & 0 \end{pmatrix}$$

for some $v \in \mathbb{R}^3$. The sharp convex cone will be defined by the conditions ${}^t v J v \geq 0$ and $v_1 \geq 0$.

We can also compute the Weyl group $W(\Theta)$ in order to define $U_\Theta^{>0}$ (and from this $G_\Theta^{>0}$ like before):

Exercise 20.7. Compute the Weyl group $W(\Theta)$ and show that this is isomorphic to the Weyl group of an A_1 root system and give a parametrization of $U_\Theta^{>0}$.

Remark 20.5. We have also learned that G is of Hermitian type of tube type. The Θ -positive structure it admits as such a group is the same one we just described.

20.1.2. Another Θ -positive structure?

Exercise 20.8. Show that for the choice $\Theta = \{\alpha_2\}$, G does **not** admit a Θ -positive structure.

20.2. The symplectic group

Define $J_{2,2}$ as the block matrix

$$J_{2,2} = \begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix}$$

and recall that the symplectic group is $G' := \mathrm{Sp}(4, \mathbb{R}) = \{M \in \mathrm{GL}(4, \mathbb{R}) : {}^t M J_{2,2} M = J_{2,2}\}$. Its Lie algebra is $\mathfrak{g}' = \mathfrak{sp}(4, \mathbb{R}) = \{X \in \mathfrak{gl}(4, \mathbb{R}) : {}^t X J_{2,2} + J_{2,2} X = 0\}$, so an element in \mathfrak{g}' is of the form

$$X = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

where C, B are symmetric (2×2) -matrices and $D = -{}^t A$. A Cartan subspace is given by

$$\mathfrak{h}' = \{\mathrm{diag}(\lambda, \mu, -\lambda, -\mu) : \lambda, \mu \in \mathbb{R}\}$$

Exercise 20.9. Compute the root decomposition of \mathfrak{g}' and prove that \mathfrak{g}' has a B_2 root system.

20.2.1. A local isomorphism with $\mathrm{SO}(2, 3)$

We observed that the root systems of G and G' are both of type B_2 and thus isomorphic. It is not hard to write this isomorphism down explicitly and starting from this we can construct an isomorphism of the Lie algebras \mathfrak{g} and \mathfrak{g}' . The resulting isomorphism $\psi: \mathfrak{g} \rightarrow \mathfrak{g}'$ is defined on the generators of \mathfrak{g} by:

$$\begin{aligned} \psi(h_1) &= \mathrm{diag}(1/2, 1/2, -1/2, -1/2) =: h'_1, \\ \psi(h_2) &= \mathrm{diag}(1/2, -1/2, 1/2, -1/2) =: h'_2, \\ \psi(e_1) &= \left(\begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) =: e'_1, \\ \psi(e_2) &= \left(\begin{array}{cc|cc} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array} \right) =: e'_2, \\ \psi(f_1) &= (e'_1)^T =: f'_1, \\ \psi(f_2) &= (e'_2)^T/2 =: f'_2. \end{aligned}$$

Under this isomorphism the simple roots α_1 and α_2 correspond to $\alpha'_1, \alpha'_2 \in \mathfrak{h}'$, respectively, which are defined by

$$\begin{aligned} \alpha'_1(\mathrm{diag}(\lambda, \mu, -\lambda, -\mu)) &= 2\mu, \\ \alpha'_2(\mathrm{diag}(\lambda, \mu, -\lambda, -\mu)) &= \lambda - \mu. \end{aligned}$$

Remark 20.6. This isomorphism of Lie algebras induces a local isomorphism between the Lie groups G and G' .

20. Exercises

20.2.2. Θ -positive structures on $\mathrm{Sp}(4, \mathbb{R})$

Here we only consider the Θ -positive structure, which G' carries as a group locally isomorphic to $\mathrm{SO}(2, 3)$ (see above), which is actually the same as the one which it carries as a Hermitian group of tube type. We can easily carry over our previous calculations using the above isomorphism.

Exercise 20.10. Use the isomorphism $\psi: \mathfrak{g} \rightarrow \mathfrak{g}'$ to prove that G' admits a Θ -positive structure for the choice $\Theta = \{\alpha'_1\}$.

Hint 20.3. You should find that the weight space $\mathfrak{u}_{\alpha'_1}$ is isomorphic to the space of symmetric matrices and that the invariant cone is given by the positive semi-definite matrices.

Some notes (almost solutions) on the exercises.

20.3. Partial solutions

20.3.1. $\mathrm{SO}(2, 3)$

The totally positive structure

Define a quadratic form on \mathbb{R}^5 by $x \mapsto {}^t x Q x$ with

$$Q = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Then $G := \mathrm{SO}(Q) = \{M \in \mathrm{SL}(5, \mathbb{R}) : {}^t M Q M = Q\} \cong \mathrm{SO}(2, 3)$ and matrices in the Lie algebra $\mathfrak{g} = \mathfrak{so}(2, 3)$ are of the form

$$\left(\begin{array}{cc|c|cc} a_{11} & a_{12} & v & b & 0 \\ a_{21} & a_{22} & w & 0 & b \\ \hline r & s & 0 & w & -v \\ \hline c & 0 & s & -a_{22} & a_{12} \\ 0 & c & -r & a_{21} & -a_{11} \end{array} \right).$$

To work out the root decomposition we pick the maximal toral subalgebra

$$\mathfrak{h} = \{\mathrm{diag}(\lambda, \mu, 0, -\mu, -\lambda) : \lambda, \mu \in \mathbb{R}\} = \underbrace{\mathbb{R} \mathrm{diag}(1, 0, 0, 0, -1)}_{=: h_1} \oplus \underbrace{\mathbb{R} \mathrm{diag}(0, 1, 0, -1, 0)}_{=: h_2},$$

i.e. the diagonal matrices in \mathfrak{g} .

Now we will investigate the root spaces: For this, define the following matrices:

$$e_1 := \left(\begin{array}{c|c|c|c} 0 & 1 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \end{array} \right), \quad e_2 := \left(\begin{array}{c|c|c|c} 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 \\ \hline 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right),$$

$$e_3 := \left(\begin{array}{c|c|c|c} 0 & 0 & 1 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & -1 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right), \quad e_4 := \left(\begin{array}{c|c|c|c} 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right).$$

These form a linear basis for the subspace of strictly upper triangular matrices in \mathfrak{g} . In fact this subspace is a subalgebra and as such is generated by e_1 and e_2 .

Moreover, each e_i spans a positive root spaces: Define the functions $\alpha_1, \alpha_2 \in \mathfrak{h}^*$ by

$$\alpha_1(\text{diag}(\lambda, \mu, 0, -\mu, -\lambda)) := \lambda - \mu,$$

$$\alpha_2(\text{diag}(\lambda, \mu, 0, -\mu, -\lambda)) := \mu,$$

and hence the positive root spaces are

$$\mathfrak{g}_{\alpha_1} = \mathbb{R}e_1, \quad \mathfrak{g}_{\alpha_2} = \mathbb{R}e_2, \quad \mathfrak{g}_{\alpha_1+\alpha_2} = \mathbb{R}e_3, \quad \mathfrak{g}_{\alpha_1+2\alpha_2} = \mathbb{R}e_4.$$

The corresponding negative root spaces are spanned by the matrices $f_i := {}^t e_i$ for $i \in \{1, 2, 3, 4\}$. We thus find that this is a B_2 root system with α_1 and α_2 a pair of simple roots.

We first pick $\Theta = \Delta$, so that

$$\mathfrak{u}_\Theta = \sum_{\alpha \in \Sigma^+} \mathfrak{g}_\alpha,$$

$$\mathfrak{u}_\Theta^{\text{opp}} = \sum_{\alpha \in \Sigma^+} \mathfrak{g}_{-\alpha},$$

and clearly in this case $\mathfrak{l}_\Theta = \mathfrak{z}_\Theta = \mathfrak{g}_0 = \mathfrak{h}$. Consequently, the weight spaces for the action $\mathfrak{z}_\Theta \curvearrowright \mathfrak{u}_\Theta$ coincide with the root spaces and we find $\mathfrak{u}_{\alpha_i} = \mathfrak{g}_{\alpha_i}$. By exponentiating elements in \mathfrak{l}_Θ we find that the identity component of the Levi subgroup is

$$L_\Theta^\circ = \{\text{diag}(a, b, 1, b^{-1}, a^{-1}) : a, b \in \mathbb{R}_+\}.$$

If we let this act on \mathfrak{u}_{α_1} by the adjoint action, we find that $X = \text{diag}(a, b, 1, b^{-1}, a^{-1})$ maps e_1 to $ab^{-1}e_1$. Thus, if we identify the one-dimensional space \mathfrak{u}_{α_1} with \mathbb{R} , we find that the sharp convex cone \mathbb{R}_+ is preserved by this action ($a, b > 0$). Analogously for \mathfrak{u}_{α_2} , and we proved that G admits a Θ -positive structure.

Exponentiating the root spaces for roots in Θ yields two maps

$$x_i : \mathbb{R} \rightarrow G$$

$$t \mapsto \exp(te_i),$$

for $i \in \{1, 2\}$. More explicitly, these map to the subgroup U_Θ of G of unipotent upper triangular matrices. As we choose $\Theta = \Delta$, we find that $W(\Theta) = W$, i.e. the Weyl group of the root

20. Exercises

system of G , which is of type B_2 . The longest word in such a root system can be written as $\sigma_1\sigma_2\sigma_1\sigma_2$, and therefore we obtain a parametrization of the totally positive subsemigroup $U_\Theta^{>0}$ of U_Θ , namely

$$U_\Theta^{>0} := \{x_1(x)x_2(v)x_1(y)x_2(w) =: F_{1212}(x, v, y, w) : x, v, y, w \in \mathbb{R}^{>0}\}.$$

An element in this can be explicitly computed to be

$$F_{1212}(x, v, y, w) = \begin{pmatrix} 1 & x+y & xv + (x+y)w & x(v+w)^2/2 + yw^2/2 & xyv^2/2 \\ 0 & 1 & v+w & (v+w)^2/2 & yv^2/2 \\ 0 & 0 & 1 & v+w & yv \\ 0 & 0 & 0 & 1 & x+y \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Analogously one can compute the positive part $O^{>0}$ of the subgroup O of unipotent lower triangular matrices. The totally positive subsemigroup $G^{>0}$ is then

$$G^{>0} = O^{>0}A^\circ U^{>0},$$

where A° is the connected component of the identity in the subgroup $A < G$ of diagonal matrices in G .

It is not hard to see that the positive semigroup defined this way is just Lusztig's totally positive subgroup and much of the discussion actually mirrors Lusztig's description. However, this is only possible because we are dealing with a split real group.

If we naively tried to extend this notion to non-split groups we run into problems as the following example shows: Consider $G_2 = \text{SO}(2, q)$ with $q > 3$. The Lie algebra then has the form

$$\mathfrak{so}(2, q) = \left\{ \left(\begin{array}{cc|cc|cc} a_{11} & a_{12} & {}^t v & b & 0 & \\ a_{21} & a_{22} & {}^t w & 0 & b & \\ \hline r & s & 0 & w & -v & \\ \hline c & 0 & {}^t s & -a_{22} & a_{12} & \\ 0 & c & -{}^t r & a_{21} & -a_{11} & \end{array} \right) : a_{ij}, b, c \in \mathbb{R}, v, w, r, s \in \mathbb{R}^{q-2} \right\},$$

and we recall that the restricted root system of this group is also of type B_2 and indeed that calculation is very similar to the above but we find that the (restricted) root space \mathfrak{g}_{α_2} is $(q-2)$ -dimensional in this case and contains all elements of the Lie algebra of the above form with all but the w -entries 0. Now, if we take $\Theta = \Delta$, \mathfrak{u}_Θ and $\mathfrak{u}_\Theta^{\text{opp}}$ are again just the sum of all positive and negative root spaces, respectively, and we also find $\mathfrak{l}_\Theta = \mathfrak{g}_0$ again. However, this takes a slightly different form since G_2 is not split.

$$\mathfrak{l}_\Theta = \mathfrak{g}_0 = \mathfrak{a} \oplus C_{\mathfrak{k}}(\mathfrak{a}) = \left\{ \left(\begin{array}{ccc} \lambda & & \\ & \mu & \\ & M & -\mu \\ & & & -\lambda \end{array} \right) : M = -{}^t M \right\},$$

where \mathfrak{a} is a Cartan subspace and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the Cartan decomposition. The center clearly is $\mathfrak{z}_\Theta = \mathfrak{a}$ and thus the weight spaces for the action $\mathfrak{z}_\Theta \curvearrowright \mathfrak{u}_\Theta$ are just the root spaces again. Exponentiating the above expression, we find that a general element in L_Θ° has the form

$$Y = \begin{pmatrix} a & & & & \\ & b & & & \\ & & S & & \\ & & & b^{-1} & \\ & & & & a^{-1} \end{pmatrix}$$

with $S \in \text{SO}(q-2)$. If we consider the adjoint action on the weight space $\mathfrak{u}_{\alpha_2} = \mathfrak{g}_{\alpha_2} \cong \mathbb{R}^{q-2}$, we find $Y.v = bSv$, so that we are effectively dealing with an $\text{SO}(q-2)$ -action on \mathbb{R}^{q-2} , which clearly has no invariant sharp convex cone, so that G_2 does not admit a Θ -positive structure for this particular choice of Θ .

Another Θ -positive structure

The group $G = \text{SO}(2,3)$ can also be endowed with another Θ -positive structure. For this, we mostly use the same setup as before and have $\Delta = \{\alpha_1, \alpha_2\}$. Now we pick $\Theta = \{\alpha_1\}$ and thus have (with $\Sigma_{\Theta}^+ = \Sigma^+ \setminus (\text{Span}(\Delta - \Theta))$, i.e. all positive roots that contain some contribution from Θ)

$$\begin{aligned} \mathfrak{u}_{\Theta} &:= \sum_{\alpha \in \Sigma_{\Theta}^+} \mathfrak{g}_{\alpha} = \mathfrak{g}_{\alpha_1} \oplus \mathfrak{g}_{\alpha_1 + \alpha_2} \oplus \mathfrak{g}_{\alpha_1 + 2\alpha_2} = \mathbb{R}e_1 \oplus \mathbb{R}e_3 \oplus \mathbb{R}e_4, \\ \mathfrak{u}_{\Theta}^{\text{opp}} &:= \sum_{\alpha \in \Sigma_{\Theta}^+} \mathfrak{g}_{-\alpha} = \mathfrak{g}_{-\alpha_1} \oplus \mathfrak{g}_{-\alpha_1 - \alpha_2} \oplus \mathfrak{g}_{-\alpha_1 - 2\alpha_2} = \mathbb{R}f_1 \oplus \mathbb{R}f_3 \oplus \mathbb{R}f_4, \\ \mathfrak{l}_{\Theta} &:= \mathfrak{g}_0 \oplus \sum_{\alpha \in \text{Span}(\Delta - \Theta) \cap \Sigma^+} (\mathfrak{g}_{\alpha} \oplus \mathfrak{g}_{-\alpha}) = \mathfrak{h} \oplus \mathfrak{g}_{\alpha_2} \oplus \mathfrak{g}_{-\alpha_2}. \end{aligned}$$

In order to determine the Θ -positive subsemigroup, we need to consider the adjoint action of the center \mathfrak{z}_{Θ} of \mathfrak{l}_{Θ} on \mathfrak{u}_{Θ} . In this simple setup, we clearly have

$$\mathfrak{z}_{\Theta} = \{\text{diag}(\lambda, 0, 0, 0, -\lambda) : \lambda \in \mathbb{R}\},$$

and thus $\mathfrak{u}_{\Theta} = \mathfrak{u}_{\alpha_1}$.

By exponentiating an element in \mathfrak{l}_{Θ} we see that a general element in the identity component of the Levi subgroup L_{Θ} is of the form

$$\begin{pmatrix} a & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & a^{-1} \end{pmatrix}$$

where $a \in \mathbb{R}^{>0}$ and $M \in \text{SO}(J)^{\circ}$. Here we denote by J the center (3×3) -block in Q , i.e.

$$J = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0^1 \end{pmatrix}.$$

Now, elements of $\mathfrak{u}_{\alpha_1} = \mathfrak{u}_{\Theta}$ are of the form

$$\begin{pmatrix} 0 & {}^t v & 0 \\ 0 & 0 & Jv \\ 0 & 0 & 0 \end{pmatrix}$$

for some $v \in \mathbb{R}^3$. Therefore, we identify this space with \mathbb{R}^3 and letting L_{Θ}° act on \mathfrak{u}_{α_1} by conjugation, we find that the sharp convex cone c defined by the conditions ${}^t v J v \geq 0$ and $v_1 \geq 0$ is invariant under this action. So G admits a Θ -positive structure for the above choice of Θ .

Our goal is to find the Θ -positive subsemigroup $G_{\Theta}^{>0}$ and for this we need to consider the Weyl group $W(\Theta)$. In our case this is very simple as we have $\Theta = \{\alpha_1\}$, which only contains a single

20. Exercises

element. Thus $\beta_\Theta = \alpha_1$ and $W(\Theta)$ is the subgroup of the full Weyl group W that is generated by the longest element s of the Weyl group $W_{\{\beta_\Theta\} \cup (\Delta - \Theta)} = W$. This is isomorphic to the Weyl group of an A_1 root system and thus the longest element in $W(\Theta)$ is simply s .

Accordingly the Θ -positive subsemigroup of $U_\Theta = \exp(\mathfrak{u}_\Theta)$ is the image of the map

$$F : c^\circ \rightarrow U_\Theta$$

$$v \mapsto \exp(v),$$

i.e. $U_\Theta^{>0} = F(c^\circ)$.

Similarly, we can compute $U_\Theta^{\text{opp}, >0}$ and thus find $G_\Theta^{>0}$, which is generated by $U_\Theta^{>0}, U_\Theta^{\text{opp}, >0}$ and L_Θ° .

Another Θ -positive structure?

Finally, we want to see whether we can find another Θ -positive structure on G by taking $\Theta = \{\alpha_2\}$ (keep the notation from before). The calculation works much the same as in the previous case but we find $\mathfrak{u}_\Theta = \mathfrak{u}_{\alpha_2} \oplus \mathfrak{u}_{2\alpha_2}$, where $\mathfrak{u}_{\alpha_2} = \mathbb{R}e_2 \oplus \mathbb{R}e_3 \cong \mathbb{R}^2$. A calculation shows that the action $L_\Theta^\circ \curvearrowright \mathfrak{u}_{\alpha_2}$ corresponds to an $\text{SL}(2, \mathbb{R})$ -action on \mathbb{R}^2 , which has no sharp invariant cone and so G does not admit a Θ -positive structure for this choice of Θ .

20.3.2. $\mathbf{Sp}(4, \mathbb{R})$

A local isomorphism with \mathbf{G}

Define $J_{2,2}$ as the block matrix

$$J_{2,2} = \begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix}$$

and recall that the symplectic group is $G' := \text{Sp}(4, \mathbb{R}) = \{M \in \text{GL}(4, \mathbb{R}) : {}^t M J_{2,2} M = J_{2,2}\}$. Its Lie algebra is $\mathfrak{g}' = \mathfrak{sp}(4, \mathbb{R}) = \{X \in \mathfrak{gl}(4, \mathbb{R}) : {}^t X Q + Q X = 0\}$, so an element in \mathfrak{g}' is of the form

$$X = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

where C, B are symmetric (2×2) -matrices and $D = -{}^t A$. A maximal toral subalgebra is given by

$$\mathfrak{h}' = \{\text{diag}(\lambda, \mu, -\lambda, -\mu) : \lambda, \mu \in \mathbb{R}\} = \underbrace{\mathbb{R} \text{diag}(1, 0, -1, 0)}_{=: \tilde{h}_1} \oplus \underbrace{\mathbb{R} \text{diag}(0, 1, 0, -1)}_{=: \tilde{h}_2}.$$

As a vector space \mathfrak{g}' is spanned by the matrices

$$\begin{aligned}\tilde{e}_1 &:= \left(\begin{array}{cc|cc} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array} \right), & \tilde{e}_2 &:= \left(\begin{array}{cc|cc} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right), \\ \tilde{e}_3 &:= \left(\begin{array}{cc|cc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right), & \tilde{e}_4 &:= \left(\begin{array}{cc|cc} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),\end{aligned}$$

and their transposed matrices $\tilde{f}_i = \tilde{e}_i^T$. If we set $X = \lambda\tilde{h}_1 + \mu\tilde{h}_2 = \text{diag}(\lambda, \mu, -\lambda, -\mu)$ and compute the bracket of X with the \tilde{e}_i , we find

$$\begin{aligned}[X, \tilde{e}_1] &= (\lambda - \mu)\tilde{e}_1, \\ [X, \tilde{e}_2] &= 2\lambda\tilde{e}_2, \\ [X, \tilde{e}_3] &= 2\mu\tilde{e}_3, \\ [X, \tilde{e}_4] &= (\lambda + \mu)\tilde{e}_4,\end{aligned}$$

and thus if we set

$$\begin{aligned}\tilde{\alpha}_1(X) &= \lambda - \mu, \\ \tilde{\alpha}_2(X) &= 2\mu,\end{aligned}$$

we have the following root spaces

$$\mathfrak{g}_{\tilde{\alpha}_1} = \mathbb{R}\tilde{e}_1, \quad \mathfrak{g}_{\tilde{\alpha}_2} = \mathbb{R}\tilde{e}_3, \quad \mathfrak{g}_{\tilde{\alpha}_1 + \tilde{\alpha}_2} = \mathbb{R}\tilde{e}_4, \quad \mathfrak{g}_{2\tilde{\alpha}_1 + \tilde{\alpha}_2} = \mathbb{R}\tilde{e}_2.$$

The corresponding negative root spaces are spanned by the \tilde{f}_i and we are clearly dealing with another root system of type B_2 .

Therefore the root systems of G and G' are isomorphic, which means that the Lie algebras \mathfrak{g} and \mathfrak{g}' are isomorphic. The isomorphism of the root systems is given in terms of the simple roots as

$$\begin{aligned}\alpha_1 &\mapsto \tilde{\alpha}_2 =: \alpha'_1, \\ \alpha_2 &\mapsto \tilde{\alpha}_1 =: \alpha'_2,\end{aligned}$$

which can be seen by comparing the expressions for the positive roots in terms of the simple roots in both cases. In order to construct a Lie algebra isomorphism $\psi : \mathfrak{g} \rightarrow \mathfrak{g}'$, recall that \mathfrak{g} is generated by $h_1, h_2, e_1, e_2, f_1, f_2$. In order to define ψ it suffices to map these generators in such a way that the bracket is preserved (this gives as a Lie algebra homomorphism).

We begin by mapping e_1, e_2 : These span the root spaces \mathfrak{g}_{α_1} and \mathfrak{g}_{α_2} , respectively, and thus we need to map these to elements in $\mathfrak{g}'_{\alpha'_2}$ and $\mathfrak{g}'_{\alpha'_1}$, respectively. Thus we define

$$\begin{aligned}\psi(e_1) &= \tilde{e}_3 =: e'_1, \\ \psi(e_2) &= \tilde{e}_1 =: e'_2.\end{aligned}$$

20. Exercises

Next, let us choose images for h_1, h_2 . We want $[h_i, e_j] = [\psi(h_i), \psi(e_j)]$ and by a quick calculation, we find

$$\begin{aligned}\psi(h_1) &= \text{diag}(1/2, 1/2, -1/2, -1/2) =: h'_1, \\ \psi(h_2) &= \text{diag}(1/2, -1/2, 1/2, -1/2) =: h'_2.\end{aligned}$$

Finally, we find images for f_1, f_2 . These again need to span the corresponding root spaces and need to be chosen in such a way that we get $[e_i, f_j] = [\psi(e_i), \psi(f_j)]$. Another round of calculations leads us to

$$\begin{aligned}\psi(f_1) &= \tilde{f}_3 =: f'_1, \\ \psi(f_2) &= \frac{\tilde{f}_1}{2} =: f'_2.\end{aligned}$$

If we furthermore observe that $h'_1, h'_2, e'_1, e'_2, f'_1, f'_2$ span the Lie algebra \mathfrak{g}' , we conclude that ψ is an isomorphism of Lie algebras as desired. As such it induces through exponentiating a local isomorphism of the Lie groups $G = \text{SO}(2, 3)$ and $G' = \text{Sp}(4, \mathbb{R})$.

Θ -positive structures on G'

Finally, let us consider Θ -positive structures on G' . Due to the local isomorphism with G , which we just described, we expect to find two such structures. Indeed, G' is both split real and of Hermitian type of tube type and by the above also locally isomorphic to $\text{SO}(2, 3)$, which are three cases for which a Θ -positive structure can be found but it is not hard to see that the last two give the same structure in this case.

We will not investigate the totally positive structure on G' , instead let us focus on the other Θ -positive structure on G' , which is obtained by picking $\Theta' = \{\alpha'_1\}$. Using the Lie algebra isomorphism $\psi : \mathfrak{g} \rightarrow \mathfrak{g}'$, we can basically copy the above formulas (for G , in our notation we just need to add a lot of 's) and find

$$\begin{aligned}\mathfrak{u}'_{\Theta'} &:= \sum_{\alpha \in \Sigma'^+_{\Theta'}} \mathfrak{g}'_{\alpha} = \mathfrak{g}'_{\alpha'_1} \oplus \mathfrak{g}'_{\alpha'_1 + \alpha'_2} \oplus \mathfrak{g}'_{\alpha'_1 + 2\alpha'_2} = \mathbb{R}e'_1 \oplus \mathbb{R}e'_3 \oplus \mathbb{R}e'_4, \\ \mathfrak{u}'^{\text{opp}}_{\Theta'} &:= \sum_{\alpha \in \Sigma'^+_{\Theta'}} \mathfrak{g}'_{-\alpha} = \mathfrak{g}'_{-\alpha'_1} \oplus \mathfrak{g}'_{-\alpha'_1 - \alpha'_2} \oplus \mathfrak{g}'_{-\alpha'_1 - 2\alpha'_2} = \mathbb{R}f'_1 \oplus \mathbb{R}f'_3 \oplus \mathbb{R}f'_4, \\ \mathfrak{l}'_{\Theta'} &:= \mathfrak{g}'_0 \oplus \sum_{\alpha \in \text{Span}(\Delta' - \Theta') \cap \Sigma'^+} (\mathfrak{g}'_{\alpha} \oplus \mathfrak{g}'_{-\alpha}) = \mathfrak{h}' \oplus \mathfrak{g}'_{\alpha'_2} \oplus \mathfrak{g}'_{-\alpha'_2}.\end{aligned}$$

The center $\mathfrak{z}'_{\Theta'}$ of $\mathfrak{l}'_{\Theta'}$ can simply be read off or obtained using ψ and we find

$$\mathfrak{z}'_{\Theta'} = \{\text{diag}(\lambda, \lambda, -\lambda, -\lambda) : \lambda \in \mathbb{R}\}$$

and thus $\mathfrak{u}'_{\Theta'} = \mathfrak{u}'_{\alpha'_1}$ again only consists of a single weight space.

By exponentiating an element in $\mathfrak{l}'_{\Theta'}$, we see that a general element $X \in L'_{\Theta'}{}^{\circ}$ has the form

$$X = \begin{pmatrix} A & 0 \\ 0 & {}^t A^{-1} \end{pmatrix}$$

for $A \in \text{GL}(2, \mathbb{R})^{\circ}$. Now $Y \in \mathfrak{u}'_{\alpha'_1}$ is of the form

$$Y = \begin{pmatrix} 0 & M \\ 0 & 0 \end{pmatrix},$$

for some symmetric matrix M . Therefore we identify this weight space with the space of symmetric matrices.

As we want to consider the action of L_{Θ}° on $\mathfrak{u}'_{\alpha'_1}$ by conjugation, we calculate

$$XYX^{-1} = \begin{pmatrix} 0 & AM^tA \\ 0 & 0 \end{pmatrix}.$$

Thus we see that the cone c' of positive (semi-)definite matrices in $\mathfrak{u}'_{\alpha'_1}$ is preserved by the action of L_{Θ}° .

From here the computation of the Θ -positive semigroup is analogous to the one for G .

Bibliography

- [BCGP⁺21] Steve Bradlow, Brian Collier, Oscar Garcia-Prada, Peter Gothen, and André Oliveira, *A general Cayley correspondence and higher Teichmüller spaces*, 2021.
- [BD14] Francis Bonahon and Guillaume Dreyer, *Parameterizing Hitchin components*, *Duke Math. J.* **163** (2014), no. 15, 2935–2975. MR 3285861
- [BD17] ———, *Hitchin characters and geodesic laminations*, *Acta Math.* **218** (2017), no. 2, 201–295. MR 3733100
- [Ben00] Yves Benoist, *Automorphismes des cônes convexes*, *Invent. Math.* **141** (2000), no. 1, 149–193. MR 1767272
- [BIW07] Marc Burger, Alessandra Iozzi, and Anna Wienhard, *Hermitian symmetric spaces and Kähler rigidity*, *Transform. Groups* **12** (2007), no. 1, 5–32. MR 2308026
- [BIW10] ———, *Surface group representations with maximal Toledo invariant*, *Ann. of Math. (2)* **172** (2010), no. 1, 517–566. MR 2680425
- [BIW14] ———, *Higher Teichmüller spaces: from $SL(2, \mathbb{R})$ to other Lie groups*, *Handbook of Teichmüller theory. Vol. IV, IRMA Lect. Math. Theor. Phys.*, vol. 19, Eur. Math. Soc., Zürich, 2014, pp. 539–618. MR 3289711
- [Bor91] Armand Borel, *Linear algebraic groups*, second ed., *Graduate Texts in Mathematics*, vol. 126, Springer-Verlag, New York, 1991. MR 1102012
- [Bou75] N. Bourbaki, *Éléments de mathématique. Fasc. XXXVIII: Groupes et algèbres de Lie. Chapitre VII: Sous-algèbres de Cartan, éléments réguliers. Chapitre VIII: Algèbres de Lie semi-simples déployées*, Hermann, Paris, 1975, *Actualités Sci. Indust.*, No. 1364. MR 0453824
- [BP21] Jonas Beyrer and Beatrice Pozzetti, *A collar lemma for partially hyperconvex surface group representations*, *Trans. Amer. Math. Soc.* **374** (2021), no. 10, 6927–6961. MR 4315593
- [Bum13] Daniel Bump, *Lie groups*, second ed., *Graduate Texts in Mathematics*, vol. 225, Springer, New York, 2013. MR 3136522
- [CG93] Suhyoung Choi and William M. Goldman, *Convex real projective structures on closed surfaces are closed*, *Proc. Amer. Math. Soc.* **118** (1993), no. 2, 657–661. MR 1145415
- [CG97] ———, *The classification of real projective structures on compact surfaces*, *Bull. Amer. Math. Soc. (N.S.)* **34** (1997), no. 2, 161–171. MR 1414974
- [Cle04] Jean-Louis Clerc, *L'indice de Maslov généralisé*, *J. Math. Pures Appl. (9)* **83** (2004), no. 1, 99–114. MR 2032583
- [CØ01] J. L. Clerc and B. Ørsted, *The Maslov index revisited*, *Transform. Groups* **6** (2001), no. 4, 303–320. MR 1870049

- [Ebe96] Patrick B. Eberlein, *Geometry of nonpositively curved manifolds*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1996. MR 1441541
- [FG06] Vladimir Fock and Alexander Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. (2006), no. 103, 1–211. MR 2233852
- [FK94] Jacques Faraut and Adam Korányi, *Analysis on symmetric cones*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1994, Oxford Science Publications. MR 1446489
- [GLW21] Olivier Guichard, François Labourie, and Anna Wienhard, *Positivity and representations of surface groups*, 2021.
- [Gol88] William M. Goldman, *Topological components of spaces of representations*, Invent. Math. **93** (1988), no. 3, 557–607. MR 952283
- [Gui08] Olivier Guichard, *Composantes de Hitchin et représentations hyperconvexes de groupes de surface*, J. Differential Geom. **80** (2008), no. 3, 391–431. MR 2472478
- [GW12] Olivier Guichard and Anna Wienhard, *Anosov representations: domains of discontinuity and applications*, Invent. Math. **190** (2012), no. 2, 357–438. MR 2981818
- [GW18] ———, *Positivity and higher Teichmüller theory*, European Congress of Mathematics, Eur. Math. Soc., Zürich, 2018, pp. 289–310. MR 3887772
- [GW22] ———, *Generalizing Lusztig’s total positivity*, 2022.
- [Hel01] Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics, vol. 34, American Mathematical Society, Providence, RI, 2001, Corrected reprint of the 1978 original. MR 1834454
- [Hit92] N. J. Hitchin, *Lie groups and Teichmüller space*, Topology **31** (1992), no. 3, 449–473. MR 1174252
- [Hum72] James E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics, Vol. 9, Springer-Verlag, New York-Berlin, 1972. MR 0323842
- [Kna02] Anthony W. Knaapp, *Lie groups beyond an introduction*, second ed., Progress in Mathematics, vol. 140, Birkhäuser Boston, Inc., Boston, MA, 2002. MR 1920389
- [Kos10] Bertram Kostant, *Root systems for Levi factors and Borel-de Siebenthal theory*, Symmetry and spaces, Progr. Math., vol. 278, Birkhäuser Boston, Boston, MA, 2010, pp. 129–152. MR 2562626
- [KW65] Adam Korányi and Joseph A. Wolf, *Realization of hermitian symmetric spaces as generalized half-planes*, Ann. of Math. (2) **81** (1965), 265–288. MR 174787
- [Lab06] François Labourie, *Anosov flows, surface groups and curves in projective space*, Invent. Math. **165** (2006), no. 1, 51–114. MR 2221137
- [LT20] François Labourie and Jérémy Toulisse, *Quasicircles and quasiperiodic surfaces in pseudo-hyperbolic spaces*, 2020.
- [Lus94] G. Lusztig, *Total positivity in reductive groups*, Lie theory and geometry, Progr. Math., vol. 123, Birkhäuser Boston, Boston, MA, 1994, pp. 531–568. MR 1327548

Bibliography

- [LV80] Gérard Lion and Michèle Vergne, *The Weil representation, Maslov index and theta series*, Progress in Mathematics, vol. 6, Birkhäuser, Boston, Mass., 1980. MR 573448
- [OV94] A. L. Onishchik and È. B. Vinberg (eds.), *Lie groups and Lie algebras, III*, Encyclopaedia of Mathematical Sciences, vol. 41, Springer-Verlag, Berlin, 1994.
- [PS69] I. I. Pyateskii-Shapiro, *Automorphic functions and the geometry of classical domains*, Mathematics and its Applications, Vol. 8, Gordon and Breach Science Publishers, New York-London-Paris, 1969, Translated from the Russian. MR 0252690
- [Sch30] Isac Schoenberg, *Über variationsvermindernde lineare Transformationen*, Math. Z. **32** (1930), no. 1, 321–328. MR 1545169
- [Whi52] A. M. Whitney, *A reduction theorem for totally positive matrices*, J. Analyse Math. **2** (1952), 88–92. MR 53173
- [Wie04] Anna Wienhard, *Bounded cohomology and geometry*, Bonner Mathematische Schriften [Bonn Mathematical Publications], vol. 368, Universität Bonn, Mathematisches Institut, Bonn, 2004, Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 2004. MR 2205508
- [Wie18] ———, *An invitation to higher Teichmüller theory*, Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures, World Sci. Publ., Hackensack, NJ, 2018, pp. 1013–1039. MR 3966798
- [WK65] Joseph A. Wolf and Adam Korányi, *Generalized Cayley transformations of bounded symmetric domains*, Amer. J. Math. **87** (1965), 899–939. MR 192002